

RATIONAL CURVES AND STRICTLY NEF DIVISORS
ON CALABI–YAU THREEFOLDS

HAIDONG LIU AND ROBERTO SVALDI

Received: September 13, 2021

Revised: September 8, 2022

Communicated by Mihai Păun

ABSTRACT. We give a criterion for a nef divisor D to be semi-ample on a Calabi–Yau threefold X when $D^3 = 0 = c_2(X) \cdot D$ and $c_3(X) \neq 0$. As a direct consequence, we show that on such a variety X , if D is strictly nef and $\nu(D) \neq 1$, then D is ample; we also show that if there exists a Caritser divisor $D \neq 0$ in the boundary of the nef cone of X , then X contains a rational curve when its topological Euler characteristic is not 0.

2020 Mathematics Subject Classification: Primary 14J32; Secondary 14E30, 14J30

Keywords and Phrases: rational curves, strictly nef divisors, Calabi–Yau threefolds

1 INTRODUCTION

From the point of view of the birational classification of algebraic varieties, varieties with trivial canonical bundle and mild singularities (in short, \mathbb{K} -trivial varieties) are one of the fundamental building blocks. Hence, their study is central to the understanding of the structure of algebraic varieties as a whole. By a well-known result of Beauville and Bogomolov [Bea83], every smooth variety with trivial canonical bundle can be decomposed, after a finite étale cover, as a product of abelian, hyperkähler and Calabi–Yau manifolds. For the purpose of this note, a smooth projective variety X with trivial canonical bundle is Calabi–Yau if it is simply connected and $H^i(X, \mathcal{O}_X) = 0$ for $0 < i < \dim X$.

One of the central problems in modern birational geometry is the so-called Abundance Conjecture, see [Kol13, §8.7].

CONJECTURE 1.1. *Let (X, Δ) be a log canonical pair, where Δ is a \mathbb{Q} -divisor. If $K_X + \Delta$ is nef, then it is semi-ample.*

For K-trivial varieties, the above conjecture is expected to hold in even greater generality, cf. [Kol15, Conjecture 51], [LOP18, Section 4] and [LP20] for conventions and a survey of several related conjectures.

CONJECTURE 1.2 (Semi-ampleness conjecture for K-trivial varieties). *Let X be a projective klt variety with $K_X \equiv 0$. Assume that $H^1(X, \mathcal{O}_X) = 0$. Let D be a nef divisor on X . Then D is semi-ample.*

The major difference between Conjecture 1.1 for a K-trivial variety and Conjecture 1.2 is that the divisor D , while nef, is not required to be effective. The assumption that $H^1(X, \mathcal{O}_X) = 0$ implies that $\text{Pic}^0(X)$ is 0-dimensional, which is a necessary condition, as otherwise there would be numerically trivial line bundles that are non-torsion, although one can formulate a version of the conjecture where semi-ampleness of D holds up to numerical equivalence, cf. [LP20]. The condition is automatically satisfied for Calabi–Yau manifolds and simple hyperkähler manifolds. In the realm of simple hyperkähler manifolds, Conjecture 1.2 is equivalent to the so-called Strominger–Yau–Zaslow (SYZ) conjecture, [Ver10, Conjecture 1.7].

While the Conjecture 1.1 is known to hold in a number of cases, e.g., it holds for projective K-trivial threefolds, so that if D is an effective nef divisor then D is semi-ample, cf. [KMM94, Corollary, p.100] or [Ogu93], Conjecture 1.2 is only known to hold in full generality for K-trivial varieties of dimension at most 2. In dimension 3 and higher, only very few cases of the conjecture have been verified, cf. [Kol15, LOP18].

In this paper, we study some cases of Conjecture 1.2 together with some applications to other problems on Calabi–Yau threefolds.

The first result that we present is the following theorem quantifying precisely the possible failure of the Conjecture 1.2 for a nef divisor of numerical dimension 2 on a Calabi–Yau threefold. We recall, for the reader, that a prime divisor S is said to be *orthogonal* to a Cartier divisor D if $D|_S \equiv 0$; the *numerical dimension* $\nu(D)$ of a nef divisor D is $\nu(D) := \max\{h \in \mathbb{N} \mid D^h \neq 0\}$.

THEOREM 1.3 (=Theorem 4.2). *Let X be a Calabi–Yau threefold and D a nef divisor on X with $\nu(D) = 2$. Let S_1, \dots, S_r be all the prime divisors on X orthogonal to D . Let g_j be the irregularity of a resolution of S_j for each j . If*

$$\frac{c_3(X)}{2} \neq r - \sum_{j=1}^r g_j,$$

then D is semi-ample.

Theorem 1.3 is a generalization of [LOP20, Theorem 8.5]. The important improvement in this generalization is that we drop the positivity conditions [LOP20, Theorem 8.5(i)-(ii)] that the authors were using to be able to carry out certain cohomological computations. Instead, we obtain an unconditional statement, by showing that those conditions are not needed, via a

Fujita-type vanishing trick, cf. § 3-4. The ideas and statements that appear here and in [LOP20] are influenced by early work of Wilson on this topic, cf. [Wil92, Wil94a].

Conjecture 1.2 is strictly intertwined with the existence of rational curves on K-trivial varieties, a classically hard problem particularly on Calabi–Yau varieties. This link is highlighted by the following conjecture due to Oguiso.

CONJECTURE 1.4 ([Ogu93, Conjecture, p.456]). *Let X be a Calabi–Yau threefold. Assume that there exists a non-trivial Cartier divisor D contained in the boundary of $\text{Nef}(X)$. Then X contains a rational curve.*

Indeed, it is not hard to show that if Conjecture 1.2 holds then also Oguiso’s conjecture holds, as the Iitaka fibration of the divisor D will contain rational curves in some fibers, cf. Lemma 2.5. By applying the techniques used for Theorem 1.3, we are able to prove the following almost complete solution of Conjecture 1.4, which also complements the main result of [DF14].

THEOREM 1.5 (=Theorem 6.3). *Let X be a Calabi–Yau threefold. Assume that there exists a non-trivial Cartier divisor D contained in the boundary of $\text{Nef}(X)$. If either $c_3(X) \neq 0$ or the second Betti number $b_2(X) \notin \{2, 3, 4\}$, then X contains a rational curve.*

In Section 7, we show also how the problem of the existence of rational curves on Calabi–Yau threefolds is strictly intertwined with another important conjecture regarding the birational structure and the mirror symmetry of Calabi–Yau threefolds, the Kawamata–Morrison Conjecture. In this regard, we discuss also how geometric and cohomological considerations inspired by the conjecture could be used towards a definitive solution of Oguiso’s conjecture. A different point of view on how to connect hyperbolicity problems with the Mirror Symmetry viewpoint on Calabi–Yau threefolds can be found in [KV20].

Conjecture 1.2 is also intertwined with another conjecture due to Serrano [Ser95]. A *strictly nef* divisor D on a normal variety X is a nef Cartier divisor such that $D \cdot C > 0$ for any curve C on X . In [Ser95], Serrano conjectured that by twisting a sufficiently large multiple of a strictly nef divisor with the canonical bundle, one should obtain an ample divisor. On a Calabi–Yau manifold, as the canonical bundle is trivial, Serrano’s conjecture just predicts that strictly nef divisors should be ample, in agreement with Conjecture 1.2. Here, we show an instance in which this principle holds on Calabi–Yau threefolds.

THEOREM 1.6 (=Theorem 5.2). *Let X be a Calabi–Yau threefold with $c_3(X) \neq 0$. Let D be a strictly nef divisor on X with $\nu(D) \neq 1$. Then D is ample.*

The two results above are based on the solution of the following special case of Conjecture 1.2 for Calabi–Yau threefolds, which can be seen as a corollary of Theorem 1.3.

THEOREM 1.7. *Let X be a Calabi–Yau threefold with $c_3(X) \neq 0$. Let D be a nef divisor with $D^3 = 0 = c_2(X) \cdot D$. Assume that there exists a very ample line bundle H on X and a general member $G \in |H|$ of the linear system such that $D|_G$ is ample. Then D is semi-ample.*

It is not hard to see that if the conditions $D^3 = 0 = c_2(X) \cdot D$ are not satisfied, then D is semi-ample, cf. Lemma 2.2. Moreover, those implies that numerical dimension of D is 2, i.e., $\nu(D) = 2$. A weaker statement had been obtained in [LOP20, Theorem 1.2]: in their statement, the authors used a strong positivity condition on a certain sheaf of 1-forms with poles along an ample divisor and coefficients in D . Hence, we successfully remove their hypothesis from the statement to obtain a more general result that allows us to obtain Theorems 1.5 and 1.6.

The main idea in the proof of Theorem 1.7 is to use the positivity condition on the restriction of D to an element $G \in |H|$ together with Fujita’s vanishing theorem 3.1 and the Hirzebruch–Riemann–Roch formula to produce sections of the nef divisor D .

Finally, if a Calabi–Yau threefold X satisfies $c_3(X) = e(X) = 2(h^{1,1}(X) - h^{2,1}(X)) = 0$, then such a X would lie exactly on the axis of symmetry of the famous “Zoo of Calabi–Yau threefolds”, cf. [KS00, Figure 1]. It is a folklore conjecture, based on the knowledge of existing models acquired over the past 40 years, that all Hodge numbers of Calabi–Yau threefolds should lie inside the cup shape appearing in [KS00, Figure 1]. One of the first sources for this cornucopia of examples to be explored were smooth divisors in the anti-canonical linear system of a 4-dimensional \mathbb{Q} -factorial Gorenstein toric Fano variety with terminal singularities. By Batyrev’s construction in [Bat94], this class of toric Fano varieties are completely determined by reflexive polytopes in dimension 4; Kreuzer and Skarke in [KS00] listed all elements in this class of polytopes, by means of a computer-aided classification. As a result, we now know that there are 30,108 distinct Hodge diamonds for the 473.8 million families of Calabi–Yau threefolds. Appearing in this list, there are just 136 pairs of Hodge numbers of Calabi–Yau threefolds X with $h^{1,1}(X) = h^{2,1}(X)$, hence, $c_3(X) = 0$. Thus, the condition that $c_3(X) = 0$ appears relatively rarely distributed within our current picture of the “Zoo of Calabi–Yau threefolds”.

ACKNOWLEDGMENTS

The authors wish to thank Vladimir Lazić for encouragement and useful conversations, and for providing a lot of suggestions after reading a first draft of this paper. HL would like to thank Chen Jiang for helpful discussion and suggestions. Part of this project started as an appendix to [DFM19]. RS is very grateful to Simone Diverio, Claudio Fontanari, and Diletta Martinelli for their encouragement and support as well as useful discussion and suggestions on the structure and the content of the material here included.

Part of this work was carried out while RS was visiting Professor Jacopo Stoppa at SISSA, Trieste. He would like to thank Professor Stoppa and SISSA for

providing such a congenial place to work. The visit was supported by funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant agreement no. 307119. The author was partially supported by Churchill College, Cambridge and from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 842071.

2 PRELIMINARIES

Throughout this paper, we work over the complex numbers field \mathbb{C} . A *scheme* is always assumed to be separated and of finite type over \mathbb{C} . A *variety* is a reduced and irreducible scheme. For notation and conventions, we refer to [KM98] and [Laz04a, Laz04b].

As mentioned in the introduction, we define the *numerical dimension* $\nu(D)$ of a nef \mathbb{Q} -divisor D as

$$\nu(D) := \max\{h \in \mathbb{N} \mid D^h \neq 0\}.$$

It is well-known that $\nu(D) \geq 0$, $\kappa(D) \leq \nu(D) \leq \dim X$ and $\nu(D) = \dim X$ if and only if D is big, see, for example, [Nak04]. By Kleiman's Ampleness Criterion, it is also well-known that

LEMMA 2.1. *If $\nu(D) = \dim X - 1$, then $D^{\dim X - 1} \cdot H > 0$ for any ample divisor H on X .*

2.1 CONES OF DIVISORS

We adopt the following notation to denote the various cones of divisors on a variety X :

- $N^1(X)_{\mathbb{R}}$ denotes the Néron–Severi \mathbb{R} -vector space on X , while $N_1(X)_{\mathbb{R}}$ denotes its dual, that is, the \mathbb{R} -vector space of 1-cycle classes modulo numerical equivalences.
- The *effective cone* $\text{Eff}(X) \subset N^1(X)_{\mathbb{R}}$ is the convex cone spanned by the classes of all effective \mathbb{R} -divisor classes on X .
- The *pseudo-effective cone* $\overline{\text{Eff}}(X) \subset N^1(X)_{\mathbb{R}}$ is the closure of $\text{Eff}(X)$. The interior of $\overline{\text{Eff}}(X)$ is the convex cone of all big \mathbb{R} -divisor classes.
- The *nef cone* $\text{Nef}(X) \subset N^1(X)_{\mathbb{R}}$ is the convex cone of all nef \mathbb{R} -divisor classes on X . The interior of $\text{Nef}(X)$ is the convex cone of all ample \mathbb{R} -divisor classes.
- The *effective nef cone* $\text{Nef}^e(X)$ is equal to $\text{Nef}(X) \cap \text{Eff}(X)$. It is not necessarily closed.
- The *cone of effective curves* $\overline{\text{NE}}(X) \subset N_1(X)_{\mathbb{R}}$ is the closure of the cone spanned by effective 1-cycles with coefficients in \mathbb{R} .

It is well-known that $\text{Nef}(X) \subset \overline{\text{Eff}}(X)$ and $\text{Nef}(X)$ is the dual cone of $\overline{\text{NE}}(X)$, see [Laz04a, Chapter 1].

2.2 CALABI–YAU THREEFOLDS

We collect here a few basic results on Calabi–Yau threefolds that will be used in the rest of the paper, cf. [LOP20]. In this subsection, X will denote a Calabi–Yau threefold.

In this paper, we will use the Hirzebruch–Riemann–Roch formula on a Calabi–Yau threefold X . If \mathcal{F} is a vector bundle on X , then the Hirzebruch–Riemann–Roch formula is as follows:

$$\chi(X, \mathcal{F}) = \frac{1}{12}c_1(\mathcal{F})c_2(X) + \frac{1}{6}(c_1(\mathcal{F})^3 - 3c_1(\mathcal{F})c_2(\mathcal{F}) + 3c_3(\mathcal{F})). \quad (2.1)$$

In particular, for a Cartier divisor D on X , (2.1) can be simplified to give

$$\chi(X, D) = \frac{D^3}{6} + \frac{D \cdot c_2(X)}{12}. \quad (2.2)$$

It is well-known that on X the second Chern class $c_2(X)$ has non-negative intersection with nef divisors by the Miyaoka–Yau inequality [Miy87]; moreover, $c_2(X) \neq 0$, as otherwise X would be dominated by an abelian threefold, cf. [GKP16].

LEMMA 2.2. *Let X be a Calabi–Yau threefold. Let D be a nef Cartier divisor on X . Then the following hold:*

1. D is semi-ample if and only if $\kappa(D) \geq 0$;
2. if $\nu(D) = 2$ and $c_2(X) \cdot D > 0$, then D is semi-ample;
3. if $\nu(D) = 1$ and $c_2(X) \cdot D > 0$, then D is semi-ample;

Proof. 1. This follows from the solution of Conjecture 1.1 for threefolds [KMM94, KMM04].

2. Using the Kawamata–Viehweg vanishing theorem and its generalization [Laz04a, Example 4.3.7], if $\nu(D) = 2$ and $c_2(X) \cdot D > 0$, then $\kappa(D) \geq 0$ by (2.2). The conclusion follows from (1).

3. This is [Ogu93, Main Theorem].

□

REMARK 2.3. In view of Lemma 2.2, the only case left to consider to prove Conjecture 1.2 for Calabi–Yau threefolds is when $\kappa(D) = -\infty$. As D is nef, this immediately implies that we can assume $D^3 = 0$.

In the proof of Theorems 1.7 and 1.3, we will use the following result taken from [Wil94a, Proposition 3.1], [LOP20, Proposition 3.4]. A generalization of this result to higher dimensions appears in [LP18, Theorem 8.1].

PROPOSITION 2.4. *Let X be a Calabi–Yau threefold. Let D be a nef divisor on X with $\kappa(D) = -\infty$. Then there exists a positive integer m_0 such that*

$$H^0(X, \Omega_X^q \otimes \mathcal{O}_X(mD)) = 0$$

for all $|m| \geq m_0$ and $q \geq 0$.

The following two simple observations will play an important role in our discussion of rational curves on a Calabi–Yau varieties. These results are immediate consequences of standard birational geometry results.

LEMMA 2.5. *Let X be a Calabi–Yau threefold. Let D be a semi-ample but not ample divisor on X . Then, X contains a rational curve.*

LEMMA 2.6. *Let X be a klt projective variety with $K_X \equiv 0$. Assume that X does not contain rational curves. Then, $\text{Nef}(X) = \overline{\text{Eff}}(X)$.*

In particular, Lemma 2.6 holds for a Calabi–Yau threefold.

2.3 NEF DIVISORS AND THEIR NULL LOCI

Let X be a normal projective variety and D a nef Cartier divisor. We study the set $\mathcal{C}_D := \{C \subset X \mid D \cdot C = 0, \dim C = 1, C \text{ is irreducible}\}$. We recall here the following standard general observation.

LEMMA 2.7. *Let X be a normal projective variety of dimension 3 and D a nef Cartier divisor on X with $\nu(D) = 2$. Assume that the set \mathcal{C}_D is at most countable. Then for any very ample divisor H on X and for a very general member $G \in |H|$, $D|_G$ is ample.*

Proof. The proof of [LOP20, Lemma 2.5(b)] applies in this context. □

COROLLARY 2.8. *Let X be a normal projective variety of dimension 3 and D a strictly nef divisor on X with $\nu(D) = 2$. Then for any very ample divisor H on X and for any very general member $G \in |H|$, $D|_G$ is ample.*

Proof. The result follows immediately from Lemma 2.7 as $\mathcal{C}_D = \emptyset$ since D is strictly nef. □

We will also use, particularly in § 6, the following result inspired by [LOP20, Theorem 2.7].

LEMMA 2.9. *Let X be a Calabi–Yau threefold and D a nef divisor on X with $\nu(D) = 2$. Assume that X does not contain rational curves. Then exactly one of the following conditions holds true:*

- 1) *through any point $x \in X$ there exists a curve C_x with $D \cdot C_x = 0$; in particular in this case D is semi-ample; or*
- 2) *the set \mathcal{C}_D is at most countable.*

Proof. By the existence and properness of the Hilbert scheme, cf. [Kol96, Chapter I], and the nefness of D , if through a general point $x \in X$ there exists an irreducible curve C_x with $D \cdot C_x = 0$, then the same conclusion holds for any point of X . In this case, D is semi-ample by [LOP20, Theorem 2.7(ii)].

Therefore, we can assume that through the general point of X there is no curve C such that $D \cdot C = 0$. By the structure of the Hilbert scheme, there are at most countably many schemes T_i , $i \in I$ whose general points parametrize irreducible curves C with $D \cdot C = 0$. If each T_i is 0-dimensional then property 2) in the statement of the lemma holds. Hence, we are left to consider the case that at least one of the T_i , say T_1 , contains a curve L such that curves parametrized by L span an irreducible surface S covered by curves parametrized by L . Then [LOP20, Theorem 2.7(i)] implies that $D \cdot S = 0$. On the other hand, we have the following claim.

CLAIM. The surface S is an ample divisor in X .

Proof of the claim. As X does not contain rational curves, $\text{Nef}(X) = \overline{\text{Eff}}(X)$, by Lemma 2.6. Therefore, S is an effective nef divisor on X and S is semi-ample by log abundance for threefolds. If S is not ample, then X must contain rational curves, cf. Lemma 2.5, a contradiction to the assumptions of the lemma. \square

Lemma 2.1 and the claim imply that $D^2 \cdot S > 0$, which provides the sought contradiction. \square

3 PROOF OF THEOREM 1.7

First let us recall Fujita's vanishing theorem from [Fuj83a, Theorem (1)] and [Fuj83b, (5.1) Theorem], see also [Laz04a, Theorem 1.4.35].

THEOREM 3.1 (Fujita's vanishing theorem). *Let X be a projective scheme. Let \mathcal{F} be a coherent sheaf and H an ample Cartier divisor on X . Then there exists a positive integer $t_0 = t_0(\mathcal{F}, H)$ such that*

$$H^i(X, \mathcal{F} \otimes \mathcal{O}_X(tH + D)) = 0$$

for all $i > 0$, $t \geq t_0$ and any nef Cartier divisor D on X .

We use Fujita's vanishing theorem to derive the following vanishing result for a general coherent sheaf, when we are in the presence of a nef divisor restricting to an ample divisor along a hyperplane section.

LEMMA 3.2. *Let X be a projective scheme. Let \mathcal{F} be a torsion-free coherent sheaf, H a very ample Cartier divisor and D a nef Cartier divisor on X . Assume that there exists a very general member $G \in |H|$ of the linear system such that $D|_G$ is ample. Then there exists a positive integer m_0 such that*

$$H^i(X, \mathcal{F} \otimes \mathcal{O}_X(mD)) = 0$$

for all $i \geq 2$ and $m \geq m_0$.

Proof. By Fujita’s vanishing theorem, we know that there exists $t_0 = t_0(\mathcal{F}, H)$ such that

$$H^i(X, \mathcal{F} \otimes \mathcal{O}_X(tH + mD)) = 0 \tag{3.1}$$

for all $i > 0$, $m \geq 0$ and $t \geq t_0$. Moreover, we can assume that $D|_{G'}$ is ample for a general $G' \in |t_0H|$. In fact, since $D|_G$ is ample, D is also ample when restricted to the non-reduced scheme associated to $t_0G \in |t_0H|$ by [Laz04a, Proposition 1.2.16]. Since being ample is an open property along the fibers of a proper morphism, see [Laz04a, Theorem 1.2.17], the claim follows. Let us consider the following short exact sequence

$$0 \rightarrow \mathcal{F} \otimes \mathcal{O}_X(mD) \rightarrow \mathcal{F} \otimes \mathcal{O}_X(mD + G') \rightarrow \mathcal{G} \otimes \mathcal{O}_{G'}((mD + G')|_{G'}) \rightarrow 0,$$

where $\mathcal{G} := \mathcal{F} / \mathcal{F} \otimes \mathcal{O}_X(-G')$. The exactness on the left follows from the torsion-freeness assumption for \mathcal{F} . The vanishing in (3.1) implies that

$$H^i(G', \mathcal{G} \otimes \mathcal{O}_{G'}((mD + G')|_{G'})) \simeq H^{i+1}(X, \mathcal{F} \otimes \mathcal{O}_X(mD))$$

for all $i \geq 1$ and $m \geq 0$. Since $D|_{G'}$ is ample, by Serre’s vanishing theorem there exists a positive integer m_0 such that

$$H^i(G', \mathcal{G} \otimes \mathcal{O}_{G'}((mD + G')|_{G'})) = 0$$

for all $i \geq 1$ and $m \geq m_0$, thus concluding the proof of the lemma. □

REMARK 3.3. When X is a Calabi–Yau manifold of dimension n and \mathcal{F} is its cotangent bundle, we can give a quick alternative proof of the vanishing result in Lemma 3.2 without referring to Fujita’s vanishing theorem.

In fact, let $\pi: Y := \mathbb{P}(\Omega_X^1) \rightarrow X$ be the projection morphism such that $\pi_*\mathcal{O}_Y(1) = \Omega_X^1$. Note that $\mathcal{O}_Y(1)$ is π -ample. Therefore, there exists a fixed integer t_1 such that $\mathcal{O}_Y(1) \otimes \mathcal{O}_Y(t\pi^*H)$ is ample for all $t \geq t_1$. Equivalently, for such t , $\mathcal{G}_t := \Omega_X^1 \otimes \mathcal{O}_X(tH)$ is ample. Hence,

$$H^i(X, \omega_X \otimes \mathcal{G}_t \otimes \det \mathcal{G}_t \otimes \mathcal{O}_X(D')) = 0 \tag{3.2}$$

for all $i \geq 1$ and any nef divisor D' by [LOP20, Lemma 2.11]. By assumption, $\det \Omega_X^1 \simeq \omega_X \simeq \mathcal{O}_X$. Therefore, $\det \mathcal{G}_t \simeq \det \Omega_X^1 \otimes \mathcal{O}_X(ntH) \simeq \mathcal{O}_X(ntH)$. Taking $\bar{t} = (n + 1)t_1$, it follows from (3.2) that

$$H^i(X, \Omega_X^1 \otimes \mathcal{O}_X(\bar{t}H + D')) = H^i(X, \omega_X \otimes \mathcal{G}_{t_1} \otimes \det \mathcal{G}_{t_1} \otimes \mathcal{O}_X(D')) = 0$$

for all $i \geq 1$ and any nef divisor D' . Taking $D' = D + (t - \bar{t})H$ then completes the argument.

Proof of Theorem 1.7. Let us assume by contradiction that D is not semi-ample. Then, by [LOP20, Proposition 2.2(iv)] and Serre duality, for all $s \in \mathbb{Z}$,

$$\chi(X, \Omega_X^q \otimes \mathcal{O}_X(sD)) = (-1)^q \frac{c_3(X)}{2}, \quad \text{for } q = 1, 2. \tag{3.3}$$

The assumption in the statement of the theorem implies that there exists a positive integer s_0 such that for any $s \geq s_0$

$$H^i(X, \Omega_X^q \otimes \mathcal{O}_X(sD)) = 0, \text{ for } i \geq 2, q = 1, 2, \quad (3.4)$$

by Lemma 3.2. Hence, Proposition 2.4 and (3.4) imply that for any $s \geq s_0$, we can rewrite (3.3) in the following ways

$$\begin{aligned} -\frac{c_3(X)}{2} &= \chi(X, \Omega_X^1 \otimes \mathcal{O}_X(sD)) = -h^1(X, \Omega_X^1 \otimes \mathcal{O}_X(sD)) \leq 0, \\ \frac{c_3(X)}{2} &= \chi(X, \Omega_X^2 \otimes \mathcal{O}_X(sD)) = -h^1(X, \Omega_X^2 \otimes \mathcal{O}_X(sD)) \leq 0. \end{aligned}$$

These two inequalities together imply that $c_3(X) = 0$, which gives the desired contradiction. \square

REMARK 3.4. Let X be a Calabi–Yau threefold and let \mathcal{F} be the cotangent bundle of X . The result of Theorem 1.7 was implicitly claimed by Wilson in the proof of [Wil94a, Theorem 2.3]. As the proof of Wilson’s claim is sketched loosely, Lazić–Oguiso–Peternell tried to reprove this result in [LOP20], but ended up needing to add the extra assumption (i) in [LOP20, Proposition 5.2] to obtain the desired statement. Our result shows that this assumption is indeed not necessary.

4 PROOF OF THEOREM 1.3

In this section, we aim to prove Theorem 1.3. The variety X will be a Calabi–Yau threefold and D will be a nef divisor of numerical dimension 2 on X . Let us recall that a prime divisor $S \subset X$ is orthogonal to D if $S \cdot D = 0$, or, equivalently, $D|_S \equiv 0$.

A birational morphism $\phi: X \rightarrow Z$ is said to be a *good Calabi–Yau model* for a nef divisor D provided that, cf. [LOP20, Definition 8.2],

- Z is a \mathbb{Q} -factorial K -trivial threefold with canonical singularities;
- the exceptional locus of ϕ contains all and only the divisors orthogonal to D ; and
- $D \sim \phi^* D_Z$, for some Cartier divisor M on Z .

In the following theorem, we recall the existence of good Calabi–Yau models of nef divisors, proven in [LOP20, Theorem 8.3], together with some of its immediate consequences.

THEOREM 4.1. *Let X be a Calabi–Yau threefold. Let D be a nef Cartier divisor on X and let S_1, \dots, S_r be the prime divisors orthogonal to D on X . Then, the following hold:*

1. there exists a smooth Calabi–Yau threefold X' and an isomorphism in codimension one $\alpha: X \dashrightarrow X'$ such that $D' := \alpha_*D$ is nef and X' admits a good Calabi–Yau model $\phi': X' \rightarrow Z$ for D' ;
2. the orthogonal divisors for D' on X' are given by the strict transforms S'_1, \dots, S'_r of S_1, \dots, S_r on X' . Moreover, for all $j \in \{1, \dots, r\}$ the irregularity g_j of a resolution of S_j coincide with the irregularity of a resolution of S'_j .
3. D and D' are crepant birational: if

$$\begin{array}{ccc}
 & Y & \\
 \swarrow p & & \searrow q \\
 X & \dashrightarrow \alpha \dashrightarrow & X'
 \end{array} \tag{4.1}$$

is any common projective resolution of indeterminacies for α , then $p^*D = q^*D'$;

4. D is semi-ample if and only if D' is semi-ample; and,
5. $c_3(X) = c_3(X')$.

Proof. 1. This is [LOP20, Theorem 8.3]. Let us recall that α is constructed $\alpha = \gamma^{-1} \circ \beta$, where $\beta: X \dashrightarrow X''$ is a minimal model for the $(K_X + \epsilon(\sum_{i=1}^r S_i))$ -MMP on X , for $0 < \epsilon \ll 1$ and $\gamma: X' \rightarrow X''$ is the terminalization of $(X'', 0)$ and $D' = \gamma^*\beta_*D$.

2. As at each step of the $(K_X + \epsilon(\sum_{i=1}^r S_i))$ -MMP constructing β , the curves contracted have intersection 0 with the strict transform of D , then $D = \beta^*\beta_*D$. Since $\beta_*\sum_{i=1}^r S_i = 0$, then X'' does not contain any orthogonal surfaces for D'' and the exceptional divisor of γ is given by the sum of the S'_i . As $D' = \gamma^*\beta_*D$, then the conclusion follows.

3. Since

$$D' = \alpha_*, \quad D' = \gamma^*\beta_*D, \quad \text{and} \quad D = \beta^*\beta_*D,$$

the conclusion follows immediately from the commutativity of the diagram (4.1)

4. This follows immediately from (3).
5. As X and X' are isomorphic in codimension one, their Hodge numbers coincide, cf. [Bat99]; in particular, $c_3(X) = c_3(X')$. □

In view of Theorem 4.1, in order to prove Theorem 1.3, we can substitute X (resp. D) with X' (resp. D') and assume that X is endowed with a good Calabi–Yau model $\phi: X \rightarrow Z$ for D . Thus, it suffices to prove the following theorem.

THEOREM 4.2. *Let X be a Calabi–Yau threefold and D a nef divisor on X with $\nu(D) = 2$, such that there exists a birational morphism $\phi: X \rightarrow Z$ which is a good Calabi–Yau model for D . Let S_1, \dots, S_r be all the prime divisors on X orthogonal to D . Let g_j be the irregularity of a resolution of S_j for each j . If*

$$\frac{c_3(X)}{2} \neq r - \sum_{j=1}^r g_j,$$

then D is semi-ample.

In order to prove Theorem 4.2, we will need the following generalization of [LOP20, Proposition 8.13]. Let us recall that since ϕ is a good Calabi–Yau model for D , we can assume, by definition of such model, that $D \sim \phi^* D_Z$.

LEMMA 4.3. *Notation as in Theorem 4.2. Fix a resolution $\tau: Y \rightarrow X$ such that the exceptional set of the morphism $\pi: Y \rightarrow Z$ where $\pi := \phi \circ \tau$, is a simple normal crossings divisor $E = \sum_{j=1}^s E_j$ on Y . Suppose that D is not semi-ample. Then*

$$H^q(Y, \Omega_Y^1(\log E) \otimes \pi^* \mathcal{O}_Z(mD_Z)) = 0$$

for any $q \geq 0$ and $m \gg 0$,

Proof. As showed in [LOP20, Proposition 8.13], it suffices to prove the following vanishings:

- (i) $H^1(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) = H^3(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) = 0$ for any $m \ll 0$,
- (ii) $\chi(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) = 0$ for all integers m ,
- (iii) $H^q(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) = 0$ for all q and all integers m such that $|m| \gg 0$, and
- (iv) $R^i \pi_* \Omega_Y^1(\log E) = 0$ for $i = 1, 2$.

We only sketch the proof of (i) and (ii), since the proof of the other two is exactly the same as that in [LOP20, Lemma 8.10 and Proposition 8.13] and does not use the extra hypotheses in the statement of the proposition. Let us recall that

$$\pi_* \Omega_Y^p(\log E) \simeq \Omega_Z^{[p]} \quad \text{and} \quad \phi_* \Omega_X^p \simeq \Omega_Z^{[p]} \quad (4.2)$$

for $0 \leq p \leq 3$, as in [LOP20, Equation (37)].

- (i) As $K_Z \sim 0$, by Serre duality, it follows that for any $m \in \mathbb{Z}$,

$$H^1(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) \simeq H^2(Z, \Omega_Z^{[2]} \otimes \mathcal{O}_Z(-mD_Z)).$$

Proposition 4.4 and Lemma 3.2 imply that

$$H^2(Z, \Omega_Z^{[2]} \otimes \mathcal{O}_Z(mD_Z)) = 0, \quad \forall m \gg 0. \tag{4.3}$$

Hence, the first part of (i) follows by combining these two observations. The second vanishing claimed in (i) follows instead from the following isomorphisms

$$\begin{aligned} & H^3(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) \\ & \simeq \text{Hom}(\Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z), \mathcal{O}_Z) && \text{[by Serre duality]} \\ & \simeq H^0(Z, \Omega_Z^{[2]} \otimes \mathcal{O}_Z(-mD_Z)) && \text{[since } K_Z \sim 0\text{]} \\ & \simeq H^0(X, \Omega_X^2 \otimes \mathcal{O}_X(-mD)) = 0 && \text{[by (4.2)]} \\ & = 0. && \text{[by Proposition 2.4]} \end{aligned}$$

(ii) By the same proof as [LOP20, Proposition 8.13, (iii)], we obtain that

$$\chi(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) \text{ is independent of } m.$$

By Proposition 2.4 and (4.2), $H^0(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) = 0$, for all $|m| \gg 0$. Moreover, by Proposition 4.4 and Lemma 3.2, for all $m \gg 0$,

$$H^i(Z, \Omega_Z^{[j]} \otimes \mathcal{O}_Z(mD_Z)) = 0, \quad i = 2, 3, \quad j = 1, 2. \tag{4.4}$$

Hence, (4.4) and Serre duality, together with (i), imply that

$$\chi(Z, \Omega_Z^{[1]} \otimes \mathcal{O}_Z(mD_Z)) \begin{cases} \leq 0 & \text{for } m \gg 0, \\ \geq 0 & \text{for } m \ll 0, \end{cases}$$

which concludes the proof. □

Proof of Theorem 4.2. With the notation introduced above, the proof of [LOP20, Theorem 8.5] applies almost verbatim after replacing [LOP20, Theorem 8.13] with our Lemma 4.3. □

In the proof of Lemma 4.3, we used the following result.

PROPOSITION 4.4. *Let X be a Calabi–Yau threefold. Let D be a nef divisor of numerical dimension 2 on X which is not semi-ample. Assume that there exists a good Calabi–Yau model $\phi: X \rightarrow Z$ for D . Let D_Z be a \mathbb{Q} -Cartier divisor on Z such that $D \sim_{\mathbb{Q}} \phi^*D_Z$. Let H_Z be a very ample divisor on Z . Then for a very general $G \in |H_Z|$, $D_Z|_G$ is ample.*

Proof. We give a proof by contradiction.

Assume that for all $G \in |H_Z|$, $D_Z|_G$ is not ample. By Lemma 2.1, $D_Z^2 \cdot G > 0$. Hence, by the Nakai–Moishezon criterion, for any $G \in |H_Z|$ there exists a curve $C_G \subset G$ such that $D_Z \cdot C_G = 0$. In particular, as the Hilbert scheme of Z with respect to the polarization given by H_Z has countably many components, there exists a family \mathcal{C} of curves that have intersection 0 with D_Z on Z , such that the dimension of \mathcal{C} is ≥ 2 . Let

$$S_{\mathcal{C}} = \bigcup_{C \in \mathcal{C}} C \subset Z.$$

Then $\dim S_{\mathcal{C}} \geq 2$.

If $S_{\mathcal{C}} = Z$, we show that D is semi-ample, which prompts the desired contradiction. Indeed, by taking the strict transform C_X of the general curve $C \in \mathcal{C}$ on X , it follows that X itself is covered by curves having intersection 0 with D and we can apply then [LOP20, Theorem 2.7(ii)] to prove that D is semi-ample. The strict transform C_X of the general element $C \in \mathcal{C}$ exists, since \mathcal{C} covers Z and so C_X is not contained in the exceptional locus of ϕ . If $S_{\mathcal{C}} \subsetneq Z$, let S' be an irreducible component of $S_{\mathcal{C}}$. Then, it follows from [LOP20, Theorem 2.7(i)] that

$$S'_X \cdot D = 0, \quad \text{where } S'_X := \phi_*^{-1} S'. \quad (4.5)$$

This prompts the desired contradiction, since (4.5) implies that S'_X would have to be contracted by ϕ , by definition of a good Calabi–Yau model. \square

5 STRICTLY NEF DIVISORS

Let us recall that a strictly nef divisor D on a normal variety X is a nef Cartier divisor such that $D \cdot C > 0$ for any curve C on X . In general, a strictly nef divisor D is not necessarily ample, as shown by many classical example, e.g., the examples of Mumford and Ramanujam, see [Har70, appendix to Chapter I]. On the other hand, given a strictly nef divisor D on a projective variety X , the ampleness of D is implied by D being semi-ample, as then strict nefness of D forces the Iitaka fibration $\pi: X \rightarrow Z$ of D to be finite.

While, as we have just discussed, we cannot expect strictly nef divisors to always be ample, the situation is expected to improve if we allow twisting with the canonical bundle, as it is often the case when working with positivity properties in algebraic geometry. In this context, Serrano [Ser95] proposed the following conjecture that aims to bound the distance between ample and strictly nef divisors once we allow such twisting. This conjecture can be also thought as a weak analogue of Fujita’s Conjectures, cf. [Laz04a, § 10.4.A], for strictly nef divisors.

CONJECTURE 5.1. *Let X be a projective manifold of dimension n and D a strictly nef divisor on X . Then $K_X + tD$ is ample for any real number $t > n+1$.*

In [Ser95], Serrano showed that Conjecture 5.1 holds in dimension 2, and that in dimension 3 the only unknown cases are the following two:

- (i) X is a Calabi–Yau threefold and $c_2(X) \cdot D = 0$, and
- (ii) $\kappa(K_X) = -\infty$, and either the irregularity $q(X) \leq 1$ or else $q(X) = 2$ and $\chi(\mathcal{O}_X) = 0$.

A decade later, Campana–Chen–Peternell [CCP08] ruled out the latter case and proved some partial results for projective manifolds of dimension higher than 3. When X is a Calabi–Yau threefold and D is a strictly nef divisor, by Lemma 2.2, we can assume that $\kappa(D) = -\infty$ and $c_2(X) \cdot D = 0$, which in turn implies that $D^3 = 0$ and $1 \leq \nu(D) \leq 2$.

The results proven in the previous sections allow us to prove that Serrano’s conjecture holds on a Calabi–Yau threefold if we assume that the numerical dimension $\nu(D)$ of the strictly nef divisor D is $\neq 1$, when $c_3(X) \neq 0$.

THEOREM 5.2. *Let X be a Calabi–Yau threefold with $c_3(X) \neq 0$. Let D be a strictly nef divisor on X with $\nu(D) \neq 1$. Then D is ample.*

Proof. The result follows at once from Corollary 2.8 and Theorem 1.7. □

In the case $\nu(D) = 1$, instead, it is possible to prove the following result, valid for Calabi–Yau varieties of any dimension ≥ 3 , which is a special case of [LP18, Theorem 6.5].

THEOREM 5.3. *Let X be a Calabi–Yau manifold of dimension $n \geq 3$ and D a nef divisor on X with $\nu(D) = 1$. Assume that there is a singular hermitian metric h on $\mathcal{O}_X(D)$ with a semipositive curvature current such that the multiplier ideal sheaf $\mathcal{I}(h) \subsetneq \mathcal{O}_X$. Then $\kappa(D) \geq 0$.*

Note that for a pseudo-effective Cartier divisor D , thus including the case where D is nef, we can always find a singular hermitian metric h on $\mathcal{O}_X(D)$ with a semipositive curvature current, cf. [Dem92]. In general, it is rather hard to determine whether the multiplier ideal sheaf $\mathcal{I}(h)$ of the metric h is contained in \mathcal{O}_X strictly or not.

COROLLARY 5.4. *Let X be a Calabi–Yau threefold and D a strictly nef divisor on X . If there is a singular metric h on $\mathcal{O}_X(D)$ with semipositive curvature current such that $\mathcal{I}(h) \subsetneq \mathcal{O}_X$, then $\nu(D) \neq 1$.*

We conclude this section with a final remark on the case $\nu(D) = 1$.

REMARK 5.5. Let X be a Calabi–Yau threefold and D a strictly nef divisor on X with $\nu(D) = 1$. Then X cannot contain any del Pezzo, $K3$ or abelian surfaces.

Indeed, if $S \subset X$ is a del Pezzo surface, then $D|_S$ is semi-ample by the basepoint-free theorem. As $D|_S$ must also be strictly nef, $D|_S$ is ample which contradicts the assumption that $\nu(D) = 1$. It is well-known that, on abelian

and $K3$ surfaces strictly nef divisors are semi-ample, hence ample; thus, the same argument as in the del Pezzo case shows that such types of surfaces cannot be contained in X .

As a consequence of this simple observation, the existence of a strictly nef divisor D with $\nu(D) = 1$ implies that there cannot possibly exist morphisms $\phi: X \rightarrow Y$ such that:

- ϕ is a birational morphism contracting a del Pezzo surface to a point; or
- ϕ is a morphism whose generic fibre is an abelian or $K3$ surface.

In particular, there is no nef effective divisor E on X such that $\nu(E) = 1$: indeed, E would then be semi-ample and its Iitaka fibration would give a fibration in abelian/ $K3$ surfaces, cf. the proof of Lemma 2.5.

Conversely, if there is a contraction ϕ of X as above, then there is no strictly nef divisor of numerical dimension 1 on X . This type of observation will also be used in the analysis of the existence of rational curves in the next section.

6 EXISTENCE OF RATIONAL CURVES

Another classical problem in the study of K -trivial varieties is to determine whether or not they contain rational curves. Existence of rational curves is fully established only in dimension 2 thanks to Bogomolov and Mumford [MM83], while only partial results are known in dimension 3 and higher, as we discuss below. Even though it is widely believed that every Calabi–Yau manifold should contain rational curves, already in the threefold case this appears to be a very difficult problem.

In his study of the birational structure of Calabi–Yau threefolds, Oguiso proposed Conjecture 1.4 which is a weakened version of the classical question on the existence of rational curves. We recall it here for the reader’s convenience.

CONJECTURE 6.1. *Let X be a Calabi–Yau threefold. Assume that there exists a non-trivial Cartier divisor D contained in the boundary of $\text{Nef}(X)$. Then X contains a rational curve.*

The existence of a divisor D satisfying the hypotheses of Conjecture 6.1 is a rather strong assumption. First of all, it requires that $b_2(X) > 1$. It also implies that the boundary of the nef cone contains rational points other than the origin. The lack of rational points in the boundary of the nef cone is a rather typical situation if we do not assume that $\pi_1(X) = \{1\}$ or $h^i(X, \mathcal{O}_X) = 0$ for $0 < i < \dim X$. For example, the nef cone of an abelian variety contains non-zero rational points if and only if it admits a morphism to a lower dimensional abelian variety; in [Ogu14], Oguiso has constructed examples of hyperkähler manifolds and of Calabi–Yau threefolds of Picard number 2 whose nef cones do not contain rational points other than zero in the boundary. Nonetheless, abelian varieties do not contain rational curves, while Verbitsky proved in [Ver15] that hyperkähler manifolds are not Kobayashi hyperbolic provided

that their second Betti number is > 3 . Moreover, the analogous of Conjecture 6.1 is expected to hold also for hyperkähler manifolds.

In a fundamental series of works [Wil89, Wil92, Wil93], Wilson studied the structure of the nef cone of a Calabi–Yau threefold. In particular, he showed that if $b_2(X)$ is sufficiently big ($b_2(X) \geq 13$), then there always exists a birational morphism $f: X \rightarrow Y$. In particular, this forces the existence of a non-zero rational point in the boundary of the nef cone of X . Moreover, as already mentioned in Lemma 2.5, the exceptional locus of f is uniruled, so that rational curves exist in X . Thus, to solve Conjecture 6.1, we just need to focus on Calabi–Yau varieties of low Picard rank, that is, of small second Betti number. Building on the seminal works of Wilson, Diverio–Ferretti proved the following result in [DF14, Theorem 1.2] which partially solves Conjecture 6.1.

THEOREM 6.2. *Let X be a Calabi–Yau threefold. Assume that there exists a non-trivial Cartier divisor D contained in the boundary of $\text{Nef}(X)$ and $b_2(X) \geq 5$. Then X contains a rational curve.*

We briefly explain some of the arguments contained in the proof of the above theorem.

By contradiction, if there is no rational curve on X , we can assume that $\text{Nef}(X) = \overline{\text{Eff}}(X)$ and $D^3 = 0 = c_2(X) \cdot D$, cf. the proof of Lemma 2.5 and Lemma 2.6; thus, we can assume that the cohomology class of D lies on the intersection between the cubic hypersurface

$$W := \{T \in H^2(X, \mathbb{R}) \mid T^3 = 0\} \subset H^2(X, \mathbb{R}) \tag{6.1}$$

and the hyperplane

$$c_2(X)^\perp := \{H \in H^2(X, \mathbb{R}) \mid c_2(X) \cdot H = 0\}. \tag{6.2}$$

In this case, the Hirzebruch–Riemann–Roch formula in (2.2) implies that for all $m \in \mathbb{Z}$,

$$\chi(X, \mathcal{O}_X(mD)) = 0.$$

It is then not clear in general how to produce sections in any of the linear systems $|mD|$. Let $\widetilde{W} \subset \mathbb{P}(H^2(X, \mathbb{R}))$ be the projectivization of W . If $D^2 = 0$, then the point p_D associated to D in $\mathbb{P}(H^2(X, \mathbb{R}))$ is a singular point of \widetilde{W} . Hence, either \widetilde{W} is the union of a hyperplane and a conic both defined over \mathbb{Q} and their intersection contains the rational point p_D , or \widetilde{W} is irreducible. In the latter case, in turn, projecting from p_D to a hyperplane in $\mathbb{P}(H^2(X, \mathbb{R}))$, then either \widetilde{W} is a rational variety over \mathbb{Q} or \widetilde{W} is a cone over a cubic hypersurface in \mathbb{P}^{b_2-2} with vertex in p_D . This last case cannot happen, as a simple computation shows that it would imply that the numerical dimension of D is 0. As a consequence of the above discussion, \widetilde{W} contains a dense set of points defined over \mathbb{Q} . As $\text{Nef}(X) \subset W$, then X carries an elliptic fibration. By Lemma 2.5, X contains a rational curve.

When $\nu(D) = 2$, in [DF14], the authors carefully analyze the configuration of \widetilde{W} and $c_2(X)^\perp$, together with results of Wilson, to conclude the proof of Theorem 6.2.

We show that when $c_3(X) \neq 0$ it is actually possible to fully prove Oguiso's conjecture. We use the results of § 3 to show that under these assumptions the divisor D contained in the boundary of the nef cone will be semi-ample. Hence, by Lemma 2.5, X will then contain a rational curve.

THEOREM 6.3. *Let X be a Calabi–Yau threefold. Assume that there exists a non-trivial Cartier divisor D contained in the boundary of $\text{Nef}(X)$. Then, X contains a rational curve provided that either $c_3(X) \neq 0$ or the second Betti number $b_2(X) \neq 2, 3, 4$.*

Proof. By contradiction, we assume that X does not contain rational curves. We distinguish two cases.

CASE 1. *We assume that $c_3(X) \neq 0$.*

By the above discussion, we can assume that $c_2(X) \cdot D = D^3 = 0 \neq D^2$. If X does not contain any rational curve, then by Lemma 2.9 the set \mathcal{C}_D is at most countable. Hence, Lemma 2.7 and Theorem 1.7 imply that D is semi-ample, which prompts a contradiction by Lemma 2.5.

CASE 2. *We assume that $c_3(X) = 0$ and $b_2(X) \neq 2, 3, 4$.*

The result follows from Theorem 6.2. □

The material in the next section contains some considerations about possible strategies to prove the missing cases.

7 KAWAMATA–MORRISON CONJECTURE

In the final section we explain how Conjecture 6.1 is closely intertwined with another important conjecture regarding Calabi–Yau manifolds, the so-called Kawamata–Morrison Cone Conjecture, [Mor93, Kaw97].

CONJECTURE 7.1 (Kawamata–Morrison Cone Conjecture). *Let X be a Calabi–Yau manifold and $\text{Aut}(X)$ the group of automorphisms of X . There exists a rational polyhedral cone Π which is a fundamental domain for the action of $\text{Aut}(X)$ on $\text{Nef}^e(X)$, in the sense that*

- a) $\text{Nef}^e(X) = \bigcup_{g \in \text{Aut}(X)} g^*\Pi$; and
- b) $\text{Int } \Pi \cap \text{Int } g^*\Pi = \emptyset$ unless $g^* = \text{Id}_{N^1(X)}$.

Moreover, the number of $\text{Aut}(X)$ -equivalence classes of faces of the cone $\text{Nef}^e(X/Y)$ corresponding to birational contractions or fiber space structures is finite.

There is an analogous version of the conjecture involving the birational automorphism groups and its action on the cone of effective movable divisors

of X . There is now a fairly extensive literature on the Cone conjecture; for the case of Calabi–Yau threefolds that we treat here, the most relevant papers are [LP13, LOP18, FHS21].

The above conjecture predicts that if $\text{Aut}(X)$ is finite then the cone on which that group acts should be rational polyhedral. It follows that, if $\text{Nef}(X)$ is polyhedral, then those divisors contained in the facets of $\partial \text{Nef}(X)$ not contained in the hyperplane $c_2(X)^\perp$ are semi-ample (but not ample). In that case, X will contain rational curves, by Lemma 2.5. Moreover, assuming Conjecture 7.1 holds, Conjecture 6.1 holds in some of the cases that are not covered by Theorem 6.3, as we proceed to explain now.

We follow the notation introduced in the previous section.

PROPOSITION 7.2. *Let X be a Calabi–Yau threefold. Assume that Conjecture 7.1 holds true. Assume that $b_2(X) \neq 3, 4$ and that \widetilde{W} is not the union of the hyperplane $P = c_2(X)^\perp$ and a quadric Q of rank $b_2(X)$ which does not contain any rational points. Then Conjecture 6.1 holds.*

Proof. We may assume, by contradiction, that X does not contain rational curves and proceed to a case by case analysis. Thus, by Lemmata 2.5 and 2.6, $\text{Nef}(X) = \overline{\text{Eff}}(X)$, the boundary of $\text{Nef}(X)$ is contained in W and it does not contain any semi-ample divisor $D \not\sim 0$. Let D be a non-zero Cartier divisor contained in the boundary of $\text{Nef}(X)$; thus, $D^3 = 0 = c_2(X) \cdot D$. By abusing notation, we will use D to denote also the rational point in $\mathbb{P}(H^2(X, \mathbb{R}))$ associated to it, so that $D \in \widetilde{W}$.

CASE $b_2(X) = 2$. The hypersurface $W \subset H^2(X, \mathbb{R})$ is a union of three lines, l_1, l_2, l_3 ; one of the lines, say l_1 is generated by the class of D , as $D^3 = 0$; hence $c_2(X)^\perp = l_1$, while the other two lines, l_2, l_3 , are defined over a quadratic extension of \mathbb{Q} and are conjugated under the \mathbb{Z}_2 -action of Galois group. As the matrix M_g corresponding to the action of an element of $g \in \text{Aut}(X)$ on $N^1(X)_\mathbb{R}$ is defined over the integers, and l_1 is an eigenspace of eigenvalue 1, then M_g must be the identity as it has to preserve $\text{Nef}(X)$. Hence, Conjecture 7.1 implies that $\text{Nef}(X)$ is polyhedral, which then implies $\partial \text{Nef}(X) \not\subset W$.

CASE $b_2(X) = 3$. Then the possible configurations for the pair $(\widetilde{W}, c_2(X)^\perp)$, excluding the one in the statement of the proposition, are the following:

1. \widetilde{W} is the union of three distinct lines H_1, H_2, H_3 , while $c_2(X)^\perp$ is a line;
2. \widetilde{W} is the union of a line P and a conic C , and $c_2(X)^\perp$ is a line. If the rank of C is maximal, then C contains rational points;
3. \widetilde{W} is geometrically irreducible and $c_2(X)^\perp$ is a line L .

We will show that, in all these cases, the boundary of $\text{Nef}(X)$ must contain a semi-ample but not ample divisor. In case (1) either the three hyperplanes H_1, H_2, H_3 , are all defined over \mathbb{Q} and so the boundary of $\text{Nef}(X)$ contains a nef divisor E with $c_2(X) \cdot E > 0$; or, up to relabeling the H_i , H_1 is defined over \mathbb{Q} and H_2, H_3 are defined over a real quadratic extension of \mathbb{Q} and are conjugated

under the \mathbb{Z}_2 -action of the Galois group. In this case, either $H_1 \neq c_2(X)^\perp$, so that, again, H_1 will contain a nef and semi-ample divisor, or the point $p_F = H_2 \cap H_3$ is defined over \mathbb{Q} and it corresponds to projectivization of a nef divisor F with $c_2(X) \cdot F > 0$, and again we can conclude as above.

In case (2), the boundary of the nef cone is contained in \widetilde{W} , hence either P is tangent to C and the nef cone is contained in the oval given by the interior of C , or the nef cone is contained in one of the subsets in which the line P partitions the oval given by the interior of C . In the former case, then $P \cap C$ is a rational point, hence C is rational over \mathbb{Q} and we are done. In the latter case, if $c_2(X)^\perp \neq P$ then P will contain rational points corresponding to semi-ample divisors. Hence, we can assume that $c_2(X)^\perp = P$. Moreover, if C has maximal rank and contains a rational point, then we can conclude as above. While, if $\text{rank } C = 2$ then C is a cone and its vertex corresponds to a nef divisor D' with $D'^2 = 0 \neq c_2(X) \cdot D'$. Hence D' is semi-ample.

In case (3), as \widetilde{W} is geometrically irreducible, it follows immediately that the action of $\text{Aut}(X)$ on the second cohomology embeds in the automorphisms of the pair $(\widetilde{W}, c_2(X)^\perp_{|\widetilde{W}})$, that are finite, as \widetilde{W} is a cubic hypersurface and we are furthermore fixing the linear section $c_2(X)^\perp$; hence, the finiteness of the representation of the automorphisms of X on $N^1(X)_\mathbb{R}$ implies that the nef cone of X is rational polyhedral and in particular there are semi-ample but not ample divisors on X .

CASE $b_2(X) = 4$. Since we are assuming that the only semi-ample divisors on X are the ample ones and that $c_2(X)$ is not strictly positive on $\text{Nef}(X) \setminus \{0\}$, then [Wil94b, Proposition 2.1] implies that $\widetilde{W} = Q \cup P \subset \mathbb{P}_\mathbb{R}^3$, where Q is a rank 4 quadric with no rational points and $P = c_2(X)^\perp$. \square

To conclude, let us explain how Conjecture 7.1 is also connected to the question of whether or not Calabi–Yau threefolds are Kobayashi hyperbolic.

By [Bro78], a compact complex space is Kobayashi hyperbolic if and only if it does not contain any non-constant holomorphic curve $f: \mathbb{C} \rightarrow X$. A now classical conjecture of Kobayashi predicts that if a smooth projective variety is Kobayashi hyperbolic then the canonical class K_X is ample on X . In particular, according to this conjecture, no Calabi–Yau manifold is Kobayashi hyperbolic. Kobayashi conjecture is known to hold for smooth projective surfaces. In dimension 3, the only missing case for the verification of the conjecture is that of Calabi–Yau threefold, see [Div12] for more details about the state of the art.

It is not hard to see that if Conjecture 6.1 holds then also Kobayashi conjecture holds in dimension 3. Even though we are not able to show that if Conjecture 7.1 holds then also Conjecture 6.1 holds, it is still possible to show that X is indeed not Kobayashi hyperbolic at least when $b_2 \geq 2$.

PROPOSITION 7.3. [Div12, Proposition] *Assume that Conjecture 7.1 holds. Then the Kobayashi conjecture holds true in dimension 3, except possibly if there exists a Calabi–Yau threefold of Picard number one which is hyperbolic.*

REFERENCES

- [Bat94] V. V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi–Yau hypersurfaces in toric varieties*, J. Algebraic Geom. 3 (1994), no. 3, 493–535. ↑1584
- [Bat99] V. V. Batyrev, *Birational Calabi–Yau n -folds have equal Betti numbers*, New trends in algebraic geometry (Warwick, 1996), London Math. Soc. Lecture Note Ser., 264. Cambridge Univ. Press, Cambridge, 1999, pp. 1–11. ↑1591
- [Bea83] A. Beauville, *Variétés Kähleriennes dont la première classe de Chern est nulle*, J. Differential Geom. 18 (1983), no. 4, 755–782 (1984) (French). ↑1581
- [Bro78] R. Brody, *Compact manifolds and hyperbolicity*, Trans. Amer. Math. Soc. 235 (1978), 213–219. ↑1600
- [CCP08] F. Campana, J. A. Chen, and T. Peternell, *Strictly nef divisors*, Math. Ann. 342 (2008), no. 3, 565–585. ↑1595
- [Dem92] J.-P. Demailly, *Singular Hermitian metrics on positive line bundles*, Complex algebraic varieties (Bayreuth, 1990), Lecture Notes in Math., 1507. Springer, Berlin, 1992, pp. 87–104, DOI 10.1007/BFb0094512. MR1178721 ↑1595
- [Div12] S. Diverio, *Rational curves on Calabi–Yau threefolds and a conjecture of Oguiso*, Komplexe Analysis. Oberwolfach Rep. 9 (2012), 2612–2615, DOI 10.48550/arXiv.1311.6612. ↑1600
- [DF14] S. Diverio and A. Ferretti, *On a conjecture of Oguiso about rational curves on Calabi–Yau threefolds*, Comment. Math. Helv. 89 (2014), no. 1, 157–172. ↑1583, 1597, 1598
- [DFM19] S. Diverio, C. Fontanari, and D. Martinelli, *Rational curves on fibered Calabi–Yau manifolds*, Doc. Math. 24 (2019), 663–675. ↑1584
- [FHS21] S. Filipazzi, C. D. Hacon, and R. Svaldi, *Boundedness of elliptic Calabi–Yau threefolds* (2021), arXiv manuscript, <https://arxiv.org/abs/2112.01352>. ↑1599
- [Fuj83a] T. Fujita, *Vanishing theorems for semipositive line bundles*, Algebraic geometry (Tokyo/Kyoto, 1982), Lecture Notes in Math., 1016. Springer, Berlin, 1983, pp. 519–528. ↑1588
- [Fuj83b] T. Fujita, *Semipositive line bundles*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 30 (1983), no. 2, 353–378. ↑1588
- [GKP16] D. Greb, S. Kebekus, and T. Peternell, *Étale fundamental groups of Kawamata log terminal spaces, flat sheaves, and quotients of abelian varieties*, Duke Math. J. 165 (2016), no. 10, 1965–2004. ↑1586
- [HM07] C. D. Hacon and J. Mckernan, *On Shokurov’s rational connectedness conjecture*, Duke Math. J. 138 (2007), no. 1, 119–136, DOI 10.1215/S0012-7094-07-13813-4. MR2309156 ↑

- [Har70] R. Hartshorne, *Ample subvarieties of algebraic varieties*, Lecture Notes in Math., 156. Springer, Berlin-New York, 1970. Notes written in collaboration with C. Musili. [↑1594](#)
- [Huy16] D. Huybrechts, *Lectures on K3 surfaces*, Cambridge Studies in Advanced Mathematics, 158. Cambridge University Press, Cambridge, 2016. [MR3586372](#) [↑](#)
- [KV20] L. Kamenova and C. Vafa, *Kobayashi non-hyperbolicity of Calabi–Yau manifolds via mirror symmetry*, Comm. Math. Phys. 378 (2020), no. 1, 329–334. [↑1583](#)
- [Kaw97] Y. Kawamata, *On the cone of divisors of Calabi–Yau fiber spaces*, Internat. J. Math. 8 (1997), no. 5, 665–687. [↑1598](#)
- [Kol96] J. Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 32. Springer, Berlin, Heidelberg, 1996. [↑1588](#)
- [Kol13] J. Kollár, *Singularities of the minimal model program*, Cambridge Tracts in Mathematics, 200. Cambridge University Press, Cambridge, 2013. With a collaboration of Sándor Kovács. [↑1581](#)
- [Kol15] J. Kollár, *Deformations of elliptic Calabi–Yau manifolds*, Recent advances in algebraic geometry, London Math. Soc. Lecture Note Ser., 417. Cambridge Univ. Press, Cambridge, 2015, pp. 254–290. [↑1582](#)
- [KM98] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, 134. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti; Translated from the 1998 Japanese original. [↑1585](#)
- [KMM94] S. Keel, K. Matsuki, and J. McKernan, *Log abundance theorem for threefolds*, Duke Math. J. 75 (1994), no. 1, 99–119. [↑1582](#), [1586](#)
- [KMM04] S. Keel, K. Matsuki, and J. McKernan, *Corrections to: “Log abundance theorem for threefolds” [Duke Math. J. 75 (1994), no. 1, 99–119]*, Duke Math. J. 122 (2004), no. 3, 625–630. [↑1586](#)
- [KS00] M. Kreuzer and H. Skarke, *Complete classification of reflexive polyhedra in four dimensions*, Adv. Theor. Math. Phys. 4 (2000), no. 6, 1209–1230. [↑1584](#)
- [Laz04a] R. Lazarsfeld, *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. [Results in Mathematics and Related Areas. 3rd Series], 48. Springer, Berlin, 2004. Classical setting: line bundles and linear series. [↑1585](#), [1586](#), [1588](#), [1589](#), [1594](#)
- [Laz04b] R. Lazarsfeld, *Positivity in algebraic geometry. II*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. [Results in Mathematics and Related Areas. 3rd Series], 49. Springer, Berlin, 2004. Positivity for vector bundles, and multiplier ideals. [↑1585](#)

- [LOP18] V. Lazić, K. Oguiso, and T. Peternell, *The Morrison-Kawamata cone conjecture and abundance on Ricci flat manifolds*, Uniformization, Riemann–Hilbert correspondence, Calabi–Yau manifolds & Picard–Fuchs equations, Adv. Lect. Math. (ALM), 42. Int. Press, Somerville, MA, 2018, pp. 157–185. ↑1582, 1599
- [LOP20] V. Lazić, K. Oguiso, and T. Peternell, *Nef line bundles on Calabi–Yau threefolds. I*, Int. Math. Res. Not. IMRN 19 (2020), 6070–6119. ↑1582, 1583, 1584, 1586, 1587, 1588, 1589, 1590, 1591, 1592, 1593, 1594
- [LP13] V. Lazić and T. Peternell, *On the cone conjecture for Calabi–Yau manifolds with Picard number two*, Math. Res. Lett. 20 (2013), no. 6, 1103–1113. ↑1599
- [LP18] V. Lazić and T. Peternell, *Abundance for varieties with many differential forms*, Épijournal Géom. Algébrique 2 (2018), Art. 1, 35 p. ↑1586, 1595
- [LP20] V. Lazić and T. Peternell, *On generalised abundance. I*, Publ. Res. Inst. Math. Sci. 56 (2020), no. 2, 353–389. ↑1582
- [Miy87] Y. Miyaoka, *The Chern classes and Kodaira dimension of a minimal variety*, Algebraic geometry, Sendai, 1985, Adv. Stud. Pure Math., 10. North-Holland, Amsterdam, 1987, pp. 449–476. ↑1586
- [MM83] S. Mori and S. Mukai, *The uniruledness of the moduli space of curves of genus 11*, Algebraic geometry (Tokyo/Kyoto, 1982), Lecture Notes in Math., 1016. Springer, Berlin, 1983, pp. 334–353. ↑1596
- [Mor93] D. R. Morrison, *Compactifications of moduli spaces inspired by mirror symmetry*, Astérisque 218 (1993), 243–271. Journées de Géométrie Algébrique d’Orsay (Orsay, 1992). ↑1598
- [Nak04] N. Nakayama, *Zariski-decomposition and abundance*, MSJ Memoirs, 14. Mathematical Society of Japan, Tokyo, 2004. MR2104208 ↑1585
- [Ogu93] K. Oguiso, *On algebraic fiber space structures on a Calabi–Yau 3-fold*, Internat. J. Math. 4 (1993), no. 3, 439–465. With an appendix by Noboru Nakayama. ↑1582, 1583, 1586
- [Ogu14] K. Oguiso, *Automorphism groups of Calabi–Yau manifolds of Picard number 2*, J. Algebraic Geom. 23 (2014), no. 4, 775–795. ↑1596
- [Ser95] F. Serrano, *Strictly nef divisors and Fano threefolds*, J. Reine Angew. Math. 464 (1995), 187–206. ↑1583, 1594, 1595
- [Ver10] M. Verbitsky, *HyperKähler SYZ conjecture and semipositive line bundles*, Geom. Funct. Anal. 19 (2010), no. 5, 1481–1493. ↑1582
- [Ver15] M. Verbitsky, *Ergodic complex structures on hyperkähler manifolds*, Acta Math. 215 (2015), no. 1, 161–182. ↑1596
- [Wil89] P. M. H. Wilson, *Calabi–Yau manifolds with large Picard number*, Invent. Math. 98 (1989), no. 1, 139–155. ↑1597

- [Wil92] P. M. H. Wilson, *The Kähler cone on Calabi–Yau threefolds*, Invent. Math. 107 (1992), no. 3, 561–583. ↑1583, 1597
- [Wil93] P. M. H. Wilson, *Erratum: “The Kähler cone on Calabi–Yau threefolds” [Invent. Math. 107 (1992), no. 3, 561–583]*, Invent. Math. 114 (1993), no. 1, 231–233. ↑1597
- [Wil94a] P. M. H. Wilson, *The existence of elliptic fibre space structures on Calabi–Yau threefolds*, Math. Ann. 300 (1994), no. 4, 693–703. ↑1583, 1586, 1590
- [Wil94b] P. M. H. Wilson, *Minimal models of Calabi–Yau threefolds*, Classification of algebraic varieties (L’Aquila, 1992), Contemp. Math., 162. Amer. Math. Soc., Providence, RI, 1994, pp. 403–410. ↑1600

Haidong Liu
Sun Yat-sen University
Department of Mathematics
Guangzhou, 510275
China
liuhd35@mail.sysu.edu.cn
jiuguiaqi@gmail.com

Roberto Svaldi
Università degli Studi di Milano
Dipartimento di Matematica
“F. Enriques”
Via Saldini 50
Milano (MI) 20133
Italy
and
EPFL
SB MATH-GE, MA B1 497
(Bâtiment MA), Station 8
CH-1015 Lausanne
Switzerland
roberto.svaldi@unimi.it