

RESEARCH ARTICLE

# Uniqueness of enhancements for derived and geometric categories

Alberto Canonaco<sup>1</sup>, Amnon Neeman<sup>2</sup> and Paolo Stellari<sup>3</sup>

<sup>1</sup>Dipartimento di Matematica ‘F. Casorati’, Università degli Studi di Pavia, Via Ferrata 5, 27100 Pavia, Italy;  
E-mail: [alberto.canonaco@unipv.it](mailto:alberto.canonaco@unipv.it).

<sup>2</sup>Centre for Mathematics and Its Applications, Mathematical Sciences Institute, Building 145, The Australian National University, Canberra, ACT 2601, Australia; E-mail: [Amnon.Neeman@anu.edu.au](mailto:Amnon.Neeman@anu.edu.au).

<sup>3</sup>Dipartimento di Matematica ‘F. Enriques’, Università degli Studi di Milano, Via Cesare Saldini 50, 20133 Milano, Italy;  
E-mail: [paolo.stellari@unimi.it](mailto:paolo.stellari@unimi.it); URL: <https://sites.unimi.it/stellari>.

Received: 27 April 2022; Accepted: 15 September 2022

2020 Mathematics Subject Classification: 14F08, 18E10, 18G80

## Abstract

We prove that the derived categories of abelian categories have unique enhancements—all of them, the unbounded, bounded, bounded above and bounded below derived categories. The unseparated and left completed derived categories of a Grothendieck abelian category are also shown to have unique enhancements. Finally, we show that the derived category of complexes with quasi-coherent cohomology and the category of perfect complexes have unique enhancements for quasi-compact and quasi-separated schemes.

## Contents

|  |           |
|--|-----------|
| <b>Introduction</b>  | <b>1</b>  |
| <b>1 The triangulated categories</b>                         | <b>7</b>  |
| <b>2 Generation</b>  | <b>11</b> |
| <b>3 Dg categories and enhancements</b>                      | <b>14</b> |
| <b>4 A special zigzag of dg functors</b>                     | <b>25</b> |
| <b>5 Uniqueness for <math>D^2(\mathcal{A})</math></b>        | <b>38</b> |
| <b>6 The unseparated and completed derived categories</b>    | <b>46</b> |
| <b>7 Homotopy pullbacks and enhancements</b>                 | <b>53</b> |
| <b>8 Uniqueness of enhancements for geometric categories</b> | <b>60</b> |
| <b>References</b>  | <b>64</b> |

## Introduction

The relation between triangulated categories and their higher categorical enhancements—pretriangulated dg, pretriangulated  $A_\infty$  or stable  $\infty$ -categorical—has been under investigation for several years now. One reason is that while triangulated categories have grown remarkably important in representation theory and algebraic geometry, many of the constructions one wants to make rely on the functoriality that comes with an enhancement. Many instances of this phenomenon appeared in the

recent developments of derived algebraic geometry: for example, in relation to deformation theory and moduli problems.

In this paper, we stick to the language of dg (differential graded) categories. We recall that, roughly, a dg enhancement (or simply an enhancement) of a triangulated category  $\mathcal{T}$  is a pretriangulated dg category whose homotopy category is equivalent to  $\mathcal{T}$ . It is relevant to note that the ‘natural’ triangulated categories of algebra and geometry come with ‘natural’ dg enhancements. For example, the derived categories  $\mathbf{D}^?(A)$  of an abelian category  $A$ , where  $? = \emptyset, b, +, -$  (i.e., where the cohomology is assumed unbounded, bounded, bounded below or bounded above), as well as the category  $\mathbf{Perf}(X)$  of perfect complexes on a quasi-compact and quasi-separated scheme, all have ‘natural’ dg enhancements by construction. This existence does not hold in general. Indeed, there are well-known examples of ‘topological’ triangulated categories that do not admit dg enhancements (see, for instance, [9, Section 3.2]). More recently, it has been proved in [40] that there exist triangulated categories that are linear over a field and without a dg enhancement.

A priori, there is no good reason to expect different enhancements of the same triangulated category to be ‘comparable’. This is important because constructions that take place in an enhancement may depend on the choice of enhancement. And the right notion of ‘comparability’ turns out to be that two enhancements are declared equivalent if they agree up to isomorphism in the homotopy category  $\mathbf{Hqe}$  of the category of (small) dg categories. Consequently, one says that a triangulated category has a unique enhancement if any two enhancements are isomorphic in  $\mathbf{Hqe}$ . As  $\mathbf{Hqe}$  is the localisation of the category of dg categories by quasi-equivalences, two dg categories that are isomorphic in  $\mathbf{Hqe}$  have equivalent homotopy categories, but the converse need not be true. So far, very few examples of triangulated categories admitting non-unique enhancements have been produced. A ‘classical’ one is reported in [9, Section 3.3] (see also Corollary 6.12). If one requires that everything be linear over a field, the first example was recently found by Rizzardo and Van den Bergh [39].

Back to  $\mathbf{Hqe}$ : one can describe all morphisms in this category thanks to the seminal work of Toën [46] (see also [8, 11]). Indeed, for the natural enhancements of geometric categories, such as the bounded derived categories of coherent sheaves on smooth projective schemes, the morphisms in  $\mathbf{Hqe}$  between them are all lifts of exact functors of a special form: the so-called Fourier–Mukai functors (see [28, 46] and [9, 10] for a survey).

The way the triangulated and dg sides of this picture should be related was pinned down, in the geometric setting, in the seminal work [6] by Bondal–Larsen–Lunts, where it is conjectured that

- (C1) The geometric triangulated categories  $\mathbf{D}^b(\mathbf{Coh}(X))$ ,  $\mathbf{D}(\mathbf{Qcoh}(X))$  and  $\mathbf{Perf}(X)$  should have a unique dg enhancement when  $X$  is a quasi-projective scheme (i.e., any two dg enhancements should be isomorphic in  $\mathbf{Hqe}$ ).
- (C2) If  $X_1$  and  $X_2$  are smooth projective schemes, then all exact functors between  $\mathbf{D}^b(\mathbf{Coh}(X_1))$  and  $\mathbf{D}^b(\mathbf{Coh}(X_2))$  should lift to morphisms in  $\mathbf{Hqe}$ .

Conjecture (C2) has recently been disproved in [38], and even when a lift exists, it is not unique in general by [12]. On the other hand, special cases of (C1) have been proved to be correct, in increasing generality, by several authors over the last decade. Let us briefly go through this part of the story; after all, this article belongs to this string of results.

The first breakthrough in the direction of (C1) came from the beautiful work by Lunts and Orlov [27], which proved, among other things, that  $\mathbf{D}(\mathcal{S})$  has a unique enhancement when  $\mathcal{S}$  is a Grothendieck abelian category with a small set of compact generators. This result implies that (C1) holds true for  $\mathbf{D}(\mathbf{Qcoh}(X))$  when  $X$  is a quasi-compact and separated scheme with enough locally free sheaves (see [27, Theorem 2.10]). This was extended to all Grothendieck abelian categories in [13] by using the theory of well-generated triangulated categories. Hence (C1) holds for  $\mathbf{D}(\mathbf{Qcoh}(X))$  when  $X$  is any scheme or algebraic stack.

As for  $\mathbf{D}^b(\mathbf{Coh}(X))$  and  $\mathbf{Perf}(X)$ , Lunts and Orlov show in [27] that they have unique enhancements when  $X$  is a quasi-projective scheme (see Theorems 2.12 and 2.13 in [27]). Clearly, together with the previous result, this implies that (C1) holds even in greater generality. Actually, an additional

improvement of the argument in [27] allowed the first and third author to prove that  $\mathbf{D}^b(\mathbf{Coh}(X))$  and  $\mathbf{Perf}(X)$  have unique enhancements when  $X$  is any noetherian scheme with enough locally free sheaves (see Corollaries 6.11 and 7.2 in [13]).

Fresh air was brought into the subject with the advent of the powerful theory of  $\infty$ -categories. Specifically, Antieau [1] reconsidered the problem of uniqueness of enhancements, taking a completely different approach employing Lurie's work on prestable  $\infty$ -categories (see [30, Appendix C]). Using this amazing machinery, he proved the beautiful result that  $\mathbf{D}^?(A)$  has a unique enhancement when  $? = b, +, -$  and  $A$  is any small abelian category. It should be noted that restricting to small categories is a minor issue, as explained in Section 3.5.

If  $A$  is not only abelian but also a Grothendieck category, one can, following Lurie, construct three interesting triangulated categories out of  $A$ : the usual derived category  $\mathbf{D}(A)$ , the *unseparated derived category*  $\check{\mathbf{D}}(A)$  and the *left completed derived category*  $\widehat{\mathbf{D}}(A)$ . We know all about  $\mathbf{D}(A)$ , and, in particular, we know that it has a unique enhancement by [13]. The triangulated category  $\check{\mathbf{D}}(A)$  is nothing but the homotopy category of injectives in  $A$  and has been extensively studied by Krause in [23]. The uniqueness of enhancements for  $\check{\mathbf{D}}(A)$ , when  $A$  is not only Grothendieck but also locally coherent, is one of the main results of Antieau (see [1, Theorem 1]). The triangulated category  $\widehat{\mathbf{D}}(A)$  is more mysterious. It does not seem to have a purely triangulated description, and it should be thought of as a remedy to the fact that, in general,  $\mathbf{D}(A)$  is not left complete (see [32]). In [1], the uniqueness of the enhancement for  $\widehat{\mathbf{D}}(A)$  is stated as an open and challenging problem (see [1, Question 8.1]).

Antieau's striking achievements offered what appeared to be conclusive evidence of the superiority of the  $\infty$ -category machine, and our initial, humble goal was to try to find out how much of his opus could be obtained by more primitive methods. The authors were surprised by the outcome of what started as a modest project: beginning with a few simple new ideas, we ended up not only improving on Antieau's results, but also solving most of the open problems in the literature.

Our first precise statement is the following:

**Theorem A.** *Let  $A$  be an abelian category:*

- (1) *The triangulated category  $\mathbf{D}^?(A)$  has a unique dg enhancement when  $? = b, +, -, \emptyset$ .*
- (2) *If  $A$  is a Grothendieck abelian category, then  $\check{\mathbf{D}}(A)$  and  $\widehat{\mathbf{D}}(A)$  have unique dg enhancements.*

The striking and new part of (1) is the uniqueness for  $\mathbf{D}(A)$  for every abelian category  $A$ . Nonetheless, our approach will uniformly and harmlessly produce uniqueness of enhancements for  $\mathbf{D}^?(A)$ , for  $? = b, -, +$ , thus recovering Antieau's results in a completely different way. Part (2) of Theorem A on the one hand generalises [1, Theorem 1] and, on the other hand, answers the questions in [1] about  $\widehat{\mathbf{D}}(A)$  that we mentioned above. In particular, Theorem A (2) gives a positive answer to Question 4.7 in [9].

Back to the geometric setting. If  $X$  is a quasi-compact and quasi-separated scheme, then the category  $\mathbf{D}(\mathbf{Qcoh}(X))$  is in general not equivalent to  $\mathbf{D}_{\mathbf{qc}}(X)$ , the full triangulated subcategory of the category of complexes of  $\mathcal{O}_X$ -modules consisting of all complexes with quasi-coherent cohomology. Thus the uniqueness of dg enhancements for  $\mathbf{D}_{\mathbf{qc}}(X)$  cannot directly be deduced from Theorem A. Nonetheless, this category is of primary interest: for example, because the category of perfect complexes on  $X$  coincides with the subcategory of compact objects in  $\mathbf{D}_{\mathbf{qc}}(X)$ .

The uniqueness of dg enhancements for  $\mathbf{D}_{\mathbf{qc}}(X)$  and  $\mathbf{Perf}(X)$  was formulated as an open problem by Antieau in [1, Question 8.16], and our second main result positively answers his question in the context of dg enhancements:

**Theorem B.** *Let  $X$  be a quasi-compact and quasi-separated scheme. Then the categories  $\mathbf{D}_{\mathbf{qc}}^?(X)$  and  $\mathbf{Perf}(X)$  have a unique dg enhancement for  $? = b, +, -, \emptyset$ .*

The case of  $\mathbf{Perf}(X)$  is covered by [1, Corollary 9] under the stronger assumption that the scheme is quasi-compact, quasi-separated and 0-complicial. The latter condition, which we do not need to make explicit here, roughly refers to a property of  $\mathbf{Perf}(X)$  induced by the t-structure on  $\mathbf{D}_{\mathbf{qc}}(X)$  and predicts how perfect complexes with only non-negative cohomologies are generated by perfect quasi-coherent

sheaves. As we will explain below, part of the interest of Theorem B is that the proof introduces a new technique to study the uniqueness of enhancements based on homotopy limits.

### *Applications: past, present and future*

We have already said that the key to our approach lies in a few simple new ideas. One can ask if these ideas might be relevant or useful in other contexts. Hence we should mention that a linchpin to our approach, namely Proposition 2.9 and its proof, has already been used by the second author in [31] to provide a counterexample to conjectures by Schlichting [43] and by Antieau, Gepner and Heller [2] about vanishing in negative  $K$ -theory. Note that this application is totally unrelated to the content of the current article.

Within the present paper, there are further, easy applications of our main results. In particular, we deduce that the following triangulated categories have unique enhancements:

- $\mathbf{D}^?(Qcoh(X))$ , for  $X$  any scheme or algebraic stack, and  $\mathbf{D}^?(Coh(X))$ , for any scheme or any locally noetherian algebraic stack, for  $? = \emptyset, b, +, -$  (see Corollary 5.9);
- $\mathbf{D}(\mathcal{G})^\alpha$ , where  $\mathcal{G}$  is a Grothendieck abelian category and  $\alpha$  is a large regular cardinal (see Corollary 5.8). Recall here that  $\mathbf{D}(\mathcal{G})$  is a well-generated triangulated category and  $\mathbf{D}(\mathcal{G})^\alpha$  is the full triangulated subcategory consisting of its  $\alpha$ -compact objects;
- $\mathbf{K}^?(\mathcal{A})$ , for any abelian category  $\mathcal{A}$  and for  $? = \emptyset, b, +, -$  (see Corollary 5.6).

The first two items together yield a complete positive answer to Question 4.8 in [10].

Another interesting and surprising application, discussed in Section 5.5, is that we recover and generalise the construction of the *realisation functor* of Beilinson, Bernstein and Deligne [4]. This functor plays a key role in the study of triangulated categories with  $t$ -structures. The original, involved proof is replaced in Section 5.5 by a different approach, combining the vital Proposition 2.9 with the techniques in Section 4 to deliver the results easily.

In future work, we will analyse how our uniqueness results apply to study the liftability of exact functors along the lines of (C2) above. More specifically, we will investigate a classical conjecture by Rickard asserting that all autoequivalences of  $\mathbf{D}(\text{Mod}(R))$ , where  $R$  is a commutative ring, are liftable. We have already (briefly) discussed the case of projective schemes. The affine case is still very challenging and essentially open. We will explain how our techniques provide simplifications and generalisations of the existing results.

Still with an eye to the future, we conclude this discussion by pointing out that there remain a couple of situations of high algebro-geometric interest where the (non-)uniqueness of the enhancements needs to be fully understood: the categories of matrix factorisations and the case of admissible subcategories of triangulated categories admitting a unique enhancement. If we work with categories and functors linear over  $\mathbb{Z}$ , then there are examples of categories of matrix factorisations with non-unique enhancements (see [17, 42]). Similarly, in Section 6.4, we provide an example of a  $\mathbb{Z}$ -linear triangulated category with a unique enhancement (by Theorem A) but with an admissible subcategory with non-unique  $\mathbb{Z}$ -linear enhancements (see Corollary 6.12). It remains open to understand if similar examples can be found for categories linear over a field and if one can find admissible subcategories with non-unique enhancements in  $\mathbf{D}^b(Coh(X))$  when  $X$  is a smooth projective scheme.

### *The strategy of the proofs*

The one-sentence summary of the proof of Theorem A (1) would be that it is an elaborate study of special generators for  $\mathbf{D}^?(\mathcal{A})$ , coupled with a suitable description of  $\mathbf{D}^?(\mathcal{A})$  as a Verdier quotient. The same principle underlies all the existing papers in the literature proving the uniqueness of enhancements of derived categories of abelian categories. The many papers differ in which generators they use and what quotient they study.

In the current article, we realise  $\mathbf{D}^2(\mathcal{A})$  as a quotient of the homotopy category  $\mathbf{K}^2(\mathcal{A})$  and show that  $\mathbf{K}^2(\mathcal{A})$  is generated in 3 steps by objects that are direct sums of shifts of objects in  $\mathcal{A}$  (Proposition 2.9 and Corollary 2.10). The key fact here, namely that  $\mathbf{K}^2(\mathcal{A})$  is generated in 3 steps by the simple objects described above, can be seen by combining Krause's [23, Lemma 3.1 and its proof] with Max Kelly's old result [22] (see [37, Theorem 7.5 and its proof] for a modern account). But we include a full proof in this article because we make use of the explicit three steps that suffice.

This is different from and, in a sense, more natural than the point of view of [27] and [13, 14]. In those earlier papers, to prove that  $\mathbf{D}(\mathcal{G})$  has a unique enhancement for  $\mathcal{G}$  a Grothendieck abelian category, one uses the strong and special property that  $\mathcal{G}$  has a generator. Thus one can take generators for  $\mathbf{D}(\mathcal{G})$ , which all live in degree 0, and  $\mathbf{D}(\mathcal{G})$  is a suitable quotient of the derived category of modules over the category formed by these generators.

The technical complications of our approach involve, at the triangulated level, a careful analysis of certain special products and coproducts. It is discussed in Section 1 and reverberates at the dg level where one has to construct suitable dg enhancements of  $\mathbf{K}^2(\mathcal{A})$  and  $\mathbf{D}^2(\mathcal{A})$  and an intricate zigzag of dg functors linking them. The dg work is carried out in Section 4. Again, the dg part of the argument is simpler in [27, 13] and involves a short zigzag diagram consisting of one roof of dg functors. The last part of the proof in Section 5 is then very close in spirit to the argument in Section 4 and 5 of [27] (and thus in Section 4 of [13]).

The proof of the uniqueness of dg enhancements for  $\check{\mathbf{D}}(\mathcal{A})$  in Section 6.1 is a reduction to Theorem C in [13, 14] (see Theorem 6.2). It uses the work of Krause [23] to show that  $\check{\mathbf{D}}(\mathcal{A})$  is a well-generated triangulated category and can be written as a quotient of the derived category of the abelian category of modules over the abelian subcategory  $\mathcal{A}^\alpha$  of  $\mathcal{A}$ , which consists of the  $\alpha$ -presentable objects in  $\mathcal{A}$  (here,  $\alpha$  is a sufficiently large regular cardinal).

Finally, the case of  $\widehat{\mathbf{D}}(\mathcal{A})$  is studied in Section 6.3, and the proof makes use of the natural t-structure induced on  $\widehat{\mathbf{D}}(\mathcal{A})$  by  $\mathbf{D}(\mathcal{A})$ . With this t-structure, we have a natural equivalence  $\widehat{\mathbf{D}}(\mathcal{A})^+ \cong \mathbf{D}^+(\mathcal{A})$ . We can invoke Theorem A (1) and then deduce the result by a careful analysis of the compatibility with homotopy colimits. It should be noted that here we need to use that  $\mathbf{D}^2(\mathcal{A})$  has a semi-strongly unique dg enhancement (see Remark 5.4). Roughly, this means if  $\mathcal{E}_1$  and  $\mathcal{E}_2$  are two dg enhancements of  $\mathbf{D}^2(\mathcal{A})$  (i.e., there are exact equivalences  $E_i: H^0(\mathcal{E}_i) \xrightarrow{\sim} \mathbf{D}^2(\mathcal{A})$ ), then the isomorphism  $f: \mathcal{E}_1 \xrightarrow{\sim} \mathcal{E}_2$  in  $\mathbf{Hqe}$  provided by Theorem A (1) is such that  $H^0(f)(X) \cong E_2^{-1} \circ E_1(X)$ , for all  $X$  in  $H^0(\mathcal{E}_1)$ .

The strategy of the proof of Theorem B is new and is based on the idea of realising a dg enhancement of  $\mathbf{D}_{\text{qc}}^2(X)$  (and of  $\mathbf{Perf}(X)$ ) as the homotopy limit of dg enhancements of the derived category of the open subschemes in an affine open cover of  $X$ . More precisely: in Section 7, we prove that given any enhancement  $\mathcal{C}$  of  $\mathbf{D}_{\text{qc}}^2(X)$  (or  $\mathbf{Perf}(X)$ ), one can produce an isomorphism in  $\mathbf{Hqe}$  between  $\mathcal{C}$  and the homotopy limit of induced enhancements of the derived categories of the affine schemes in the cover (and of their finite intersections). This can be deduced from Theorem 7.4, which is a general criterion involving the simpler case of homotopy pullbacks.

This has the clear advantage that for each Zariski open subset  $U$  in the covering of  $X$  and all their finite intersections, one knows that  $\mathbf{D}_{\text{qc}}^2(U) \cong \mathbf{D}^2(\mathbf{Qcoh}(U))$ , since  $U$  is quasi-compact and separated. Thus the uniqueness of their enhancements is guaranteed by Theorem A (1). The hard work comes in showing that the constructions in Section 4, and thus the proof of Theorem A (1) in Section 5, are compatible with restriction to appropriate open subschemes. In Section 8, we show this compatibility with the special homotopy limits we are considering, concluding the proof.

### Related work

Throughout this article, we work with dg enhancements. Since Antieau's enhancements are stable  $\infty$ -categories, a comparison requires one to invoke results like [1, Meta Theorem 13] and [15]. These results assert that for most of the triangulated categories we study, the difference is immaterial; the uniqueness of enhancement problems in the two different settings are equivalent. This is clear for the triangulated categories in Theorem A. As for the categories  $\mathbf{Perf}(X)$  and  $\mathbf{D}_{\text{qc}}^2(X)$  in

Theorem B, the situation is a bit more delicate. Indeed, as pointed out in Section 8.2 of [1], Meta Theorem 13 may not automatically apply, and, by [15], Theorem B only implies that the two triangulated categories mentioned above have unique  $\mathbb{Z}$ -linearised stable  $\infty$ -enhancements. We leave this to the interested reader.

The reason we stick to (pre)triangulated dg categories is for convenience—it makes the key Proposition 2.9 and Corollary 2.10 easier and cleaner to use. In the triangulated category  $\mathbf{D}^?(A)$ , the generators given by Proposition 2.9 and Corollary 2.10 have (non-canonical) direct sum decompositions, and using this is easier in enhancements that are additive. In fact, the interested reader can check that for us, the case where  $? = b$  is easy. The subtlety comes in dealing with  $? \in \{+, -, \infty\}$ , where issues of limits come up. And as far as we can tell, the  $\infty$ -category machine would not help much here.

Of course, Antieau [1] has taught us that a clever application of the  $\infty$ -category machine can lead to spectacular progress. It is entirely possible that our new ideas, combined with the powerful machinery, will produce startling advances. For example, the local-to-global approach of Theorem B might readily be amenable to the  $\infty$ -category methods.

We leave this to the experts.

### *Structure of the paper*

In Section 1, we deal with some foundational questions about products and coproducts in the triangulated categories  $\mathbf{K}^?(A)$  and  $\mathbf{D}^?(A)$ . We also discuss their behaviour with respect to exact functors.

Section 2 introduces various notions of generation for triangulated categories. In particular, in Section 2.1, we define well-generated triangulated categories, and we set the stage for our study of  $\check{\mathbf{D}}(\mathcal{G})$ . Section 2.2 is about the approach to the generation of  $\mathbf{K}^?(A)$  and  $\mathbf{D}^?(A)$ , which is crucial in the proof of Theorem A (1).

In Section 3, we introduce some standard material about dg categories and, at the same time, slightly extend known results and constructions such as Drinfeld’s notion of homotopy flat resolution (see Section 3.2). In Section 3.3, we study homotopy limits and homotopy pullbacks. We also reconsider localisations in the dg context (Section 3.4) and carefully define the notion of dg enhancement and why their uniqueness is independent of the universe (see Section 3.5).

Section 4 is devoted to the construction of appropriate enhancements for  $\mathbf{K}^?(A)$  and  $\mathbf{D}^?(A)$ . This naturally leads to the proof of Theorem A (1) in Section 5. The second part of Theorem A is proved in the subsequent section and uses somewhat different techniques.

The proof of Theorem B occupies Section 7 and Section 8. The first of the two sections sets up the general technique and criterion that link homotopy pullbacks and limits to dg enhancements. The second combines this with Theorem A to finish the argument.

### *Notation and conventions*

All preadditive categories and all additive functors are assumed to be  $\mathbb{k}$ -linear for a fixed commutative ring  $\mathbb{k}$ . By a  $\mathbb{k}$ -linear category, we mean a category whose Hom-spaces are  $\mathbb{k}$ -modules and such that the compositions are  $\mathbb{k}$ -bilinear, not assuming that finite coproducts exist.

With the small exception of Section 3.5, throughout the paper, we assume that a universe containing an infinite set is fixed. Several definitions concerning dg categories need special care because they may, in principle, require a change of universe. The major subtle logical issues in this sense can be overcome in view of [27, Appendix A] and Section 3.5. A careful reader should look at them; but in this paper, after these delicate issues are appropriately discussed, we will not explicitly mention the universe we are working in. The members of this universe will be called small sets. Unless stated otherwise, we always assume that the Hom-spaces in a category form a small set. A category is called *small* if the isomorphism classes of its objects form a small set.

If  $\mathcal{T}$  is a triangulated category and  $\mathcal{S}$  a full triangulated subcategory of  $\mathcal{T}$ , we denote by  $\mathcal{T}/\mathcal{S}$  the Verdier quotient of  $\mathcal{T}$  by  $\mathcal{S}$ . In general,  $\mathcal{T}/\mathcal{S}$  is not a category according to our convention (meaning

the Hom-spaces in  $\mathcal{T}/\mathcal{S}$  need not be small sets), but it is in many common situations, for instance when  $\mathcal{T}$  is small.

### 1. The triangulated categories

In this section, we discuss some properties of most of the triangulated categories whose dg enhancements are studied in this paper. The focus is on the existence of (co)products of special objects and the commutativity of such (co)products with exact functors.

#### 1.1. The categories

Let us recall that when  $\mathcal{A}$  is a small additive category, then  $\mathbf{K}(\mathcal{A})$  denotes the homotopy category of complexes. Namely, its objects are cochain complexes of objects in  $\mathcal{A}$ , while its morphisms are homotopy equivalence classes of morphisms of complexes. For  $A^* \in \text{Ob}(\mathbf{K}(\mathcal{A}))$ , we denote by  $A^i$  its  $i$ th component. We can then define the full subcategories  $\mathbf{K}^b(\mathcal{A})$ ,  $\mathbf{K}^+(\mathcal{A})$ ,  $\mathbf{K}^-(\mathcal{A})$  of the category  $\mathbf{K}(\mathcal{A})$  whose objects are

$$\begin{aligned} \text{Ob}(\mathbf{K}^b(\mathcal{A})) &= \{A^* \in \mathbf{K}(\mathcal{A}) : A^i = 0 \text{ for all } |i| \gg 0\} \\ \text{Ob}(\mathbf{K}^+(\mathcal{A})) &= \{A^* \in \mathbf{K}(\mathcal{A}) : A^i = 0 \text{ for all } i \ll 0\} \\ \text{Ob}(\mathbf{K}^-(\mathcal{A})) &= \{A^* \in \mathbf{K}(\mathcal{A}) : A^i = 0 \text{ for all } i \gg 0\}. \end{aligned}$$

For  $? = b, +, -, \emptyset$ , we single out the full subcategory  $\mathbf{V}^?( \mathcal{A}) \subseteq \mathbf{K}^?( \mathcal{A})$  consisting of objects with zero differentials. The properties of such a subcategory will be studied in Section 2 and will be crucial in the rest of this paper. Here we just point out that for an object  $A^* \in \mathbf{V}^?( \mathcal{A})$ , we will use the shorthand

$$\bigoplus_{i \in \mathbb{Z}} A^i[-i]$$

to remind that the object  $A^i \in \mathcal{A}$  is placed in degree  $i$ .

**Remark 1.1.** It is not difficult to prove that  $\bigoplus_{i \in \mathbb{Z}} A^i[-i]$  satisfies both the universal property of product and coproduct in  $\mathbf{K}^?( \mathcal{A})$ . Namely, there are canonical isomorphisms

$$\bigoplus_{i \in \mathbb{Z}} A^i[-i] \cong \prod_{i \in \mathbb{Z}} A^i[-i] \cong \coprod_{i \in \mathbb{Z}} A^i[-i].$$

When  $\mathcal{A}$  is an abelian category, the full triangulated subcategory  $\mathbf{K}_{\text{acy}}^?( \mathcal{A}) \subseteq \mathbf{K}^?( \mathcal{A})$  consists of *acyclic complexes*: that is, objects in  $\mathbf{K}(\mathcal{A})$  with trivial cohomology. The triangulated category  $\mathbf{D}^?( \mathcal{A})$  is then the Verdier quotient of  $\mathbf{K}^?( \mathcal{A})$  by  $\mathbf{K}_{\text{acy}}^?( \mathcal{A})$ , and it comes with a quotient functor

$$\mathbf{Q} : \mathbf{K}^?( \mathcal{A}) \longrightarrow \mathbf{D}^?( \mathcal{A}). \tag{1.1}$$

We can then consider the full subcategory  $\mathbf{B}^?( \mathcal{A}) \subseteq \mathbf{D}^?( \mathcal{A})$  as

$$\mathbf{B}^?( \mathcal{A}) := \mathbf{Q}(\mathbf{V}^?( \mathcal{A})).$$

**Remark 1.2.** By the definition of the Verdier quotient,  $\mathbf{B}^?( \mathcal{A})$  has the same objects as  $\mathbf{V}^?( \mathcal{A})$ . Thus we will freely denote them by  $\bigoplus A^i[-i]$ . But since the morphisms in  $\mathbf{D}^?( \mathcal{A})$  differ from those in  $\mathbf{K}^?( \mathcal{A})$  in a significant way, we should not expect  $\bigoplus A^i[-i]$  to automatically satisfy the universal properties of either product or coproduct in  $\mathbf{D}^?( \mathcal{A})$ . This will be discussed later.

**Remark 1.3.** It is interesting to observe that if  $\mathcal{A}$  is a small abelian category, then [23, Lemma 3.1] implies that there is a (small) abelian category  $\mathcal{B}$  and an exact equivalence  $\mathbf{K}^?( \mathcal{A}) \cong \mathbf{D}^?( \mathcal{B})$  for

$? = b, +, -, \emptyset$ . To be precise, the category  $\mathcal{B}$  is the abelian category of coherent  $\mathcal{A}$ -modules. The result follows from [47, Proposition III 2.4.4 (c)], once we observe that any coherent  $\mathcal{A}$ -module has a projective resolution of length at most 2 (see the proof of [23, Lemma 3.1]).

If  $\mathcal{G}$  is a Grothendieck abelian category, then it contains enough injectives, and one can take the full subcategory  $\text{Inj}(\mathcal{G}) \subseteq \mathcal{G}$  of injective objects. According to Lurie’s terminology [29], the *unseparated derived category* of  $\mathcal{G}$  is the triangulated category

$$\check{\mathbf{D}}(\mathcal{G}) := \mathbf{K}(\text{Inj}(\mathcal{G})).$$

This category has been extensively studied by the second author [35] and Krause [23]. Some of its properties will be recalled later. For the moment, we content ourselves with the simple observation that it fits in the following localisation sequence

$$\mathbf{K}_{\text{acy}}(\text{Inj}(\mathcal{G})) \xrightarrow{\mathbf{J}} \check{\mathbf{D}}(\mathcal{G}) \xrightarrow{\mathbf{Q}} \mathbf{D}(\mathcal{G}).$$

Namely,  $\mathbf{Q}$  and  $\mathbf{J}$  have right adjoints  $\mathbf{Q}^\rho$  and  $\mathbf{J}^\rho$ , respectively. Under some additional assumptions (e.g., if  $\mathbf{D}(\mathcal{G})$  is compactly generated),  $\mathbf{Q}$  and  $\mathbf{J}$  have left adjoints  $\mathbf{Q}^\lambda$  and  $\mathbf{J}^\lambda$  as well.

**Example 1.4.** A case where the latter situation is realised can be obtained as follows. Let  $p$  be a prime number and  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$  the field with  $p$  elements. Consider the rings  $R_1 := \mathbb{Z}/p^2\mathbb{Z}$  and  $R_2 := \mathbb{F}_p[\varepsilon]$  (where  $\varepsilon^2 = 0$ ). For  $i = 1, 2$ , set  $\mathcal{T}_i := \check{\mathbf{D}}(\text{Mod}(R_i))$ , and denote by  $\mathcal{S}_i$  the full subcategory of  $\mathcal{T}_i$  consisting of acyclic complexes. The ring  $R_i$  is noetherian and  $\mathbf{D}(\text{Mod}(R_i))$  is compactly generated. Thus the inclusion of  $\mathcal{S}_i$  has left adjoint, and  $\mathcal{S}_i$  is a localising and admissible subcategory of  $\mathcal{T}_i$ .

The last triangulated category we study in this paper is the completed derived category of a Grothendieck category. Since, to the best of our knowledge, its definition intrinsically involves pre-triangulated dg categories, this discussion is postponed to Section 6.2.

### 1.2. More on products and coproducts

As we observed in Remark 1.2, objects with zero differentials need not always agree with the obvious products or coproducts in  $\mathbf{D}^2(\mathcal{A})$ . In this section, we provide sufficient conditions for agreement.

Let us introduce some notation, slightly generalising the problem. Let  $\mathcal{A}$  be a small abelian category, and let  $\{A_n^*\}_{n \geq 0}$  be a sequence of objects in  $\mathbf{D}^2(\mathcal{A})$ . If either  $A_n^i = 0$  for all  $i > -n$  or  $A_n^i = 0$  for all  $i < n$ , we can consider the complex  $\bigoplus_{n=0}^\infty A_n^* \in \mathbf{D}(\mathcal{A})$ , which is the termwise direct sum of the complexes  $A_n$ . Note that this makes sense because under our assumptions, each term of the complex  $\bigoplus_{n=0}^\infty A_n^* \in \mathbf{D}(\mathcal{A})$  consists of a finite direct sum.

With this in mind, we can prove the following.

**Lemma 1.5.** *Let  $\mathcal{A}$  be a small abelian category, and let  $\{A_n^*\}_{n \geq 0} \subseteq \text{Ob}(\mathbf{D}^2(\mathcal{A}))$ :*

- (1) *If  $A_n^i = 0$  for all  $i > -n$  and  $? = -, \emptyset$ , then  $\bigoplus_{n=0}^\infty A_n^*$  is a coproduct in  $\mathbf{D}^2(\mathcal{A})$ : that is, there is a canonical isomorphism*

$$\bigoplus_{n=0}^\infty A_n^* \cong \coprod_{n=0}^\infty A_n^*.$$


- (2) *If  $A_n^i = 0$  for all  $i < n$  and  $? = +, \emptyset$ , then  $\bigoplus_{n=0}^\infty A_n^*$  is a product in  $\mathbf{D}^2(\mathcal{A})$ : that is, there is a canonical isomorphism*

$$\bigoplus_{n=0}^\infty A_n^* \cong \prod_{n=0}^\infty A_n^*.$$



*Proof.* The statements in (1) and (2) are obtained one from the other by passing to the opposite categories. Thus we just need to prove (1) and show that  $\bigoplus_{n=0}^{\infty} A_n^*$  satisfies the universal property of a coproduct in  $\mathbf{D}^?(\mathcal{A})$  for  $? = -, \emptyset$ .

Suppose therefore that  $B^*$  is an object of  $\mathbf{D}^?(\mathcal{A})$  and that we are given morphisms  $\varphi_n : A_n^* \rightarrow B^*$ . For  $n \geq 0$ ,  $\varphi_n$  can be represented in  $\mathbf{K}^?(\mathcal{A})$  by a roof



This means  $\tilde{A}_n^* \rightarrow A_n^* \rightarrow N_n^*$  is a distinguished triangle in  $\mathbf{K}^?(\mathcal{A})$  with  $N_n^* \in \mathbf{K}_{\text{acy}}(\mathcal{A})$ , and  $\varphi_n = \beta \circ \alpha^{-1}$  in  $\mathbf{D}^?(\mathcal{A})$ .

As  $A_n^i = 0$  for all  $i > -n$ , the cochain map  $A_n^* \rightarrow N_n^*$  must factor as  $A_n^* \rightarrow N_n^{\leq -n} \rightarrow N_n^*$ , where  $N_n^{\leq -n}$  is still acyclic but vanishes in degrees  $> -n$ . Here  $N_n^{\leq -n}$  is the truncation


$$\dots \rightarrow N^{-n-2} \rightarrow N^{-n-1} \rightarrow K \rightarrow 0 \rightarrow 0 \rightarrow \dots,$$

where  $K$  is the kernel of the map  $N^{-n} \rightarrow N^{-n+1}$ .


Now we form in  $\mathbf{K}^?(\mathcal{A})$  the morphism of distinguished triangles

$$\begin{array}{ccccc}
 \tilde{A}_n^* & \longrightarrow & A_n^* & \longrightarrow & N_n^{\leq -n} \\
 \downarrow & & \parallel & & \downarrow \\
 \tilde{A}_n^* & \longrightarrow & A_n^* & \longrightarrow & N_n^*
 \end{array}$$

which allows us to represent the morphism  $\varphi_n : A_n^* \rightarrow B^*$  by the roof




with  $\hat{A}_n^i = 0$  for all  $i > -n + 1$ . And now the roof



is a well-defined diagram in  $\mathbf{K}^?(\mathcal{A})$  since in each degree, the sums are finite. Hence we obtain a morphism  $\bigoplus_{n=0}^{\infty} A_n^* \rightarrow B^*$  in  $\mathbf{D}^?(\mathcal{A})$ , and obviously the composite  $A_n^* \rightarrow \bigoplus_{n=0}^{\infty} A_n^* \rightarrow B^*$  is  $\varphi_n$  for every  $n$ .

It remains to prove that such a morphism is unique. This is equivalent to showing that given a morphism  $\varphi : \bigoplus_{n=0}^{\infty} A_n^* \rightarrow B^*$  in  $\mathbf{D}^2(\mathcal{A})$  such that the composites  $A_n^* \rightarrow \bigoplus_{n=0}^{\infty} A_n^* \rightarrow B^*$  vanish for every  $n$ ,  $\varphi$  vanishes.

Such a  $\varphi$  may be represented in  $\mathbf{K}^2(\mathcal{A})$  by a roof



meaning  $N^* \rightarrow B^* \rightarrow \tilde{B}^*$  is a distinguished triangle in  $\mathbf{K}^2(\mathcal{A})$  with  $N^* \in \mathbf{K}_{\text{acy}}(\mathcal{A})$  and  $\varphi = \beta^{-1} \circ \alpha$  in  $\mathbf{D}^2(\mathcal{A})$ .

If the composite  $A_n^* \rightarrow \bigoplus_{n=0}^{\infty} A_n^* \rightarrow B^*$  vanishes in  $\mathbf{D}^2(\mathcal{A})$ , then the morphism  $A_n^* \rightarrow \bigoplus_{n=0}^{\infty} A_n^* \rightarrow \tilde{B}^*$  is a morphism in  $\mathbf{K}^2(\mathcal{A})$  whose image in  $\mathbf{D}^2(\mathcal{A})$  vanishes, and hence it must factor in  $\mathbf{K}^2(\mathcal{A})$  as  $A_n^* \rightarrow M_n^* \rightarrow \tilde{B}^*$ , with  $M_n^* \in \mathbf{K}_{\text{acy}}(\mathcal{A})$ . As in the first part of the proof, the map  $A_n^* \rightarrow M_n^*$  must factor as  $A_n^* \rightarrow M_n^{\leq -n} \rightarrow M_n^*$ , where  $M_n^{\leq -n}$  is acyclic and vanishes in degrees  $> -n$ . But then  $\bigoplus_{n=0}^{\infty} A_n^* \rightarrow \tilde{B}^*$  factors in  $\mathbf{K}^2(\mathcal{A})$  as

$$\bigoplus_{n=0}^{\infty} A_n^* \longrightarrow \bigoplus_{n=0}^{\infty} M_n^{\leq -n} \longrightarrow \tilde{B}^*,$$

showing that  $\varphi$  vanishes in  $\mathbf{D}^2(\mathcal{A})$ . □

The following is then a straightforward consequence.

**Corollary 1.6.** *Let  $\mathcal{A}$  be a small abelian category, and let  $\{A^i\}_{i \geq 0} \subseteq \text{Ob}(\mathcal{A})$ . Then  $\bigoplus_{i=0}^{\infty} A^i[k+i]$  is a coproduct in  $\mathbf{D}^-(\mathcal{A})$  and  $\mathbf{D}(\mathcal{A})$ , while  $\bigoplus_{i=0}^{\infty} A^i[k-i]$  is a product in  $\mathbf{D}^+(\mathcal{A})$  and  $\mathbf{D}(\mathcal{A})$  for all  $k \in \mathbb{Z}$ .*

**1.3. (Co)products and functors**

We continue with some technical results that will be used later. In particular, in this section, we investigate when special exact functors commute with products and, dually, with coproducts.

If  $\mathcal{A}$  is an abelian category and  $n \in \mathbb{Z}$ , we denote by  $\mathbf{D}^2(\mathcal{A})^{\geq n}$  (respectively,  $\mathbf{D}^2(\mathcal{A})^{\leq n}$ ) the full subcategory of  $\mathbf{D}^2(\mathcal{A})$  consisting of objects with trivial cohomologies in degrees  $< n$  (respectively,  $> n$ ).

**Proposition 1.7.** *Let  $\mathcal{A}$  be a small abelian category and  $F : \mathcal{T} \rightarrow \mathbf{D}^2(\mathcal{A})$  an additive functor, where  $? = +, \emptyset$ . Assume that  $\{T_i\}_{1 \leq i < \infty} \subseteq \text{Ob}(\mathcal{T})$  is such that*

- (i) *The product  $\prod_{i=1}^{\infty} T_i$  exists in  $\mathcal{T}$ .*
- (ii) *For every integer  $n > 0$ , there exists an integer  $m(n) > 0$  such that  $F\left(\prod_{i=m(n)}^{\infty} T_i\right) \in \mathbf{D}(\mathcal{A})^{\geq n}$ .*

*Then the product  $\prod_{i=1}^{\infty} F(T_i)$  exists in  $\mathbf{D}^2(\mathcal{A})$  and the canonical map*

$$F\left(\prod_{i=1}^{\infty} T_i\right) \longrightarrow \prod_{i=1}^{\infty} F(T_i)$$

*is an isomorphism.*

*Proof.* Given  $k > 0$  and  $j \geq k$ , the object  $F(T_j)$  is a direct summand of  $F(\prod_{i=k}^\infty T_i)$ . Thus, assumption (ii) implies that for every  $n > 0$ , there exists  $m(n) > 0$  with  $F(T_j) \in \mathbf{D}(\mathcal{A})^{\geq n}$  for all  $j \geq m(n)$ . For  $n > 0$ , we set

$$A_n := F\left(\prod_{i=m(n)}^{m(n+1)-1} T_i\right) \cong \prod_{i=m(n)}^{m(n+1)-1} F(T_i).$$

By the previous discussion,  $A_n \in \mathbf{D}(\mathcal{A})^{\geq n}$ , and Lemma 1.5 implies that  $\prod_{n=1}^\infty A_n$  exists in  $\mathbf{D}^2(\mathcal{A})$ . Thus

$$\prod_{i=1}^\infty F(T_i) \cong \left(\prod_{i=1}^{m(1)-1} F(T_i)\right) \prod_{n=1}^\infty \left(\prod_{i=1}^\infty A_n\right)$$

exists in  $\mathbf{D}^2(\mathcal{A})$ .

Let us move to the second part of the statement. For  $m \geq 2$ , the natural map

$$\theta: F\left(\prod_{i=1}^\infty T_i\right) \longrightarrow \prod_{i=1}^\infty F(T_i)$$

can be identified with the product of the two natural maps

$$\theta_1: F\left(\prod_{i=1}^{m-1} T_i\right) \longrightarrow \prod_{i=1}^{m-1} F(T_i), \quad \theta_2: F\left(\prod_{i=m}^\infty T_i\right) \longrightarrow \prod_{i=m}^\infty F(T_i).$$

This clearly implies that  $\text{Cone}(\theta) \cong \text{Cone}(\theta_1) \oplus \text{Cone}(\theta_2)$ . Since  $F$  (being additive) commutes with finite products,  $\theta_1$  is an isomorphism, whence  $\text{Cone}(\theta_1) \cong 0$ . On the other hand, condition (ii) tells us that when  $m \gg 0$ , the object  $\text{Cone}(\theta_2)$  will belong to  $\mathbf{D}^2(\mathcal{A})^{\geq n}$  with  $n$  arbitrarily large. Hence  $\text{Cone}(\theta) \cong \text{Cone}(\theta_2)$  must vanish, and  $\theta$  must be an isomorphism.  $\square$

Clearly, Proposition 1.7 has the following dual version whose proof simply consists in reducing to the previous result by passing to the opposite category.

**Proposition 1.8.** *Let  $\mathcal{A}$  be a small abelian category and  $F: \mathcal{T} \rightarrow \mathbf{D}^2(\mathcal{A})$  an additive functor, where  $? = -, \emptyset$ . Assume that  $\{T_i\}_{1 \leq i < \infty} \subseteq \text{Ob}(\mathcal{T})$  is such that*

- (i) *The coproduct  $\coprod_{i=1}^\infty T_i$  exists in  $\mathcal{T}$ .*
- (ii) *For every integer  $n > 0$ , there exists an integer  $m(n) > 0$  such that  $F\left(\coprod_{i=m(n)}^\infty T_i\right) \in \mathbf{D}(\mathcal{A})^{\leq -n}$ .*

*Then the coproduct  $\coprod_{i=1}^\infty F(T_i)$  exists in  $\mathbf{D}^2(\mathcal{A})$  and the canonical map*

$$\prod_{i=1}^\infty F(T_i) \longrightarrow F\left(\prod_{i=1}^\infty T_i\right)$$

*is an isomorphism.*

## 2. Generation

The key idea pursued in this paper is that uniqueness of dg enhancements is tightly related to suitable notions of generations. Those that are of interest in this paper are explained in this section.

**2.1. Well-generated triangulated categories**

Let  $\mathcal{T}$  be a triangulated category with small coproducts. For a cardinal  $\alpha$ , an object  $S$  of  $\mathcal{T}$  is  $\alpha$ -small if every map  $S \rightarrow \coprod_{i \in I} X_i$  in  $\mathcal{T}$  (where  $I$  is a small set) factors through  $\coprod_{i \in J} X_i$  for some  $J \subseteq I$  with  $|J| < \alpha$ . A cardinal  $\alpha$  is called *regular* if it is not the sum of fewer than  $\alpha$  cardinals, all of them smaller than  $\alpha$ .

**Definition 2.1.** The category  $\mathcal{T}$  is *well-generated* if there exists a small set  $\mathcal{S}$  of objects in  $\mathcal{T}$  satisfying the following properties:

- (G1) An object  $X \in \mathcal{T}$  is isomorphic to 0 if and only if  $\text{Hom}_{\mathcal{T}}(S, X[j]) = 0$  for all  $S \in \mathcal{S}$  and all  $j \in \mathbb{Z}$ .
- (G2) For every small set of maps  $\{X_i \rightarrow Y_i\}_{i \in I}$  in  $\mathcal{T}$ , the induced map  $\text{Hom}_{\mathcal{T}}(S, \coprod_i X_i) \rightarrow \text{Hom}_{\mathcal{T}}(S, \coprod_i Y_i)$  is surjective for all  $S \in \mathcal{S}$  if  $\text{Hom}_{\mathcal{T}}(S, X_i) \rightarrow \text{Hom}_{\mathcal{T}}(S, Y_i)$  is surjective for all  $i \in I$  and all  $S \in \mathcal{S}$ .
- (G3) There exists a regular cardinal  $\alpha$  such that every object of  $\mathcal{S}$  is  $\alpha$ -small.

**Remark 2.2.** The above notion was originally developed in [36]. Here we used the equivalent formulation in [24]. A nice survey on the subject is in [25].

Part of this paper is about enhancements of triangulated categories constructed out of Grothendieck categories. For the non-expert reader, let us recall that an abelian category  $\mathcal{G}$  is a Grothendieck category if it is closed under small coproducts, has a small set of generators  $\mathcal{S}$  and the direct limits of short exact sequences are exact in  $\mathcal{G}$ . The objects in  $\mathcal{S}$  are generators in the sense that for any  $C$  in  $\mathcal{G}$ , there exists an epimorphism  $S \twoheadrightarrow C$  in  $\mathcal{G}$ , where  $S$  is a small coproduct of objects in  $\mathcal{S}$ . By taking the coproduct of all generators in  $\mathcal{S}$ , we can assume that  $\mathcal{G}$  has a single generator  $G$ .

**Example 2.3.**

- (i) If  $X$  is an algebraic stack, the abelian categories  $\text{Mod}(\mathcal{O}_X)$  and  $\mathbf{Qcoh}(X)$  of  $\mathcal{O}_X$ -modules and quasi-coherent modules are Grothendieck categories.
- (ii) If  $\mathcal{A}$  is a small,  $\mathbb{k}$ -linear category, we denote by  $\text{Mod}(\mathcal{A})$  the Grothendieck category of additive functors  $\mathcal{A}^\circ \rightarrow \text{Mod}(\mathbb{k})$ . For later use, recall that if  $\alpha$  is a regular cardinal, then we denote by  $\text{Lex}_\alpha(\mathcal{A}^\circ, \text{Mod}(\mathbb{k}))$  the full subcategory of  $\text{Mod}(\mathcal{A})$  consisting of left exact functors that commute with  $\alpha$ -coproducts.
- (iii) If  $\mathcal{A}$  is an abelian category, we denote by  $\text{Ind}(\mathcal{A})$  its Ind-category (see [19, Section 8]), which is a Grothendieck category. Roughly, it is obtained from  $\mathcal{A}$  by formally adding filtered colimits of objects in  $\mathcal{A}$ .

The following states an important property for the derived category of a Grothendieck category.

**Theorem 2.4** ([33], Theorem 0.2). *If  $\mathcal{G}$  is a Grothendieck category, then  $\mathbf{D}(\mathcal{G})$  is well-generated.*

A full triangulated subcategory of  $\mathcal{T}$  is  $\alpha$ -localising if it is closed under  $\alpha$ -coproducts and under direct summands (the latter condition is automatic if  $\alpha > \aleph_0$ ). An  $\alpha$ -coproduct is a coproduct of strictly less than  $\alpha$  summands. A full subcategory of  $\mathcal{T}$  is *localising* if it is  $\alpha$ -localising for all regular cardinals  $\alpha$ .

When the category  $\mathcal{T}$  is well-generated and we want to put emphasis on the cardinal  $\alpha$  in (G3) and on  $\mathcal{S}$ , we say that  $\mathcal{T}$  is  $\alpha$ -well-generated by the set  $\mathcal{S}$ . In this situation, following [24], we denote by  $\mathcal{T}^\alpha$  the smallest  $\alpha$ -localising subcategory of  $\mathcal{T}$  containing  $\mathcal{S}$ . By [24, 36],  $\mathcal{T}^\alpha$  does not depend on the choice of the set  $\mathcal{S}$ , which well-generates  $\mathcal{T}$ .

Let  $\mathcal{G}$  be a Grothendieck category, and let  $\alpha$  be a sufficiently large regular cardinal. We are interested in describing the category  $\mathbf{D}(\mathcal{G})^\alpha$ . To this end, we denote by  $\mathcal{G}^\alpha$  the full subcategory of  $\mathcal{G}$  consisting of  $\alpha$ -presentable objects. An object  $G$  in  $\mathcal{G}$  is  $\alpha$ -presentable if the functor  $\text{Hom}_{\mathcal{G}}(G, -): \mathcal{G} \rightarrow \text{Mod}(\mathbb{k})$  preserves  $\alpha$ -filtered colimits (see, for example, [25, Section 6.4] for the definition of  $\alpha$ -filtered colimit).

**Theorem 2.5** ([23], Corollary 5.5 and Theorem 5.10). *Let  $\mathcal{G}$  be a Grothendieck category, and let  $\alpha$  be a sufficiently large regular cardinal.*

- (1) The category  $\mathcal{G}^\alpha$  is abelian.
- (2) There is a natural exact equivalence  $\mathbf{D}(\mathcal{G})^\alpha \cong \mathbf{D}(\mathcal{G}^\alpha)$ .

The objects in  $\mathcal{T}^\alpha$  are called  $\alpha$ -compact. Thus we will sometimes say that  $\mathcal{T}$  is  $\alpha$ -compactly generated by the set of  $\alpha$ -compact generators  $\mathcal{S}$ . A very interesting case is when  $\alpha = \aleph_0$ . Indeed, with this choice,  $\mathcal{T}^\alpha = \mathcal{T}^c$ , the full triangulated subcategory of compact objects in  $\mathcal{T}$ . Recall that an object  $C$  in  $\mathcal{T}$  is compact if the functor  $\mathcal{T}(C, -)$  commutes with small coproducts. Notice that the compact objects in  $\mathcal{T}$  are precisely the  $\aleph_0$ -small ones.

The analogue of Theorem 2.4 can be proven for the unseparated derived category.

**Theorem 2.6** ([23], Theorem 5.12). *If  $\mathcal{G}$  is a Grothendieck category, then  $\check{\mathbf{D}}(\mathcal{G})$  is well-generated.*

A weaker form of Theorem 2.5 (2) is also available. Indeed, by combining [23, Theorem 5.12(3)] with Theorem 2.5(2) for  $\alpha$  a sufficiently large regular cardinal, there is a quotient functor

$$\check{\mathbf{D}}(\mathcal{G})^\alpha \longrightarrow \mathbf{D}(\mathcal{G})^\alpha.$$

We will not use such a general result in this paper, but we will elaborate more on the following easier case.

**Example 2.7.** If  $\mathcal{A}$  is a small abelian category, then one takes the Ind-category  $\mathcal{G} := \text{Ind}(\mathcal{A})$  (see Example 2.3). By [23, Theorem 4.9], there is a natural exact equivalence  $\check{\mathbf{D}}(\mathcal{G})^c \cong \mathbf{D}^b(\mathcal{A})$ .

### 2.2. Generating derived categories

In the general case when  $\mathcal{A}$  is any abelian category, not necessarily Grothendieck, we need a different approach to the generation of  $\mathbf{D}^?( \mathcal{A} )$ .

Let us first recall the following rather general definition (see [7]).

**Definition 2.8.** Let  $\mathcal{T}$  be a triangulated category, and let  $\mathcal{S} \subset \text{Ob}(\mathcal{T})$ . We define

- (1)  $\langle \mathcal{S} \rangle_1$  is the collection of all direct summands of finite coproducts of shifts of objects in  $\mathcal{S}$ .
- (2)  $\langle \mathcal{S} \rangle_{n+1}$  consists of all direct summands of objects  $T \in \mathcal{T}$  for which there exists a distinguished triangle  $T_1 \rightarrow T \rightarrow T_2$  with  $T_1 \in \langle \mathcal{S} \rangle_n$  and  $T_2 \in \langle \mathcal{S} \rangle_1$ .

We set  $\langle \mathcal{S} \rangle_\infty$  for the full subcategory consisting of all objects  $T$  in  $\mathcal{T}$  contained in  $\langle \mathcal{S} \rangle_n$  for some  $n$ .

In our special case, we can prove the following.

**Proposition 2.9.** *Let  $\mathbf{V}^?( \mathcal{A} ) \subset \mathbf{K}^?( \mathcal{A} )$  be as defined in the opening paragraph of Subsection 1.1. For  $? = b, +, -, \emptyset$ , we have that  $\langle \mathbf{V}^?( \mathcal{A} ) \rangle_3 = \mathbf{K}^?( \mathcal{A} )$ .*

*Proof.* Let  $A^* \in \text{Ob}(\mathbf{K}^?( \mathcal{A} ) )$ , which we write as a complex

$$\dots \longrightarrow A^{-2} \longrightarrow A^{-1} \longrightarrow A^0 \longrightarrow A^1 \longrightarrow A^2 \longrightarrow \dots$$

Let  $K^i$  be the kernel of the differential  $A^i \rightarrow A^{i+1}$ . Then the map  $A^{i-1} \rightarrow A^i$  factors uniquely as  $A^{i-1} \xrightarrow{\alpha^i} K^i \hookrightarrow A^i$ .

This yields the morphism

$$\bigoplus_{i \in \mathbb{Z}} A^{i-1}[-i] \xrightarrow{\bigoplus_{i \in \mathbb{Z}} \alpha^i} \bigoplus_{i \in \mathbb{Z}} K^i[-i]$$

in  $\mathbf{V}^2(\mathcal{A})$ . Denote by  $C^*$  its mapping cone. It is clear that  $C^* \in \langle \mathbf{V}^2(\mathcal{A}) \rangle_2$ , and it is the direct sum over  $i \in \mathbb{Z}$  of the complexes

$$\dots \longrightarrow 0 \longrightarrow A^{i-1} \xrightarrow{\alpha^i} K^i \longrightarrow 0 \longrightarrow \dots$$

Now consider the cochain map

$$\bigoplus_{i \in \mathbb{Z}} K^i[-i] \xrightarrow{\varphi + \psi} C^*$$

whose components, out of  $K^i[-i]$ , are (respectively)  $\varphi^i$  as below

$$\begin{array}{cccccccccccc} \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & K^i & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & \downarrow & & \downarrow & & \parallel & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & 0 & \longrightarrow & A^{i-1} & \longrightarrow & K^i & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \end{array} \tag{2.1}$$

and  $\psi^i$  as below

$$\begin{array}{cccccccccccc} \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & K^i & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & A^i & \longrightarrow & K^{i+1} & \longrightarrow & 0 & \longrightarrow & \dots \end{array} \tag{2.2}$$

It can be easily checked that the mapping cone of the morphism  $\varphi + \psi$  is isomorphic to the direct sum of the complex  $A^*$  and of complexes of the form

$$\dots \longrightarrow 0 \longrightarrow K^i \longrightarrow K^i \longrightarrow 0 \longrightarrow \dots$$

In other words,  $\text{Cone}(\varphi + \psi) \cong A^*$  in  $\mathbf{K}^2(\mathcal{A})$ . Therefore, as  $C^*$  belongs to  $\langle \mathbf{V}^2(\mathcal{A}) \rangle_2$  and  $\varphi + \psi$  is a morphism from an object of  $\mathbf{V}^2(\mathcal{A})$  to  $C^*$ , we have that  $A^* \in \langle \mathbf{V}^2(\mathcal{A}) \rangle_3$ .  $\square$

The reader might wish to compare the proof of Proposition 2.9 above with the proof of [41, Proposition 7.22]; there are similarities.

As a straightforward consequence of Proposition 2.9 and of the fact that  $\mathbf{D}^2(\mathcal{A})$  is a quotient of  $\mathbf{K}^2(\mathcal{A})$ , as explained in Section 1.1, we obtain the following.

**Corollary 2.10.** *Let  $\mathbf{B}^2(\mathcal{A}) \subset \mathbf{D}^2(\mathcal{A})$  be as defined in the paragraph between Remarks 1.2 and 1.1. For  $? = b, +, -, \emptyset$ , we have that  $\langle \mathbf{B}^2(\mathcal{A}) \rangle_3 = \mathbf{D}^2(\mathcal{A})$ .*

### 3. Dg categories and enhancements

We briefly introduce dg categories and some basic machinery in Section 3.1. Next, we describe some constructions that will be important in the rest of the paper: Drinfeld quotients and h-flat resolutions (Section 3.2), the model structure and homotopy pullbacks (Section 3.3) and, finally, localisations for

dg categories (Section 3.4). Dg enhancements, their uniqueness and dependence on the universe where the categories are defined are the contents of Section 3.5.

### 3.1. Dg categories

A dg category is a  $\mathbb{k}$ -linear category  $\mathcal{C}$  such that the morphism spaces  $\text{Hom}_{\mathcal{C}}(A, B)$  are complexes of  $\mathbb{k}$ -modules and the composition maps  $\text{Hom}_{\mathcal{C}}(B, C) \otimes_{\mathbb{k}} \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{C}}(A, C)$  are morphisms of complexes for all  $A, B, C$  in  $\text{Ob}(\mathcal{C})$ .

A dg functor  $F: \mathcal{C}_1 \rightarrow \mathcal{C}_2$  between two dg categories is a  $\mathbb{k}$ -linear functor such that the maps  $\text{Hom}_{\mathcal{C}_1}(A, B) \rightarrow \text{Hom}_{\mathcal{C}_2}(F(A), F(B))$  are morphisms of complexes for all  $A, B$  in  $\text{Ob}(\mathcal{C}_1)$ .

The underlying category  $Z^0(\mathcal{C})$  (respectively, the homotopy category  $H^0(\mathcal{C})$ ) of a dg category  $\mathcal{C}$  is the  $\mathbb{k}$ -linear category with the same objects of  $\mathcal{C}$  and such that  $\text{Hom}_{Z^0(\mathcal{C})}(A, B) := Z^0(\text{Hom}_{\mathcal{C}}(A, B))$  (respectively,  $\text{Hom}_{H^0(\mathcal{C})}(A, B) := H^0(\text{Hom}_{\mathcal{C}}(A, B))$ ) for all  $A, B$  in  $\text{Ob}(\mathcal{C})$  (with composition of morphisms naturally induced from the one in  $\mathcal{C}$ ). Two objects of  $\mathcal{C}$  are dg isomorphic (respectively, homotopy equivalent) if they are isomorphic in  $Z^0(\mathcal{C})$  (respectively,  $H^0(\mathcal{C})$ ). One can also define  $Z(\mathcal{C})$  (respectively  $H(\mathcal{C})$ ) to be the graded (namely, dg with trivial differential) category whose objects are the same as those of  $\mathcal{C}$ , while  $\text{Hom}_{Z(\mathcal{C})}(A, B) := \bigoplus_{i \in \mathbb{Z}} Z^i(\text{Hom}_{\mathcal{C}}(A, B))$  (respectively,  $\text{Hom}_{H(\mathcal{C})}(A, B) := \bigoplus_{i \in \mathbb{Z}} H^i(\text{Hom}_{\mathcal{C}}(A, B))$ ) for all  $A, B$  in  $\text{Ob}(\mathcal{C})$ .

**Example 3.1.** If  $\mathcal{A}$  is a  $\mathbb{k}$ -linear category, there is a natural dg category  $\mathbf{C}_{\text{dg}}^?( \mathcal{A} )$  such that  $H^0(\mathbf{C}_{\text{dg}}^?( \mathcal{A} )) = \mathbf{K}^?( \mathcal{A} )$  for  $? = b, +, -, \emptyset$ . Explicitly,

$$\text{Hom}_{\mathbf{C}_{\text{dg}}^?( \mathcal{A} )}(A^*, B^*)^n := \prod_{i \in \mathbb{Z}} \text{Hom}_{\mathcal{A}}(A^i, B^{n+i})$$

for every  $A^*, B^* \in \text{Ob}(\mathbf{C}_{\text{dg}}^?( \mathcal{A} ))$  and for every  $n \in \mathbb{Z}$ . While the composition of morphisms is the obvious one, the differential is defined on a homogeneous element  $f \in \text{Hom}_{\mathbf{C}_{\text{dg}}^?( \mathcal{A} )}(A^*, B^*)^n$  by  $d(f) := d_B \circ f - (-1)^n f \circ d_A$ .

A dg functor  $F: \mathcal{C}_1 \rightarrow \mathcal{C}_2$  induces a  $\mathbb{k}$ -linear functor  $H^0(F): H^0(\mathcal{C}_1) \rightarrow H^0(\mathcal{C}_2)$ . We say that  $F$  is a quasi-equivalence if the maps  $\text{Hom}_{\mathcal{C}_1}(A, B) \rightarrow \text{Hom}_{\mathcal{C}_2}(F(A), F(B))$  are quasi-isomorphisms for all  $A, B$  in  $\text{Ob}(\mathcal{C}_1)$ , and  $H^0(F)$  is an equivalence.

If  $\mathbb{U}$  is a universe, we denote by  $\mathbf{dgCat}_{\mathbb{U}}$  (or simply by  $\mathbf{dgCat}$ , if there can be no ambiguity about  $\mathbb{U}$ ) the category whose objects are  $\mathbb{U}$ -small dg categories and whose morphisms are dg functors. It is known (see [44]) that  $\mathbf{dgCat}$  has a model structure whose weak equivalences are quasi-equivalences and such that every object is fibrant. We denote by  $\mathbf{Hqe}$  (or  $\mathbf{Hqe}_{\mathbb{U}}$ , if needed) the corresponding homotopy category, namely the localisation of  $\mathbf{dgCat}$  with respect to quasi-equivalences. Since  $H^0$  sends quasi-equivalences to equivalences, for every morphism  $f: \mathcal{C}_1 \rightarrow \mathcal{C}_2$  in  $\mathbf{Hqe}$  there is a  $\mathbb{k}$ -linear functor  $H^0(f): H^0(\mathcal{C}_1) \rightarrow H^0(\mathcal{C}_2)$ , which is well-defined up to isomorphism.

Dg functors between two dg categories  $\mathcal{C}_1$  and  $\mathcal{C}_2$  form in a natural way the objects of a dg category  $\underline{\text{Hom}}(\mathcal{C}_1, \mathcal{C}_2)$ . For every dg category  $\mathcal{C}$ , we set  $\text{dgMod}(\mathcal{C}) := \underline{\text{Hom}}(\mathcal{C}^\circ, \mathbf{C}_{\text{dg}}(\text{Mod}(\mathbb{k})))$  and call its objects (right) dg  $\mathcal{C}$ -modules. Observe that  $\text{dgMod}(\mathbb{k})$  can be identified with  $\mathbf{C}_{\text{dg}}(\text{Mod}(\mathbb{k}))$ .

For every dg category  $\mathcal{C}$ , the map defined on objects by  $A \mapsto \text{Hom}_{\mathcal{C}}(-, A)$  extends to a fully faithful dg functor  $Y_{\text{dg}}^{\mathcal{C}}: \mathcal{C} \rightarrow \text{dgMod}(\mathcal{C})$  (the dg Yoneda embedding). It is easy to see that the image of  $Y_{\text{dg}}^{\mathcal{C}}$  is contained in the full dg subcategory  $\text{h-proj}(\mathcal{C})$  of  $\text{dgMod}(\mathcal{C})$  whose objects are  $h$ -projective dg  $\mathcal{C}$ -modules. By definition,  $M \in \text{Ob}(\text{dgMod}(\mathcal{C}))$  is  $h$ -projective if  $\text{Hom}_{H^0(\text{dgMod}(\mathcal{C}))}(M, N) = 0$  for every  $N \in \text{Ob}(\text{dgAcy}(\mathcal{C}))$ , where  $\text{dgAcy}(\mathcal{C})$  is the full dg subcategory of  $\text{dgMod}(\mathcal{C})$  whose objects are acyclic (in the sense that  $N(A)$  is an acyclic complex for every  $A \in \text{Ob}(\mathcal{C})$ ).

If  $F: \mathcal{C}_1 \rightarrow \mathcal{C}_2$  is a dg functor, composition with  $F^\circ$  yields a dg functor  $\text{Res}(F): \text{dgMod}(\mathcal{C}_2) \rightarrow \text{dgMod}(\mathcal{C}_1)$ . It turns out that there also exists a dg functor  $\text{Lnd}(F): \text{dgMod}(\mathcal{C}_1) \rightarrow \text{dgMod}(\mathcal{C}_2)$ , which is left adjoint to  $\text{Res}(F)$  and such that  $\text{Lnd}(F) \circ Y_{\text{dg}}^{\mathcal{C}_1} \cong Y_{\text{dg}}^{\mathcal{C}_2} \circ F$ . Moreover,  $\text{Lnd}(F)$  preserves  $h$ -projective dg modules, and  $\text{Lnd}(F): \text{h-proj}(\mathcal{C}_1) \rightarrow \text{h-proj}(\mathcal{C}_2)$  is a quasi-equivalence if  $F$  is. This last fact clearly

implies that a(n iso)morphism  $f: \mathcal{C}_1 \rightarrow \mathcal{C}_2$  in **Hqe** induces a(n iso)morphism  $\text{Ind}(f): \text{h-proj}(\mathcal{C}_1) \rightarrow \text{h-proj}(\mathcal{C}_2)$  in **Hqe**.

As is explained, for instance, in [6], there is a notion of formal shift by an integer  $n$  of an object  $A$  in a dg category  $\mathcal{C}$  (denoted, as usual, by  $A[n]$ ). Similarly, one can define the formal cone of a morphism  $f$  in  $Z^0(\mathcal{C})$  (denoted, as usual, by  $\text{Cone}(f)$ ). Shifts and cones need not exist in an arbitrary dg category, but when they do, they are unique up to dg isomorphism and are preserved by dg functors. The following property of the cone of a morphism will be useful later.

**Lemma 3.2.** *Let  $A \xrightarrow{f} B \xrightarrow{g} C$  be morphisms in  $Z^0(\mathcal{C})$  such that  $\text{Cone}(f)$  exists and  $g \circ f$  is a coboundary. Then there exists  $h: \text{Cone}(f) \rightarrow C$  in  $Z^0(\mathcal{C})$  such that  $g = h \circ j$ , where  $j: B \rightarrow \text{Cone}(f)$  is the natural morphism.*

*Proof.* See [18, Proposition 2.3.4]. □

**Definition 3.3.** A dg category  $\mathcal{C}$  is *strongly pretriangulated* if  $A[n]$  and  $\text{Cone}(f)$  exist (in  $\mathcal{C}$ ) for every  $n \in \mathbb{Z}$ , every object  $A$  of  $\mathcal{C}$  and every morphism  $f$  of  $Z^0(\mathcal{C})$ .

A dg category  $\mathcal{C}$  is *pretriangulated* if there exists a quasi-equivalence  $\mathcal{C} \rightarrow \mathcal{C}'$  with  $\mathcal{C}'$  strongly pretriangulated.

**Remark 3.4.** If  $\mathcal{C}$  is a pretriangulated dg category, then  $H^0(\mathcal{C})$  is a triangulated category in a natural way. If  $f$  is a morphism in **Hqe** between two pretriangulated dg categories, then the functor  $H^0(f)$  is exact.

If  $\mathcal{C}$  is a dg category,  $\text{dgMod}(\mathcal{C})$ ,  $\text{dgAcy}(\mathcal{C})$  and  $\text{h-proj}(\mathcal{C})$  are strongly pretriangulated dg categories. Moreover, the (triangulated) categories  $H^0(\text{dgMod}(\mathcal{C}))$ ,  $H^0(\text{dgAcy}(\mathcal{C}))$  and  $H^0(\text{h-proj}(\mathcal{C}))$  have arbitrary coproducts, and there is a semi-orthogonal decomposition

$$H^0(\text{dgMod}(\mathcal{C})) = \langle H^0(\text{dgAcy}(\mathcal{C})), H^0(\text{h-proj}(\mathcal{C})) \rangle. \tag{3.1}$$

This clearly implies that there is an exact equivalence between  $H^0(\text{h-proj}(\mathcal{C}))$  and the Verdier quotient  $\mathcal{D}(\mathcal{C}) := H^0(\text{dgMod}(\mathcal{C}))/H^0(\text{dgAcy}(\mathcal{C}))$  (which is by definition the *derived category* of  $\mathcal{C}$ ).

For every dg category  $\mathcal{C}$ , we will denote by  $\text{Pretr}(\mathcal{C})$  (respectively,  $\text{Perf}(\mathcal{C})$ ) the smallest full dg subcategory of  $\text{h-proj}(\mathcal{C})$  containing  $Y_{\text{dg}}^{\mathcal{C}}(\mathcal{C})$  and closed under homotopy equivalences, shifts, cones (respectively, also direct summands in  $H^0(\text{h-proj}(\mathcal{C}))$ ). It is easy to see that  $\text{Pretr}(\mathcal{C})$  and  $\text{Perf}(\mathcal{C})$  are strongly pretriangulated and that  $\mathcal{C}$  is pretriangulated if and only if  $Y_{\text{dg}}^{\mathcal{C}}: \mathcal{C} \rightarrow \text{Pretr}(\mathcal{C})$  is a quasi-equivalence. Moreover,  $\text{Pretr}(\mathcal{C}) \subseteq \text{Perf}(\mathcal{C})$  and  $H^0(\text{Perf}(\mathcal{C}))$  can be identified with the idempotent completion  $H^0(\text{Pretr}(\mathcal{C}))^{\text{ic}}$  of  $H^0(\text{Pretr}(\mathcal{C}))$ . Hence  $Y_{\text{dg}}^{\mathcal{C}}: \mathcal{C} \rightarrow \text{Perf}(\mathcal{C})$  is a quasi-equivalence if and only if  $\mathcal{C}$  is pretriangulated and  $H^0(\mathcal{C})$  is idempotent complete.

**Remark 3.5.** Recall that an additive category  $\mathcal{A}$  is *idempotent complete* if every idempotent (namely, a morphism  $e: A \rightarrow A$  in  $\mathcal{A}$  such that  $e^2 = e$ ) splits or, equivalently, has a kernel. Every additive category  $\mathcal{A}$  admits a fully faithful and additive embedding  $\mathcal{A} \hookrightarrow \mathcal{A}^{\text{ic}}$ , where  $\mathcal{A}^{\text{ic}}$  is an idempotent complete additive category, with the property that every object of  $\mathcal{A}^{\text{ic}}$  is a direct summand of an object from  $\mathcal{A}$ . The category  $\mathcal{A}^{\text{ic}}$  (or, better, the functor  $\mathcal{A} \rightarrow \mathcal{A}^{\text{ic}}$ ) is called the *idempotent completion* of  $\mathcal{A}$ . It can be proved (see [3]) that if  $\mathcal{T}$  is a triangulated category, then  $\mathcal{T}^{\text{ic}}$  is triangulated as well (and  $\mathcal{T} \hookrightarrow \mathcal{T}^{\text{ic}}$  is exact).

If  $\mathcal{C}$  is a dg category, then  $H^0(\text{Perf}(\mathcal{C}))$  is idempotent complete, and from this it is easy to deduce that  $Z^0(\text{Perf}(\mathcal{C}))$  is also idempotent complete.

If  $\mathcal{A}$  is an abelian category, it follows from [3, 43] that  $\mathbf{D}^?( \mathcal{A} )$  is idempotent complete for  $? = b, +, -, \emptyset$ . More precisely, for  $? \in \{-, +\}$ , the result may be found in [3, Lemma 2.4]; for  $? = b$ , see [3, Lemma 2.8]; and for  $? = \emptyset$ , see Theorem 6 of Section 10 in [43] combined with Lemma 7 of Section 9 in the same paper.

Observe that by Remark 1.3 combined with the paragraph above, the categories  $\mathbf{K}^?( \mathcal{A} )$  are also idempotent complete—as long as  $\mathcal{A}$  is abelian and with  $? = b, +, -, \emptyset$ .



### 3.2. Drinfeld quotients and h-flat resolutions

Let  $\mathcal{C}$  be a dg category and  $\mathcal{D} \subseteq \mathcal{C}$  a full dg subcategory. As explained in [16, Section 3.1], one can form the *Drinfeld quotient* of  $\mathcal{C}$  by  $\mathcal{D}$ , which we denote by  $\mathcal{C}/\mathcal{D}$ . This is a dg category, and its construction goes roughly as follows: given  $D \in \text{Ob}(\mathcal{D})$ , we formally add a morphism  $f_D: D \rightarrow D$  of degree  $-1$ , and we set  $d(f_D) = \text{id}_D$ .

If  $\mathcal{C}$  is pretriangulated and  $\mathcal{D}$  is a full pretriangulated dg subcategory, then  $\mathcal{C}/\mathcal{D}$  is pretriangulated. In this case, the natural dg functor  $\mathcal{C} \rightarrow \mathcal{C}/\mathcal{D}$  induces an exact functor  $H^0(\mathcal{C}) \rightarrow H^0(\mathcal{C}/\mathcal{D})$ , which sends to zero the objects of  $\mathcal{D}$ . Thus it factors through the Verdier quotient  $H^0(\mathcal{C}) \rightarrow H^0(\mathcal{C})/H^0(\mathcal{D})$ , yielding an exact functor

$$H^0(\mathcal{C})/H^0(\mathcal{D}) \longrightarrow H^0(\mathcal{C}/\mathcal{D}), \tag{3.2}$$

which need not be an equivalence, in general.

**Definition 3.6.** We remind the reader of the terminology of [16]. A dg category  $\mathcal{C}$  is *h-flat* if, for all  $C_1, C_2 \in \text{Ob}(\mathcal{C})$ , the complex  $\text{Hom}_{\mathcal{C}}(C_1, C_2)$  is homotopically flat over  $\mathbb{k}$ . The homotopic flatness of  $\text{Hom}_{\mathcal{C}}(C_1, C_2)$  means for any acyclic complex  $M$  of  $\mathbb{k}$ -modules,  $\text{Hom}_{\mathcal{C}}(C_1, C_2) \otimes_{\mathbb{k}} M$  is acyclic.

**Example 3.7.** If  $\mathbb{k}$  is a field, then every dg category is clearly h-flat.

As a special case of [16, Theorem 3.4], we have that if  $\mathcal{C}$  is an h-flat pretriangulated dg category and  $\mathcal{D}$  is a full pretriangulated subcategory of  $\mathcal{C}$ , then equation (3.2) is an exact equivalence.

If  $\mathcal{C}$  is not h-flat, Drinfeld shows in [16, Lemma B.5] that one can construct an h-flat dg category  $\tilde{\mathcal{C}}$  with a quasi-equivalence  $l_{\mathcal{C}}: \tilde{\mathcal{C}} \rightarrow \mathcal{C}$ . One can then define  $\tilde{\mathcal{D}}$  to be the full dg subcategory  $l_{\mathcal{C}}^{-1}\mathcal{D} \subset \tilde{\mathcal{C}}$  and take the morphism  $q \in \text{Hom}_{\mathbf{Hqcat}}(\mathcal{C}, \tilde{\mathcal{C}}/\tilde{\mathcal{D}})$  represented as

$$\begin{array}{ccc} & \tilde{\mathcal{C}} & \\ l_{\mathcal{C}} \swarrow & & \searrow \\ \mathcal{C} & & \tilde{\mathcal{C}}/\tilde{\mathcal{D}}, \end{array}$$

where the dg functor on the right is the natural one mentioned above.

This construction satisfies the following universal property, which is a special instance of [16, Main Theorem]. Assume that  $\mathcal{C}'$  is a pretriangulated dg category and  $f \in \text{Hom}_{\mathbf{Hqcat}}(\mathcal{C}, \mathcal{C}')$  is such that  $H^0(f)$  sends the objects of  $\mathcal{D}$  to zero. Then there is a unique  $\bar{f} \in \text{Hom}_{\mathbf{Hqcat}}(\tilde{\mathcal{C}}/\tilde{\mathcal{D}}, \mathcal{C}')$  making the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{q} & \tilde{\mathcal{C}}/\tilde{\mathcal{D}} \\ & \searrow f & \downarrow \bar{f} \\ & & \mathcal{C}' \end{array} \tag{3.3}$$

commute in  $\mathbf{Hqcat}$ .

In the rest of this section, we describe two variants of  $\tilde{\mathcal{C}}$  with properties that we will need in the rest of the paper. We start with a dg category  $\mathcal{C}$ , we let  $\mathcal{T}$  be the graded category  $\mathcal{T} = H(\mathcal{C})$ , and the aim is to produce two sequences of dg categories and faithful dg functors

$$\mathcal{C}_0 \hookrightarrow \mathcal{C}_1 \hookrightarrow \mathcal{C}_2 \hookrightarrow \mathcal{C}_3 \hookrightarrow \dots, \quad \mathcal{C}'_0 \hookrightarrow \mathcal{C}'_1 \hookrightarrow \mathcal{C}'_2 \hookrightarrow \mathcal{C}'_3 \hookrightarrow \dots,$$

together with compatible dg functors  $l_n: \mathcal{C}_n \rightarrow \mathcal{C}$  and  $l'_n: \mathcal{C}'_n \rightarrow \mathcal{C}$ , which are the identity on objects. Then we set  $\mathcal{C}^{\text{hf}}$  and  $\mathcal{C}^{\text{sm}}$  to be the respective colimits, with the induced dg functors  $l_{\mathcal{C}}^{\text{hf}}: \mathcal{C}^{\text{hf}} \xrightarrow{\sim} \mathcal{C}$  and  $l_{\mathcal{C}}^{\text{sm}}: \mathcal{C}^{\text{sm}} \xrightarrow{\sim} \mathcal{C}$ .

We define  $\mathcal{C}_0 = \mathcal{C}'_0$  to be the discrete  $\mathbb{k}$ -linear category with the same objects as  $\mathcal{C}$ . This means

$$\text{Hom}_{\mathcal{C}_0}(A, B) := \begin{cases} \mathbb{k} \cdot \text{id}_A & \text{if } A = B \\ 0 & \text{otherwise.} \end{cases}$$

The dg functor  $l_0 = l'_0$  is the obvious one that acts as the identity on objects and morphisms.

For  $n = 1$ , we set

$$D^1_{\mathcal{C}}(A, B) := \{(f, 0) \in \text{Hom}_{\mathcal{C}}(A, B) \times \text{Hom}_{\mathcal{C}_0}(A, B) : d(f) = 0\}.$$

Now the composite

$$D^1_{\mathcal{C}}(A, B) \longrightarrow Z(\text{Hom}_{\mathcal{C}}(A, B)) \longrightarrow H(\text{Hom}_{\mathcal{C}}(A, B)) \xrightarrow{\sim} \text{Hom}_{\mathcal{T}}(A, B)$$

is surjective by construction; hence we may choose a splitting. We let  $\overline{D}^1_{\mathcal{C}}(A, B) \subset D^1_{\mathcal{C}}(A, B)$  be a subset such that the composite  $\overline{D}^1_{\mathcal{C}}(A, B) \rightarrow D^1_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{T}}(A, B)$  is an isomorphism. And we define  $\mathcal{C}_1$  so that  $\text{Hom}_{\mathcal{C}_1}(A, B)$  is the graded  $\mathbb{k}$ -module freely generated by the basis  $D^1_{\mathcal{C}}(A, B) - \{0\}$ , with the composition being obvious on basis vectors. And  $\mathcal{C}'_1$  is the graded  $\mathbb{k}$ -linear category freely generated<sup>1</sup> over  $\mathcal{C}_0$  by the sets  $\overline{D}^1_{\mathcal{C}}(A, B)$ . The differentials of  $\mathcal{C}_1$  and  $\mathcal{C}'_1$  are trivial.

We continue for  $n \geq 2$  by defining inductively, for all  $A, B \in \text{Ob}(\mathcal{C})$ ,

$$D^n_{\mathcal{C}}(A, B) := \{(f, b) \in \text{Hom}_{\mathcal{C}}(A, B) \times \text{Hom}_{\mathcal{C}_{n-1}}(A, B) : d(f) = l_{n-1}(b)\}.$$

The definition of  $\overline{D}^n_{\mathcal{C}}(A, B)$  is slightly more delicate. We begin by copying the procedure above with  $\mathcal{C}'_{n-1}$  in place of  $\mathcal{C}_{n-1}$ , setting

$$\widehat{D}^n_{\mathcal{C}}(A, B) := \{(f, b) \in \text{Hom}_{\mathcal{C}}(A, B) \times \text{Hom}_{\mathcal{C}'_{n-1}}(A, B) : d(f) = l'_{n-1}(b)\}.$$

And then we observe that  $\widehat{D}^n_{\mathcal{C}}(A, B)$  surjects to the kernel of the surjective map

$$Z(\text{Hom}_{\mathcal{C}'_{n-1}}(A, B)) \longrightarrow \text{Hom}_{\mathcal{T}}(A, B),$$

allowing us to choose a subset  $\overline{D}^n_{\mathcal{C}}(A, B) \subset \widehat{D}^n_{\mathcal{C}}(A, B)$  so that the composite

$$\overline{D}^n_{\mathcal{C}}(A, B) \longrightarrow \widehat{D}^n_{\mathcal{C}}(A, B) \longrightarrow \text{Ker}\left(Z(\text{Hom}_{\mathcal{C}'_{n-1}}(A, B)) \rightarrow \text{Hom}_{\mathcal{T}}(A, B)\right) \quad (3.4)$$

is an isomorphism. And now we are ready:  $\mathcal{C}_n$  is the graded  $\mathbb{k}$ -linear category freely generated over  $\mathcal{C}_{n-1}$  by the sets  $D^n_{\mathcal{C}}(A, B)$ , while  $\mathcal{C}'_n$  is the graded  $\mathbb{k}$ -linear category freely generated over  $\mathcal{C}'_{n-1}$  by the sets  $\overline{D}^n_{\mathcal{C}}(A, B)$ . It is easy to show that  $\mathcal{C}_n, \mathcal{C}'_n$  become dg categories if we extend the differential on  $\mathcal{C}_{n-1}$  by sending a generator  $(f, b)$ , in either  $D_n(A, B)$  or  $\overline{D}^n_{\mathcal{C}}(A, B)$ , to  $b$ .

For  $n \geq 1$ , the dg functor  $l_n : \mathcal{C}_n \rightarrow \mathcal{C}$  is determined by setting  $l_n|_{\mathcal{C}_{n-1}} := l_{n-1}$  and  $l_n((f, b)) := f$  for all  $(f, b) \in D^n_{\mathcal{C}}(A, B)$ . Similarly, the dg functor  $l'_n : \mathcal{C}'_n \rightarrow \mathcal{C}$  is given by setting  $l'_n|_{\mathcal{C}'_{n-1}} := l'_{n-1}$  and  $l'_n((f, b)) := f$  for all  $(f, b) \in \overline{D}^n_{\mathcal{C}}(A, B)$ .

The following are easy consequences of the definitions.

<sup>1</sup>The notion of graded ( $\mathbb{k}$ -linear) category that is freely generated over another graded category, which we use here, is the same as that in [16, Lemma B.5] (see also [16, Section 3.1]).

**Lemma 3.8.** *If the category  $\mathcal{T}$  is  $\mathbb{U}$ -small, if  $\mathbb{U} \in \mathbb{V}$  is a larger universe and if  $\mathcal{C}$  is a  $\mathbb{V}$ -small dg category with  $H(\mathcal{C}) \cong \mathcal{T}$ , then the dg functor  $\mathbb{1}^{\text{sm}} : \mathcal{C}^{\text{sm}} \rightarrow \mathcal{C}$  is a quasi-equivalence with  $\mathcal{C}^{\text{sm}}$  a  $\mathbb{U}$ -small dg category. Moreover, if  $\mathcal{C}$  is pretriangulated, then so is  $\mathcal{C}^{\text{sm}}$ .*

*Proof.* Obvious by construction, the passage from  $\mathcal{C}'_{n-1}$  to  $\mathcal{C}'_n$  keeps tight control of the size of sets that come up (see, in particular, equation (3.4)). □

**Remark 3.9.** It is easy to see that if  $\mathcal{C}$  is pretriangulated (closed by shifts is enough), then  $H(\mathcal{C})$  is  $\mathbb{U}$ -small if and only if  $H^0(\mathcal{C})$  is  $\mathbb{U}$ -small.

**Proposition 3.10.** *The construction taking a dg category  $\mathcal{C}$  to the dg functor  $\mathbb{1}^{\text{hf}} : \mathcal{C}^{\text{hf}} \rightarrow \mathcal{C}$  satisfies the following properties:*

- (1) *For every dg category  $\mathcal{C}$ , the dg functor  $\mathbb{1}^{\text{hf}}$  is a quasi-equivalence and  $\mathcal{C}^{\text{hf}}$  is h-flat.*
- (2)  *$(-)^{\text{hf}}$  is a functor from the category of dg categories to itself, and  $\mathbb{1}^{\text{hf}} : (-)^{\text{hf}} \rightarrow \text{id}$  is a natural transformation. This means if  $\mathcal{C}$  and  $\mathcal{D}$  are dg categories and  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a dg functor, there is a canonically defined dg functor  $F^{\text{hf}} : \mathcal{C}^{\text{hf}} \rightarrow \mathcal{D}^{\text{hf}}$  making the following diagram commutative:*

$$\begin{array}{ccc}
 \mathcal{C}^{\text{hf}} & \xrightarrow{F^{\text{hf}}} & \mathcal{D}^{\text{hf}} \\
 \mathbb{1}^{\text{hf}} \downarrow & & \downarrow \mathbb{1}^{\text{hf}} \\
 \mathcal{C} & \xrightarrow{F} & \mathcal{D}.
 \end{array}$$

*Moreover, the construction taking  $F : \mathcal{C} \rightarrow \mathcal{D}$  to  $F^{\text{hf}} : \mathcal{C}^{\text{hf}} \rightarrow \mathcal{D}^{\text{hf}}$  respects composition and identities.*

- (3) *If  $\mathcal{C}$  is pretriangulated, then  $\mathcal{C}^{\text{hf}}$  is pretriangulated.*

*Proof.* Part (1) has the same proof as in [16, Lemma B.5] since properties (i)–(iv) in the construction of [16, Lemma B.5] are obviously satisfied by our construction as well (note that (iii) is verified starting from  $n = 2$ ). The functoriality and naturality in (2) are clear; the construction is choice-free. Indeed,  $F$  induces a natural dg functor  $F_0 : \mathcal{C}_0 \rightarrow \mathcal{D}_0$  acting as  $F$  on the objects and as the identity on the morphisms. Similarly,  $F$  induces a dg functor  $F_n : \mathcal{C}_n \rightarrow \mathcal{D}_n$  extending  $F_{n-1}$  and determined by the assignment

$$(f, b) \in D_{\mathcal{C}}^n(A, B) \mapsto (F(f), F_{n-1}(b)) \in D_{\mathcal{D}}^n(F(A), F(B)).$$

We set  $F^{\text{hf}}$  to be the colimit of the dg functors  $F_n$ . Part (3) is easy from (1). □

Suppose now that  $\mathcal{C}$  is a dg category and  $\mathcal{D} \subseteq \mathcal{C}$  is a full dg subcategory. Moreover, let  $\mathcal{D}'$  be the full dg subcategory of  $\mathcal{C}^{\text{hf}}$  with the same objects as  $\mathcal{D}$ . When  $\mathcal{C}$  and  $\mathcal{D}$  are pretriangulated, there are natural exact equivalences  $H^0(\mathcal{C})/H^0(\mathcal{D}) \cong H^0(\mathcal{C}^{\text{hf}})/H^0(\mathcal{D}') \cong H^0(\mathcal{C}^{\text{hf}}/\mathcal{D}')$ . Furthermore, not only are  $\tilde{\mathcal{C}}$  and  $\mathcal{C}^{\text{hf}}$  isomorphic in **Hqe**, but by the universal property of Drinfeld quotients mentioned above, the dg categories  $\tilde{\mathcal{C}}/\mathcal{D}$  and  $\mathcal{C}^{\text{hf}}/\mathcal{D}'$  are isomorphic in **Hqe**.

**Remark 3.11.** Now let  $\mathcal{C}$  be a pretriangulated dg category and  $\mathcal{D}$  a full pretriangulated dg subcategory of  $\mathcal{C}$ . Denoting by  $\mathbb{Q} : \mathcal{C}^{\text{hf}} \rightarrow \mathcal{C}^{\text{hf}}/\mathcal{D}'$  the natural dg functor, assume that we have a dg functor  $F : \mathcal{C}^{\text{hf}} \rightarrow \mathcal{C}'$  of pretriangulated dg categories such that  $H^0(F)$  sends the objects of  $\mathcal{D}'$  to zero. Then the universal property pictured in equation (3.3) can be made more explicit: there exists a dg functor  $\bar{F} : \mathcal{C}^{\text{hf}}/\mathcal{D}' \rightarrow \mathcal{C}'$  such that the diagram

$$\begin{array}{ccc}
 \mathcal{C}^{\text{hf}} & \xrightarrow{\mathbb{Q}} & \mathcal{C}^{\text{hf}}/\mathcal{D}' \\
 & \searrow F & \downarrow \bar{F} \\
 & & \mathcal{C}'
 \end{array}$$

is commutative in **dgCat**. The existence of  $\bar{F}$  is simple enough to see from the construction of  $\mathcal{E}^{\text{hf}}/\mathcal{D}'$  and  $\mathbf{Q}$  in [16, Section 3.1]. Indeed, as  $\mathcal{E}^{\text{hf}}/\mathcal{D}'$  has the same objects as  $\mathcal{E}^{\text{hf}}$ , one sets  $\bar{F}(C) := F(C)$  for all  $C$  in  $\mathcal{E}^{\text{hf}}/\mathcal{D}'$ . If  $D \in \mathcal{D}'$  is an object, then  $\text{Hom}_{\mathcal{E}'}(F(D), F(D))$  is acyclic, allowing us to choose a degree -1 morphism  $f(D): F(D) \rightarrow F(D)$  with  $d(f(D)) = \text{id}$ . Then the dg functor  $\bar{F}$  extends  $F$  on morphisms and takes each degree -1 morphism  $f_D: D \rightarrow D$ , in the definition of the category  $\mathcal{E}^{\text{hf}}/\mathcal{D}'$ , to the degree -1 morphism  $f(D)$  in  $\mathcal{E}'$ .

The construction of the dg functor  $\bar{F}$  in the paragraph above depends on making choices and is not unique; the uniqueness is only up to homotopy, meaning in the category **Hqe**. But if  $\mathcal{E}_1$  and  $\mathcal{E}_2$  are pretriangulated dg categories with full pretriangulated dg subcategories  $\mathcal{D}_1 \subset \mathcal{E}_1$  and  $\mathcal{D}_2 \subset \mathcal{E}_2$ , and  $G: \mathcal{E}_1 \rightarrow \mathcal{E}_2$  is a dg functor with  $G(\mathcal{D}_1) \subseteq \mathcal{D}_2$ , then  $G^{\text{hf}}$  induces a natural dg functor  $\mathcal{E}_1^{\text{hf}}/\mathcal{D}'_1 \rightarrow \mathcal{E}_2^{\text{hf}}/\mathcal{D}'_2$ . Both the passage from  $\mathcal{E}$  to  $\mathcal{E}^{\text{hf}}$  and Drinfeld’s construction of the quotient  $\mathcal{E}^{\text{hf}}/\mathcal{D}'$  are manifestly functorial.

In the rest of the paper, we will sometimes sloppily denote by  $\mathcal{E}/\mathcal{D}$  either the Drinfeld quotient of  $\mathcal{E}$  by  $\mathcal{D}$  when  $\mathcal{E}$  is h-flat or of  $\mathcal{E}^{\text{hf}}$  by  $\mathcal{D}'$  otherwise.

### 3.3. The model structure and homotopy pullbacks

Let us recall that the pullback of a diagram

$$\mathcal{E}_1 \xrightarrow{F_1} \mathcal{D} \xleftarrow{F_2} \mathcal{E}_2 \tag{3.5}$$

in **dgCat** is given by a dg category  $\mathcal{E}_1 \times_{\mathcal{D}} \mathcal{E}_2$  defined in the obvious way. Unfortunately, this notion of pullback does not behave well with respect to quasi-equivalences.

To overcome this issue, one has to note that by the work of Tabuada [44], **dgCat** has a model category structure whose weak equivalences are the quasi-equivalences. We refer to [21] for an exhaustive introduction to model categories. Here we content ourselves with some remarks about the special case of **dgCat**. In particular, in Tabuada’s model structure, all dg categories are fibrant, but not all are cofibrant. Furthermore, such a model structure is *right proper*: that is, every pullback of a weak equivalence along a fibration is a weak equivalence, thanks to the fact that all objects are fibrant (see [20, Corollary 13.1.3]). Finally, **Hqe** can be reinterpreted as the homotopy category of **dgCat** with respect to such a model structure.

As is explained, for instance, in [20, Section 13.3], one can then consider the *homotopy pullback*  $\mathcal{E}_1 \times_{\mathcal{D}}^h \mathcal{E}_2$  of equation (3.5). By definition  $\mathcal{E}_1 \times_{\mathcal{D}}^h \mathcal{E}_2 := \mathcal{E}'_1 \times_{\mathcal{D}} \mathcal{E}'_2$  is the usual pullback of a diagram

$$\mathcal{E}'_1 \xrightarrow{F'_1} \mathcal{D} \xleftarrow{F'_2} \mathcal{E}'_2, \tag{3.6}$$

where at least one among  $F'_1$  and  $F'_2$  is a fibration and (for  $i = 1, 2$ )  $F_i = F'_i \circ l_i$  with  $l_i: \mathcal{E}_i \rightarrow \mathcal{E}'_i$  a quasi-equivalence. Notice that such a factorisation of  $F_i$  always exists, and in fact one could choose  $l_i$  to be a cofibration as well. The homotopy pullback does not depend, up to isomorphism in **Hqe**, on the choice of the diagram given by equation (3.6).

Let us spell out an explicit description of  $\mathcal{E}_1 \times_{\mathcal{D}}^h \mathcal{E}_2$ . We can take  $F'_2 = F_2$  and factor only  $F_1$  as follows. Define  $\mathcal{E}'_1$  to be the dg category whose objects are triples,  $(C_1, D, f)$ , where  $C_1 \in \text{Ob}(\mathcal{E}_1)$ ,  $D \in \text{Ob}(\mathcal{D})$  and  $f: F_1(C_1) \rightarrow D$  is a homotopy equivalence. A morphism of degree  $n$  from  $(C_1, D, f)$  to  $(C'_1, D', f')$  in  $\mathcal{E}'_1$  is given by a triple  $(a_1, b, h)$  with  $a_1 \in \text{Hom}_{\mathcal{E}_1}(C_1, C'_1)^n$ ,  $b \in \text{Hom}_{\mathcal{D}}(D, D')^n$  and  $h \in \text{Hom}_{\mathcal{D}}(F_1(C_1), D')^{n-1}$ . The differential is defined by

$$d(a_1, b, h) := (d(a_1), d(b), d(h) + (-1)^n(f' \circ F_1(a_1) - b \circ f))$$

and the composition by

$$(a'_1, b', h') \circ (a_1, b, h) := (a'_1 \circ a_1, b' \circ b, b' \circ h + (-1)^n h' \circ F_1(a_1)).$$

The dg functor  $l_1$  is defined by  $l_1(C_1) := (C_1, F_1(C_1), \text{id}_{F_1(C_1)})$  on objects and  $l_1(a_1) := (a_1, F_1(a_1), 0)$  on morphisms. On the other hand, the dg functor  $F'_1$  is defined as projection on the second component both on objects and on morphisms. It is not difficult to check that  $l_1$  is a quasi-equivalence and  $F'_1$  is a fibration.

With the above choice,  $\mathcal{E}_1 \times_{\mathcal{D}}^h \mathcal{E}_2$  can be identified with the dg category whose objects are triples  $(C_1, C_2, f)$ , where  $C_i \in \text{Ob}(\mathcal{E}_i)$  for  $i = 1, 2$  and  $f: F_1(C_1) \rightarrow F_2(C_2)$  is a homotopy equivalence. A morphism of degree  $n$  from  $(C_1, C_2, f)$  to  $(C'_1, C'_2, f')$  in  $\mathcal{E}_1 \times_{\mathcal{D}}^h \mathcal{E}_2$  is given by a triple  $(a_1, a_2, h)$  with  $a_i \in \text{Hom}_{\mathcal{E}_i}(C_i, C'_i)^n$  for  $i = 1, 2$  and  $h \in \text{Hom}_{\mathcal{D}}(F_1(C_1), F_2(C'_2))^{n-1}$ . The differential is defined by

$$d(a_1, a_2, h) := (d(a_1), d(a_2), d(h) + (-1)^n (f' \circ F_1(a_1) - F_2(a_2) \circ f))$$

and the composition by

$$(a'_1, a'_2, h') \circ (a_1, a_2, h) := (a'_1 \circ a_1, a'_2 \circ a_2, F_2(a'_2) \circ h + (-1)^n h' \circ F_1(a_1)).$$

**Remark 3.12.** By the universal property of the pullback, if  $\mathcal{C}$  is a dg category with dg functors  $G_i: \mathcal{C} \rightarrow \mathcal{E}_i$  for  $i = 1, 2$  making the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{G_2} & \mathcal{E}_2 \\ G_1 \downarrow & & \downarrow F_2 \\ \mathcal{E}_1 & \xrightarrow{F_1} & \mathcal{D} \end{array} \tag{3.7}$$

commutative in **dgCat**, then there is a unique dg functor  $F: \mathcal{C} \rightarrow \mathcal{E}_1 \times_{\mathcal{D}}^h \mathcal{E}_2$  making the diagram

$$\begin{array}{ccccc} \mathcal{C} & \xrightarrow{G_2} & \mathcal{E}_2 & & \\ \downarrow G_1 & \searrow F & \downarrow l_2 & & \\ & \mathcal{E}_1 \times_{\mathcal{D}}^h \mathcal{E}_2 & \longrightarrow & \mathcal{E}'_2 & \\ & \downarrow & & \downarrow F'_2 & \\ \mathcal{E}_1 & \xrightarrow{l_1} & \mathcal{E}'_1 & \xrightarrow{F'_1} & \mathcal{D} \end{array}$$

commutative in **dgCat**. Given the explicit description of the homotopy pullback discussed above, the dg functor  $F$  is defined by

$$F(C) := (G_1(C), G_2(C), \text{id}_{F_1(G_1(C))})$$

on objects and

$$F(a) := (G_1(a), G_2(a), 0)$$

on morphisms.

**Remark 3.13.** The homotopy pullback is a special instance of the concept of homotopy limit in a model category for which exhaustive presentations are available—for example, in [20, Section 19.1]. What is important for us is that homotopy limits are well-defined once we invert weak equivalences. Concretely,

given a small category  $\mathcal{N}$  and a functor  $F: \mathcal{N} \rightarrow \mathbf{dgCat}$ , the homotopy limit  $\mathbf{Holim}(F)$  is a well-defined object of  $\mathbf{Hqe}$  up to canonical isomorphism. Moreover, suppose we are given a diagram

$$\begin{array}{ccc}
 \mathcal{N} & \begin{array}{c} \xrightarrow{F} \\ \Downarrow \theta \\ \xrightarrow{G} \end{array} & \mathbf{dgCat} \xrightarrow{\pi} \mathbf{Hqe} \\
 & & \\
 & & 
 \end{array}$$

meaning  $\mathcal{N}$  is a small category,  $F, G: \mathcal{N} \rightarrow \mathbf{dgCat}$  are functors,  $\theta: F \rightarrow G$  is a natural transformation and  $\pi: \mathbf{dgCat} \rightarrow \mathbf{Hqe}$  is the natural functor. Then  $\mathbf{Holim}(\theta)$  delivers a well-defined morphism in  $\mathbf{Hqe}$  (up to canonical isomorphism), and if  $\pi \circ \theta: \pi: F \rightarrow \pi: G$  is an isomorphism, then so is  $\mathbf{Holim}(\theta)$ . Thus  $\mathbf{Holim}: \mathbf{Hom}(\mathcal{N}, \mathbf{dgCat}) \rightarrow \mathbf{Hqe}$  takes morphisms in  $\mathbf{Hom}(\mathcal{N}, \mathbf{dgCat})$  that induce isomorphisms in  $\mathbf{Hom}(\mathcal{N}, \mathbf{Hqe})$  to isomorphisms in  $\mathbf{Hqe}$ .

What we will need in Section 7 is the following simple case. We start with a finite set of integers  $N := \{1, \dots, n\}$  and form the category  $\mathcal{N}$ , whose objects are the subsets of  $N$  and whose morphisms are the inclusions. We then consider a functor  $\mathcal{N} \rightarrow \mathbf{dgCat}$  and denote by  $\mathcal{E}_I$  the dg category corresponding to  $I \subseteq N$ . We can then form the homotopy limit

$$\mathbf{Holim}_{\emptyset \neq I \subseteq N} \mathcal{E}_I.$$

The special case when  $n = 2$  is just a homotopy pullback, specifically the homotopy pullback of  $\mathcal{E}_{\{1\}} \rightarrow \mathcal{E}_{\{1,2\}} \leftarrow \mathcal{E}_{\{2\}}$ . In the discussion just preceding Remark 3.12, we spelled out an explicit, functorial construction for homotopy pullbacks, allowing us to enhance the homotopy pullback to a functor  $\mathbf{Holim}: \mathbf{Hom}(\{1\} \rightarrow \{1, 2\} \leftarrow \{2\}, \mathbf{dgCat}) \rightarrow \mathbf{dgCat}$ . This will allow us to iterate the construction. We will end up reducing the computation of the homotopy limit, in the case where  $n > 2$ , to iterated homotopy pullbacks.

### 3.4. Localisations in dg categories

Let us first recall a basic construction in the context of triangulated categories. Let  $\mathcal{T}$  be such a category, and suppose further that  $\mathcal{T}$  is closed under arbitrary coproducts. Let  $\mathcal{S} \subset \mathcal{T}^c$  be a collection of compact objects closed under shifts. Then every object  $T \in \mathcal{T}$  sits in a distinguished triangle

$$S \longrightarrow T \longrightarrow T_S$$

with  $S \in \mathbf{Loc}(\mathcal{S})$ , the smallest full triangulated subcategory of  $\mathcal{T}$  containing  $\mathcal{S}$  and closed under arbitrary coproducts, and  $T_S$  in  $\mathcal{S}^\perp := \{T \in \mathcal{T} : \mathbf{Hom}_{\mathcal{T}}(\mathcal{S}, T) = 0\}$ .

The expert reader has certainly noticed that this observation follows from the existence of a semi-orthogonal decomposition for  $\mathcal{T}$  with factors given by  $\mathcal{S}$  and  $\mathcal{S}^\perp \cong \mathcal{T}/\mathcal{S}$ . But here, we want to stress that the construction of  $S$  and  $T_S$  is explicit. Indeed, put  $T_0 = T$ , and then inductively construct distinguished triangles

$$\coprod_{C \rightarrow T_i} C \longrightarrow T_i \longrightarrow T_{i+1},$$

where the first map is the coproduct of all morphisms  $C \rightarrow T_i$  with  $C \in \mathbf{Ob}(\mathcal{S})$ . Then we consider the map  $T \longrightarrow \mathbf{Hocolim} T_i$  and complete it to a distinguished triangle

$$S \longrightarrow T \longrightarrow \mathbf{Hocolim} T_i.$$

A direct computation shows that  $S \in \mathbf{Ob}(\mathbf{Loc}(\mathcal{S}))$  and  $T_S := \mathbf{Hocolim} T_i \in \mathbf{Ob}(\mathcal{S}^\perp)$  (see, for example, [34, Lemma 1.7]).

The following result will be used later and provides an enhancement of the above discussion to the setting of dg categories. Actually, the content of the proposition is more elaborate and deals with two distinct (and orthogonal) localisations.

**Proposition 3.14.** *Let  $\mathcal{C}$  be a dg category, and suppose that  $\mathcal{D}_1$  and  $\mathcal{D}_2$  are full dg subcategories of  $\mathcal{C}$  such that*

- (i)  $\mathcal{D}_1$  and  $\mathcal{D}_2$  are closed under shifts.
- (ii)  $\text{Hom}_{H^0(\mathcal{C})}(\mathcal{D}_i, \mathcal{D}_j) = 0$  for  $i \neq j \in \{1, 2\}$ .

Then for any  $C \in \text{Ob}(\text{h-proj}(\mathcal{C}))$ , there exists a diagram in  $Z^0(\text{h-proj}(\mathcal{C}))$

$$\begin{array}{ccccc}
 & & D_2 & \xlongequal{\quad} & D_2 & & \\
 & & \downarrow & & \downarrow & & \\
 D_1 & \longrightarrow & C & \longrightarrow & C_{D_1} & & \\
 \parallel & & \downarrow & & \downarrow & & \\
 D_1 & \longrightarrow & C_{D_2} & \longrightarrow & C_{D_1, D_2} & & 
 \end{array} \tag{3.8}$$

that is commutative in  $H^0(\text{h-proj}(\mathcal{C}))$  and such that

- (1) its rows and columns yield distinguished triangles in  $H^0(\text{h-proj}(\mathcal{C}))$ ;
- (2)  $D_1 \in \text{Ob}(\text{h-proj}(\mathcal{D}_1))$  and  $D_2 \in \text{Ob}(\text{h-proj}(\mathcal{D}_2))$ ;
- (3) the complexes

$$\text{Hom}_{\text{h-proj}(\mathcal{C})}(\text{h-proj}(\mathcal{D}_i), C_{D_i}) \quad \text{Hom}_{\text{h-proj}(\mathcal{C})}(\text{h-proj}(\mathcal{D}_i), C_{D_1, D_2})$$

are acyclic for  $i = 1, 2$ .

*Proof.* To obtain the objects  $D_i$  and  $C_{D_i}$  and the triangles in the second row and the second column in equation (3.8), we proceed by enhancing the construction at the beginning of this section to the context of dg categories. Since the situation is symmetric, we denote by  $\mathcal{D}$  either of the two dg subcategories  $\mathcal{D}_1$  and  $\mathcal{D}_2$ . Given  $C \in \text{Ob}(\text{h-proj}(\mathcal{C}))$ , we set  $C_0 = C$  and then inductively define  $C_{i+1}$  to be the cone of the map

$$\coprod_{P \rightarrow C_i} P \longrightarrow C_i,$$

where the coproduct is over all objects  $P \in \text{Ob}(\mathcal{D})$  and a set of morphisms in  $\text{Hom}_{Z^0(\text{h-proj}(\mathcal{D}))}(P, C_i)$  representing all morphisms in  $\text{Hom}_{H^0(\text{h-proj}(\mathcal{D}))}(P, C_i)$ . As above, the natural map  $C \rightarrow \text{Hocolim } C_i$  in  $Z^0(\text{h-proj}(\mathcal{C}))$  sits in the triangle

$$D \longrightarrow C \longrightarrow \text{Hocolim } C_i,$$

which is distinguished in  $H^0(\text{h-proj}(\mathcal{C}))$ . The argument above now shows that  $D$  is an object of  $\text{h-proj}(\mathcal{D})$  and  $C_D := \text{Hocolim } C_i$  is such that  $\text{Hom}_{\text{h-proj}(\mathcal{C})}(Q, C_D)$  is acyclic for all  $Q \in \text{Ob}(\text{h-proj}(\mathcal{D}))$ .

Since  $\text{h-proj}(\mathcal{C})$  is pretriangulated, the second row and second column we just constructed can be completed to the diagram of equation (3.8), where the rows and columns are cofibre sequences, in particular yielding distinguished triangles in  $H^0(\text{h-proj}(\mathcal{C}))$ .

It remains to prove that the object  $C_{D_1, D_2}$  that we constructed is such that the complex  $\text{Hom}_{\text{h-proj}(\mathcal{C})}(\text{h-proj}(\mathcal{D}_i), C_{D_1, D_2})$  is acyclic for  $i = 1, 2$ . To this end, we apply the functor

$\text{Hom}_{\text{h-proj}(\mathcal{C})}(\mathcal{D}_i, -)$  to the triangle

$$D_j \longrightarrow C_{D_i} \longrightarrow C_{D_1, D_2},$$

where  $i \neq j \in \{1, 2\}$ . The complex  $\text{Hom}_{\text{h-proj}(\mathcal{C})}(\mathcal{D}_i, C_{D_i})$  is clearly acyclic from the first part of the construction. By (ii), we know that  $\text{Hom}_{\mathcal{C}}(\mathcal{D}_i, \mathcal{D}_j)$  is acyclic, and thus  $\text{Hom}_{\text{h-proj}(\mathcal{C})}(\mathcal{D}_i, \text{h-proj}(\mathcal{D}_j))$  is acyclic as well, because the objects in  $\mathcal{D}_i$  are in  $\mathcal{C}$  and thus are compact in  $H^0(\text{h-proj}(\mathcal{C}))$ . Since  $D_j$  belongs to  $\text{h-proj}(\mathcal{D}_j)$ , we obtain that  $\text{Hom}_{\text{h-proj}(\mathcal{C})}(\mathcal{D}_i, D_j)$  is acyclic. From this, we deduce that  $\text{Hom}_{\text{h-proj}(\mathcal{C})}(\mathcal{D}_i, C_{D_1, D_2})$  is acyclic, and this implies that  $\text{Hom}_{\text{h-proj}(\mathcal{C})}(\text{h-proj}(\mathcal{D}_i), C_{D_1, D_2})$  is acyclic as well. □

### 3.5. Dg enhancements and their uniqueness

In this section, we will be careful about the universe where the triangulated and dg categories live—mostly to show that such care is superfluous.

**Definition 3.15.** Let  $\mathbb{U} \in \mathbb{V}$  be universes, and consider a  $\mathbb{U}$ -small triangulated category  $\mathcal{T}$ . A dg  $\mathbb{V}$ -enhancement (or simply a  $\mathbb{V}$ -enhancement) of  $\mathcal{T}$  is a pair  $(\mathcal{C}, E)$ , where  $\mathcal{C}$  is a  $\mathbb{V}$ -small pretriangulated dg category and  $E: H^0(\mathcal{C}) \rightarrow \mathcal{T}$  is an exact equivalence.

By abuse of notation, one says that  $\mathcal{C}$  is a  $\mathbb{V}$ -enhancement of  $\mathcal{T}$  if  $E$  is clear from the context.

**Example 3.16.** If  $\mathcal{A}$  is a  $\mathbb{U}$ -small additive category, then  $\mathbf{C}_{\text{dg}}^2(\mathcal{A})$  is in a natural way a  $\mathbb{U}$ -enhancement of  $\mathbf{K}^2(\mathcal{A})$  (see Example 3.1). If  $\mathcal{A}$  is abelian, then, since  $\mathbf{D}^2(\mathcal{A}) = \mathbf{K}^2(\mathcal{A})/\mathbf{K}_{\text{acy}}^2(\mathcal{A})$ , the discussion in Section 3.2 allows us to conclude that the Drinfeld quotient

$$\mathbf{D}_{\text{dg}}^2(\mathcal{A}) := \mathbf{C}_{\text{dg}}^2(\mathcal{A})/\mathbf{Acy}^2(\mathcal{A})$$

is a  $\mathbb{U}$ -enhancement of  $\mathbf{D}^2(\mathcal{A})$ . Here  $\mathbf{Acy}^2(\mathcal{A})$  denotes the full dg subcategory of  $\mathbf{C}_{\text{dg}}^2(\mathcal{A})$  whose objects correspond to those of  $\mathbf{K}_{\text{acy}}^2(\mathcal{A})$ .

The core of this paper is to study when dg enhancements are unique in the following sense.

**Definition 3.17.** The  $\mathbb{U}$ -small triangulated category  $\mathcal{T}$  has a unique  $\mathbb{V}$ -enhancement if, given two  $\mathbb{V}$ -enhancements  $(\mathcal{C}_1, E_1)$  and  $(\mathcal{C}_2, E_2)$  of  $\mathcal{T}$ , there is a  $\mathbb{V}$ -small pretriangulated dg category  $\mathcal{C}_3$  and quasi-equivalences  $l_i: \mathcal{C}_3 \rightarrow \mathcal{C}_i$  for  $i = 1, 2$ .

Let us now prove the following result.

**Proposition 3.18.** Let  $\mathcal{T}$  be a  $\mathbb{U}$ -small triangulated category. If  $\mathcal{T}$  has a  $\mathbb{V}$ -enhancement for some universe  $\mathbb{V}$ , with  $\mathbb{U} \in \mathbb{V}$ , then it also has a  $\mathbb{U}$ -enhancement. Furthermore, there exists a universe  $\mathbb{V}$ , with  $\mathbb{U} \in \mathbb{V}$  and such that  $\mathcal{T}$  has a unique  $\mathbb{V}$ -enhancement, if and only if  $\mathcal{T}$  has a unique  $\mathbb{U}$ -enhancement. Moreover, if  $\mathcal{T}$  has a unique  $\mathbb{U}$ -enhancement, then it has a unique  $\mathbb{V}$  enhancement for all  $\mathbb{U} \in \mathbb{V}$ .

*Proof.* Suppose  $\mathcal{T}$  has a  $\mathbb{V}$ -enhancement  $\mathcal{C}$ . In Lemma 3.8 (see also Remark 3.9), we produced a quasi-equivalence  $l_{\mathcal{C}}^{\text{sm}}: \mathcal{C}^{\text{sm}} \rightarrow \mathcal{C}$ , with  $\mathcal{C}^{\text{sm}}$  a  $\mathbb{U}$ -small pretriangulated dg category; this gives the existence of a  $\mathbb{U}$ -enhancement for  $\mathcal{T}$ .

Now let  $\mathbb{V}$  be a universe with  $\mathbb{U} \in \mathbb{V}$  and such that  $\mathcal{T}$  has a unique  $\mathbb{V}$ -enhancement. Let  $\mathcal{C}_1$  and  $\mathcal{C}_2$  be two  $\mathbb{U}$ -enhancements of  $\mathcal{T}$ . By the uniqueness of  $\mathbb{V}$ -enhancements, there exists a  $\mathbb{V}$ -enhancement  $\mathcal{C}_3$  and quasi-equivalences  $l_i: \mathcal{C}_3 \rightarrow \mathcal{C}_i$  for  $i = 1, 2$ . Lemma 3.8 produces for us a quasi-equivalence  $l_{\mathcal{C}_3}^{\text{sm}}: \mathcal{C}_3^{\text{sm}} \rightarrow \mathcal{C}_3$  with  $\mathcal{C}_3^{\text{sm}}$  a  $\mathbb{U}$ -small pretriangulated dg category, and the composites  $l_i \circ l_{\mathcal{C}_3}^{\text{sm}}: \mathcal{C}_3^{\text{sm}} \rightarrow \mathcal{C}_i$  for  $i = 1, 2$  show that the  $\mathbb{U}$ -enhancement is unique.

Finally, the ‘moreover’ part. Suppose, therefore, that  $\mathcal{T}$  has a unique  $\mathbb{U}$ -enhancement and  $\mathbb{V}$  is some universe with  $\mathbb{U} \in \mathbb{V}$ . Let  $\mathcal{C}_1$  and  $\mathcal{C}_2$  be two pretriangulated dg  $\mathbb{V}$ -categories enhancing  $\mathcal{T}$ . Lemma 3.8 produces for us quasi-equivalences  $l_i^{\text{sm}}: \mathcal{C}_i^{\text{sm}} \rightarrow \mathcal{C}_i$  for  $i = 1, 2$ , with  $\mathcal{C}_i^{\text{sm}}$  both  $\mathbb{U}$ -small. By



the uniqueness of  $\mathbb{U}$ -enhancements, there exists a  $\mathbb{U}$ -small pretriangulated dg category  $\mathcal{C}_3$  and quasi-equivalences  $l_i: \mathcal{C}_3 \rightarrow \mathcal{C}_i^{\text{sm}}$  for  $i = 1, 2$ . But now the composites  $l_i^{\text{sm}} \circ l_i: \mathcal{C}_3 \rightarrow \mathcal{C}_i$  for  $i = 1, 2$  show that the enhancements to  $\mathbb{V}$  are unique.  $\square$

Given this, in the rest of the paper, it will rarely be necessary to specify in which universe we are working. For this reason, we will freely talk about dg enhancements and their uniqueness without mentioning any universe.

**Remark 3.19.** (i) The property of having a unique dg enhancement is clearly invariant under exact equivalences. More precisely, if  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are triangulated categories such that there is an exact equivalence  $\mathcal{T}_1 \cong \mathcal{T}_2$ , then  $\mathcal{T}_1$  has a unique dg enhancement if and only if  $\mathcal{T}_2$  does.

(ii) Given a triangulated category  $\mathcal{T}$ , the category  $\mathcal{T}^\circ$  has a natural triangulated structure. Since there is a natural exact equivalence  $H^0(\mathcal{D})^\circ \cong H^0(\mathcal{D}^\circ)$  for any pretriangulated dg category  $\mathcal{D}$ , it is clear that  $\mathcal{T}$  has a unique dg enhancement if and only if  $\mathcal{T}^\circ$  does.

An equivalent way to state Definition 3.17 is by saying that there is an isomorphism  $f \in \text{Hom}_{\mathbf{Hqe}}(\mathcal{C}_1, \mathcal{C}_2)$ . With this in mind, we can state the following stronger definition.

**Definition 3.20.** A triangulated category  $\mathcal{T}$  has a *semi-strongly unique enhancement* if, given two enhancements  $(\mathcal{C}_1, E_1)$  and  $(\mathcal{C}_2, E_2)$  of  $\mathcal{T}$ , there is an isomorphism  $f \in \text{Hom}_{\mathbf{Hqe}}(\mathcal{C}_1, \mathcal{C}_2)$  such that  $E_1(C) \cong E_2(H^0(f))(C)$  in  $\mathcal{T}$  for every  $C \in \text{Ob}(\mathcal{C}_1)$ .

### 4. A special zigzag of dg functors

This slightly technical section provides a useful enhancement for the category  $\mathbf{B}^?(A)$  of Section 1.1. This is done in Section 4.1. Section 4.4 deals with the important relation between enhancements of  $\mathbf{V}^?(A)$  and  $\mathbf{B}^?(A)$ . This is achieved with a complicated argument that involves a variant of the enhancement of  $\mathbf{B}^?(A)$  constructed in Section 4.1 (see Section 4.2) and a discussion about homotopy limits in Section 4.3, where we make explicit the construction in Section 3.3 in a very concrete context.

#### 4.1. An enhancement for $\mathbf{B}^?(A)$

Remembering Example 3.16, it is clear that for  $? = b, +, -, \emptyset$ , the full dg subcategory  $\mathcal{V} = \mathcal{V}^?(A)$  of  $\mathbf{C}_{\text{dg}}^?(A)$  whose objects are complexes with trivial differential is in a natural way an enhancement of  $\mathbf{V}^?(A)$ .

Assume from now on that  $A$  is abelian, and fix an enhancement  $(\mathcal{C}, E)$  of  $\mathbf{D}^?(A)$ . We are going to define a dg category  $\mathcal{B}_{\mathcal{C}, E}^?(A)$ , which will turn out to be an enhancement of  $\mathbf{B}^?(A)$ . Recall that the latter category is the full subcategory of  $\mathbf{D}^?(A)$  consisting of complexes with trivial differential. The notation for  $\mathcal{B}_{\mathcal{C}, E}^?(A)$  alludes to the fact that its definition depends on the pair  $(\mathcal{C}, E)$ . But in the sequel, when there is no risk of confusion, we will use the shorthand  $\mathcal{B} := \mathcal{B}_{\mathcal{C}, E}^?(A)$ .

An object  $B = (B^-, B^+, B^i, \alpha^i, \beta^i)_{i \in \mathbb{Z}}$  of  $\mathcal{B}$  is given by objects  $B^-, B^+$  and  $B^i$  of  $\mathcal{C}$  together with morphisms  $B^i[-i] \xrightarrow{\alpha^i} B^? \xrightarrow{\beta^i} B^i[-i]$  of  $Z^0(\mathcal{C})$  (where  $? = +$  if  $i > 0$  and  $? = -$  if  $i \leq 0$ ) such that the following conditions are satisfied:

- (B.1) If  $i, j$  are both  $> 0$  or both  $\leq 0$ , then  $\beta^j \circ \alpha^i$  is  $\text{id}_{B^i[-i]}$  for  $i = j$  and is 0 for  $i \neq j$ .
- (B.2)  $E(B^i) \in \text{Ob}(A)$  for every  $i \in \mathbb{Z}$ .
- (B.3) The morphisms  $\alpha^i$  with  $i \leq 0$  and  $\beta^i$  with  $i > 0$  induce isomorphisms in  $H^0(\mathcal{C}) \cong \mathbf{D}^?(A)$

$$\prod_{i \leq 0} B^i[-i] \xrightarrow{\sim} B^-, \quad B^+ \xrightarrow{\sim} \prod_{i > 0} B^i[-i].$$

Given objects  $B_k = (B_k^-, B_k^+, B_k^i, \alpha_k^i, \beta_k^i)_{i \in \mathbb{Z}}$ , for  $k = 1, 2$ , we define

$$\text{Hom}_{\mathcal{B}}(B_1, B_2) := \text{Hom}_{\mathcal{C}}(B_1^- \oplus B_1^+, B_2^- \oplus B_2^+).$$

It is straightforward to check that  $\mathcal{B}$  (with the obvious composition of morphisms) is a dg category and that the map defined on objects by  $(B^-, B^+, B^i, \alpha^i, \beta^i)_{i \in \mathbb{Z}} \mapsto B^- \oplus B^+$  extends to a fully faithful dg functor  $\mathbf{B}: \mathcal{B} \rightarrow \mathcal{C}$ .

One can consider the following variant, which will be used in the proof of Theorem B. Let  $\mathcal{A}$  be an abelian category with a Serre subcategory  $\mathcal{E} \subseteq \mathcal{A}$ . Consider then the full triangulated subcategory  $\mathbf{D}_{\mathcal{E}}^?(\mathcal{A})$  of  $\mathbf{D}^?(\mathcal{A})$  consisting of all complexes with cohomology in  $\mathcal{E}$ . There is a natural exact functor

$$\pi: \mathbf{D}^?(\mathcal{E}) \longrightarrow \mathbf{D}_{\mathcal{E}}^?(\mathcal{A}),$$

which in general is not an equivalence.

Given a dg enhancement  $(\mathcal{E}, \mathbf{E})$  of  $\mathbf{D}_{\mathcal{E}}^?(\mathcal{A})$ , we define a dg category  $\widehat{\mathcal{B}}_{\mathcal{E}, \mathbf{E}}^?(\mathcal{A}, \mathcal{E})$ , for which we use the shorthand  $\widehat{\mathcal{B}}$ , in a fashion that is very similar to  $\mathcal{B}$ .

In particular, an object  $B = (B^-, B^+, B^i, \alpha^i, \beta^i)_{i \in \mathbb{Z}}$  of  $\widehat{\mathcal{B}}$  is given by objects  $B^-, B^+$  and  $B^i$  of  $\mathcal{C}$  together with morphisms  $B^i[-i] \xrightarrow{\alpha^i} B^? \xrightarrow{\beta^i} B^i[-i]$  of  $Z^0(\mathcal{C})$  (where  $? = +$  if  $i > 0$  and  $? = -$  if  $i \leq 0$ ) such that (B.1) is satisfied, while (B.2) and (B.3) are replaced, respectively, by the following:

(B.2\*)  $\mathbf{E}(B^i) \cong \pi(Q_i)$  for some  $Q_i \in \text{Ob}(\mathcal{E})$  and for every  $i \in \mathbb{Z}$ .

(B.3\*) The morphisms  $\alpha^i$  with  $i \leq 0$  and  $\beta^i$  with  $i > 0$  induce isomorphisms in  $\mathbf{D}_{\mathcal{E}}^?(\mathcal{A})$

$$\pi\left(\prod_{i \leq 0} Q^i[-i]\right) \xrightarrow{\sim} \mathbf{E}(B^-), \quad \mathbf{E}(B^+) \xrightarrow{\sim} \pi\left(\prod_{i > 0} Q^i[-i]\right).$$

As for the Hom-spaces, given objects  $B_k = (B_k^-, B_k^+, B_k^i, \alpha_k^i, \beta_k^i)_{i \in \mathbb{Z}}$ , for  $k = 1, 2$ , we keep the same definition and set

$$\text{Hom}_{\widehat{\mathcal{B}}}(B_1, B_2) := \text{Hom}_{\mathcal{C}}(B_1^- \oplus B_1^+, B_2^- \oplus B_2^+).$$

Recall the h-flat resolution  $\text{lh}^{\text{f}}: \mathcal{C}^{\text{hf}} \rightarrow \mathcal{C}$  of Proposition 3.10.

**Proposition 4.1.** *If  $(\mathcal{C}, \mathbf{E})$  is a dg enhancement of  $\mathbf{D}_{\mathcal{E}}^?(\mathcal{A})$ , then  $\text{lh}^{\text{f}}$  induces a quasi-equivalence*

$$\widehat{\text{lh}}^{\text{f}}: \widehat{\mathcal{B}}_{\mathcal{C}^{\text{hf}}, \mathbf{E} \circ H^0(\text{lh}^{\text{f}})}^?(\mathcal{A}, \mathcal{E}) \longrightarrow \widehat{\mathcal{B}}_{\mathcal{C}, \mathbf{E}}^?(\mathcal{A}, \mathcal{E}),$$

which is also surjective on objects.

*Proof.* The result follows from a direct check using the definitions, and we only briefly outline it here. Since the dg functor  $\text{lh}^{\text{f}}$  is the identity on objects, we define  $\widehat{\text{lh}}^{\text{f}}$  by sending an object  $(B^-, B^+, B^i, \widetilde{\alpha}^i, \widetilde{\beta}^i)_{i \in \mathbb{Z}}$  of  $\widehat{\mathcal{B}}^{\text{hf}} := \widehat{\mathcal{B}}_{\mathcal{C}^{\text{hf}}, \mathbf{E} \circ H^0(\text{lh}^{\text{f}})}^?(\mathcal{A}, \mathcal{E})$  to  $(B^-, B^+, B^i, \text{lh}^{\text{f}}(\widetilde{\alpha}^i), \text{lh}^{\text{f}}(\widetilde{\beta}^i))_{i \in \mathbb{Z}}$  in  $\widehat{\mathcal{B}} := \widehat{\mathcal{B}}_{\mathcal{C}, \mathbf{E}}^?(\mathcal{A}, \mathcal{E})$ . The fact that morphisms in  $\widehat{\mathcal{B}}^{\text{hf}}$  (respectively,  $\widehat{\mathcal{B}}$ ) are defined as in  $\mathcal{C}^{\text{hf}}$  (respectively,  $\mathcal{C}$ ) clearly implies that  $\widehat{\text{lh}}^{\text{f}}$  extends to a dg functor if we define it on morphisms like  $\text{lh}^{\text{f}}$ . It is also obvious that  $\widehat{\text{lh}}^{\text{f}}$  is quasi-fully faithful because  $\text{lh}^{\text{f}}$  is, and it remains to show that  $\widehat{\text{lh}}^{\text{f}}$  is surjective on objects. Given  $(B^-, B^+, B^i, \alpha^i, \beta^i)_{i \in \mathbb{Z}} \in \text{Ob}(\widehat{\mathcal{B}})$ , the closed degree-0 morphisms  $\alpha^i$  and  $\beta^i$  can be seen as morphisms in the dg category  $\mathcal{E}_1$ —with the notation used in Section 3.2 to define  $\mathcal{C}^{\text{hf}}$ , we may view  $\alpha^i$  and  $\beta^i$  as belonging to the bases  $D_{\mathcal{E}}^1(B^i[-i], B^?)$  (respectively  $D_{\mathcal{E}}^1(B^?, B^i[-i])$ ) freely generating  $\text{Hom}_{\mathcal{E}_1}(B^i[-i], B^?)$  (respectively  $\text{Hom}_{\mathcal{E}_1}(B^?, B^i[-i])$ ) as modules over  $\mathbb{k}$ —and the relations  $\beta^i \circ \alpha^i = \text{id}$  and  $\beta^j \circ \alpha^i = 0$  for  $i \neq j$ , hold in  $\mathcal{E}_1$  by definition, and therefore also in  $\mathcal{C}^{\text{hf}} = \text{colim } \mathcal{E}_i$ . Thus we obtain an object in  $\widehat{\mathcal{B}}^{\text{hf}}$  that is mapped to  $(B^-, B^+, B^i, \alpha^i, \beta^i)_{i \in \mathbb{Z}}$  by  $\widehat{\text{lh}}^{\text{f}}$ .  $\square$

Back to the dg category  $\mathcal{B}$ , in the following, we will assume that:

$$Z^0(\mathcal{C}) \text{ idempotent complete and closed under coproducts of null-homotopic objects.} \tag{4.1}$$

Notice that by Remark 3.5, this condition can be always achieved, up to replacing  $\mathcal{C}$  with  $\text{Pretr}(\mathcal{C}) = \text{Perf}(\mathcal{C})$ .

**Proposition 4.2.** *If  $(\mathcal{C}, E)$  is a dg enhancement of  $\mathbf{D}^2(\mathcal{A})$  such that  $\mathcal{C}$  satisfies equation (4.1), then the essential image of  $E \circ H^0(B): H^0(\mathcal{B}) \rightarrow \mathbf{D}^2(\mathcal{A})$  is  $\mathbf{B}^2(\mathcal{A})$ .*

*Proof.* Given  $B = (B^-, B^+, B^i, \alpha^i, \beta^i)_{i \in \mathbb{Z}}$  in  $\mathcal{B}$ , it is clear from Corollary 1.6 and the definition of  $\mathcal{B}$  that  $E(B(B)) \cong \bigoplus_{i \in \mathbb{Z}} A^i[-i] \in \text{Ob}(\mathbf{B}^2(\mathcal{A}))$ , where  $A^i := E(B^i) \in \text{Ob}(\mathcal{A})$ .

Conversely, given  $A^* = \bigoplus_{i \in \mathbb{Z}} A^i[-i] \in \text{Ob}(\mathbf{B}^2(\mathcal{A}))$ , we can take  $B^i \in \text{Ob}(\mathcal{C})$  such that  $E(B^i) \cong A^i$  for every  $i \in \mathbb{Z}$ . Moreover, let  $C^-, C^+ \in \text{Ob}(\mathcal{C})$  be such that  $E(C^-) \cong \bigoplus_{i \leq 0} A^i[-i]$  and  $E(C^+) \cong \bigoplus_{i > 0} A^i[-i]$ . Then there exist morphisms

$$B^i[-i] \xrightarrow{\tilde{\alpha}^i} C^? \xrightarrow{\tilde{\beta}^i} B^i[-i]$$

of  $Z^0(\mathcal{C})$  (where  $? = +$  if  $i > 0$  and  $? = -$  if  $i \leq 0$ ) such that if  $i, j$  are both  $> 0$  or both  $\leq 0$ , then the image in  $H^0(\mathcal{C})$  of  $\tilde{\beta}^j \circ \tilde{\alpha}^i$  is  $\text{id}_{B^i[-i]}$  for  $i = j$  and is 0 for  $i \neq j$ . Now we set

$$B^- := C^- \oplus \prod_{i \leq 0} \text{Cone}(\text{id}_{B^i[-i]}), \quad B^+ := C^+ \oplus \prod_{i > 0} \text{Cone}(\text{id}_{B^i[-i]}).$$

For every  $j \in \mathbb{Z}$ , we also define  $\alpha^j: B^j[-j] \rightarrow B^?$  as the morphism induced by  $\tilde{\alpha}^j: B^j[-j] \rightarrow C^?$  and by the natural morphism  $B^j[-j] \rightarrow \text{Cone}(\text{id}_{B^j[-j]})$ . On the other hand, we define  $\beta^j: B^? \rightarrow B^j[-j]$  as the morphism induced by  $\tilde{\beta}^j: C^? \rightarrow B^j[-j]$  and by morphisms  $\text{Cone}(\text{id}_{B^i[-i]}) \rightarrow B^j[-j]$  in  $Z^0(\mathcal{C})$  (whose existence is ensured by Lemma 3.2) with the property that the composition with the natural morphism  $B^i[-i] \rightarrow \text{Cone}(\text{id}_{B^i[-i]})$  is  $\text{id}_{B^i[-i]} - \tilde{\beta}^i \circ \tilde{\alpha}^i$  for  $i = j$  and is  $-\tilde{\beta}^j \circ \tilde{\alpha}^i$  for  $i \neq j$ . It is then clear that  $B := (B^-, B^+, B^i, \alpha^i, \beta^i)_{i \in \mathbb{Z}}$  is an object of  $\mathcal{B}$  that satisfies  $E(B(B)) \cong A^*$ .  $\square$

**Remark 4.3.** By Proposition 4.2, the dg category  $\mathcal{B}$  is an enhancement of  $\mathbf{B}^2(\mathcal{A})$ . Moreover, in view of Corollary 2.10, and taking into account that  $H^0(\mathcal{C}) \cong \mathbf{D}^2(\mathcal{A})$  is idempotent complete (see Remark 3.5), we have  $\text{Perf}(\mathcal{B}) \cong \mathcal{C}$  in  $\mathbf{Hqe}$  (using [27, Proposition 1.16]). Observe that, similarly,  $\text{Perf}(\mathcal{V}) \cong \mathbf{C}_{\text{dg}}^2(\mathcal{A})$  in  $\mathbf{Hqe}$ . Hence  $\text{Perf}(\mathcal{B})$  (respectively,  $\text{Perf}(\mathcal{V})$ ) is an enhancement of  $\mathbf{D}^2(\mathcal{A})$  (respectively,  $\mathbf{K}^2(\mathcal{A})$ ).

**4.2. A technical interlude**

We first describe a variant of  $\mathcal{B}$ . We start by introducing new Hom spaces between objects of  $\mathcal{B}$ , depending on an integer  $n > 0$ .

Given  $B = (B^-, B^+, B^i, \alpha^i, \beta^i)_{i \in \mathbb{Z}}$  in  $\mathcal{B}$ , the two compositions of morphisms (induced by the  $\alpha^i$  and the  $\beta^i$ )

$$\bigoplus_{i=-n}^0 B^i[-i] \longrightarrow B^- \longrightarrow \bigoplus_{i=-n}^0 B^i[-i] \qquad \bigoplus_{i=1}^n B^i[-i] \longrightarrow B^+ \longrightarrow \bigoplus_{i=1}^n B^i[-i]$$

are obviously the identities. As  $Z^0(\mathcal{C})$  is idempotent complete, this implies that there exist objects  ${}^n B^{-n-1}$  and  ${}^n B^{n+1}$  of  $\mathcal{C}$  such that (setting also  ${}^n B^i := B^i$  for  $-n \leq i \leq n$ )

$$B^- \cong \bigoplus_{i=-n-1}^0 {}^n B^i[-i] \qquad B^+ \cong \bigoplus_{i=1}^{n+1} {}^n B^i[-i]$$

in  $Z^0(\mathcal{C})$ . Notice that if  $n < m$ , then

$${}^n B^{-n-1}[n+1] \cong \bigoplus_{i=-m-1}^{-n-1} {}^m B^i[-i] \qquad {}^n B^{n+1}[-n-1] \cong \bigoplus_{i=n+1}^{m+1} {}^m B^i[-i]$$

in  $Z^0(\mathcal{C})$ . Now, given two objects  $B_1$  and  $B_2$  of  $\mathcal{B}$ , there is an isomorphism of complexes

$$\text{Hom}_{\mathcal{B}}(B_1, B_2) \cong \prod_{i=-n-1}^{n+1} \prod_{j=-n-1}^{n+1} \text{Hom}_{\mathcal{C}}({}^n B_1^i[-i], {}^n B_2^j[-j]),$$

and we define

$$\text{Hom}_n(B_1, B_2) := \prod_{i=-n-1}^{n+1} \prod_{j=-n-1}^{n+1} \text{Hom}_n^{i,j}(B_1, B_2),$$

where (for  $-n-1 \leq i, j \leq n+1$ ),

$$\text{Hom}_n^{i,j}(B_1, B_2) := \begin{cases} \text{Hom}_{\mathcal{C}}({}^n B_1^i[-i], {}^n B_2^j[-j]) & \text{if } i = -n-1 \text{ or } j = n+1 \\ \text{Hom}_{\mathcal{C}}({}^n B_1^i[-i], {}^n B_2^j[-j])^{\leq j-i} & \text{if } i \geq -n \text{ and } j \leq n. \end{cases}$$

Obviously we can identify  $\text{Hom}_n(B_1, B_2)$  with a subcomplex of  $\text{Hom}_{\mathcal{B}}(B_1, B_2)$ .

**Remark 4.4.** One might hope to obtain a dg category  $\mathcal{B}_n$  with the same objects as  $\mathcal{B}$  and such that  $\text{Hom}_{\mathcal{B}_n}(B_1, B_2) := \text{Hom}_n(B_1, B_2)$  for every  $B_1, B_2 \in \text{Ob}(\mathcal{B})$ . Unfortunately, given also  $B_3 \in \text{Ob}(\mathcal{B})$ , the composition map (which is a morphism of complexes)

$$\text{Hom}_{\mathcal{B}}(B_1, B_2) \otimes_{\mathbb{k}} \text{Hom}_{\mathcal{B}}(B_2, B_3) \longrightarrow \text{Hom}_{\mathcal{B}}(B_1, B_3) \tag{4.2}$$

does not send  $\text{Hom}_n(B_1, B_2) \otimes_{\mathbb{k}} \text{Hom}_n(B_2, B_3)$  to  $\text{Hom}_n(B_1, B_3)$ , in general. More precisely,  $\text{Hom}_n^{h,i}(B_1, B_2) \otimes_{\mathbb{k}} \text{Hom}_n^{j,j'}(B_2, B_3)$  need not be sent to  $\text{Hom}_n^{h,j'}(B_1, B_3)$  if  $-n \leq h \leq n+1$ ,  $-n-1 \leq j \leq n$  and  $i = -n-1$  or  $i = n+1$ .

**Remark 4.5.** If  $n < m$ , we can regard  $\text{Hom}_m(B_1, B_2)$  as a subcomplex of  $\text{Hom}_n(B_1, B_2)$ . Indeed, it is clear that for  $m-1 \leq i, j \leq m+1$ ,  $\text{Hom}_{\mathcal{C}}({}^m B_1^i[-i], {}^m B_2^j[-j])$  can be viewed as a subcomplex of  $\text{Hom}_{\mathcal{C}}({}^n B_1^{i'}[-i'], {}^n B_2^{j'}[-j'])$ , where we define

$$i' := \begin{cases} -n-1 & \text{if } -m-1 \leq i \leq -n-1 \\ i & \text{if } -n-1 \leq i \leq n+1 \\ n+1 & \text{if } n+1 \leq i \leq m+1 \end{cases}$$

and similarly for  $j'$ . Since  $j-i \leq j'-i'$  when  $i' \geq -n$  (which implies  $i' \leq i$ ) and  $j' \leq n$  (which implies  $j' \geq j$ ), it follows that in any case,  $\text{Hom}_m^{i,j}(B_1, B_2)$  can be viewed as a subcomplex of  $\text{Hom}_n^{i',j'}(B_1, B_2)$ .

By the above, there is a natural morphism  $\text{Holim Hom}_n(B_1, B_2) \rightarrow \text{Hom}_{\mathcal{B}}(B_1, B_2)$  in  $\mathbf{D}(\text{Mod}(\mathbb{k}))$ . Composing it with the morphism of complexes

$$\text{Hom}_{\mathcal{B}}(B_1, B_2) = \text{Hom}_{\mathcal{C}}(B_1^- \oplus B_1^+, B_2^- \oplus B_2^+) \longrightarrow \prod_{i,j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j])$$

(induced by precomposition with the  $\beta_2^j$  and postcomposition with the  $\alpha_1^i$ ), we obtain a morphism in  $\mathbf{D}(\text{Mod}(\mathbb{k}))$

$$\text{Holim Hom}_n(B_1, B_2) \longrightarrow \prod_{i,j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j]). \tag{4.3}$$

**Lemma 4.6.** For every  $B_1, B_2 \in \text{Ob}(\mathcal{B})$ , equation (4.3) factors (uniquely) through a morphism

$$\text{Holim Hom}_n(B_1, B_2) \longrightarrow \prod_{i,j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j])^{\leq j-i},$$

which is an isomorphism in  $\mathbf{D}(\text{Mod}(\mathbb{k}))$ . Moreover, this isomorphism is compatible with truncations, meaning for every  $k \in \mathbb{Z}$ , the induced morphism

$$\text{Holim Hom}_n(B_1, B_2)^{\leq k} \longrightarrow \prod_{i,j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j])^{\leq \min(k, j-i)}$$

is also an isomorphism in  $\mathbf{D}(\text{Mod}(\mathbb{k}))$ .

*Proof.* Let us fix an integer  $l$ . Notice first that since  $H^l(\text{Hom}_{\mathcal{C}}(C_1, C_2)) \cong \text{Hom}_{H^0(\mathcal{C})}(C_1, C_2[l])$  for every  $C_1, C_2 \in \text{Ob}(\mathcal{C})$ , we have

$$H^l(\text{Hom}_{\mathcal{C}}(C_1, C_2)^{\leq k}) \cong \begin{cases} \text{Hom}_{H^0(\mathcal{C})}(C_1, C_2[l]) & \text{if } l \leq k \\ 0 & \text{if } l > k \end{cases}$$

for every  $k \in \mathbb{Z}$ . In particular, we obtain

$$H^l(\text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j])^{\leq j-i}) \cong \begin{cases} \text{Hom}_{H^0(\mathcal{C})}(B_1^i, B_2^j[l+i-j]) & \text{if } l \leq j-i \\ 0 & \text{if } l > j-i \end{cases}$$

for every  $i, j \in \mathbb{Z}$ . Taking into account that  $\text{Hom}_{H^0(\mathcal{C})}(B_1^i, B_2^j[k]) = 0$  for  $k < 0$  (because  $E(B_1^i), E(B_2^j) \in \text{Ob}(\mathcal{A})$ ), this implies that

$$H^l(\text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j])^{\leq j-i}) \cong \begin{cases} \text{Hom}_{H^0(\mathcal{C})}(B_1^i, B_2^j) & \text{if } l = j-i \\ 0 & \text{if } l \neq j-i. \end{cases}$$

As  $H^l : \mathbf{D}(\text{Mod}(\mathbb{k})) \rightarrow \text{Mod}(\mathbb{k})$  commutes with products, it follows that

$$H^l \left( \prod_{i,j \in \mathbb{Z}} \text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j])^{\leq j-i} \right) \cong \prod_{i \in \mathbb{Z}} \text{Hom}_{H^0(\mathcal{C})}(B_1^i, B_2^{i+l}). \tag{4.4}$$

On the other hand, given integers  $n, i, j$  with  $n > 0$  and  $-n - 1 \leq i, j \leq n + 1$ , we have

$$H^l(\text{Hom}_n^{i,j}(B_1, B_2)) \cong \begin{cases} \text{Hom}_{H^0(\mathcal{C})}({}^n B_1^i[-i], {}^n B_2^j[l-j]) & \text{if } l \leq j-i \text{ or } i = -n-1 \text{ or } j = n+1 \\ 0 & \text{if } l > j-i, i \geq -n \text{ and } j \leq n. \end{cases}$$

We shall henceforth assume  $n > -l/2 - 1$ . Then

$$H^l(\text{Hom}_n^{n+1, -n-1}(B_1, B_2)) = 0$$

(since  $i = n + 1 \geq -n$ ,  $j = -n - 1 \leq n$  and  $l > j - i = -2n - 2$ ). Therefore,

$$\begin{aligned}
 H^l(\text{Hom}_n(B_1, B_2)) &\cong \prod_{i=-n-1}^{n+1} \prod_{j=-n-1}^{n+1} H^l(\text{Hom}_n^{i,j}(B_1, B_2)) \\
 &\cong \prod_{i=-n-1}^n \prod_{j=-n}^{n+1} H^l(\text{Hom}_n^{i,j}(B_1, B_2)) \oplus \prod_{j=-n}^{n+1} H^l(\text{Hom}_n^{n+1,j}(B_1, B_2)) \oplus \prod_{i=-n-1}^n H^l(\text{Hom}_n^{i,-n-1}(B_1, B_2)).
 \end{aligned}
 \tag{4.5}$$

Now, observe that  ${}^nB_1^{-n-1}[n + 1] \cong \prod_{i < n} B_1^i[-i]$  and  ${}^nB_2^{n+1}[-n - 1] \cong \prod_{i > n} B_2^i[-i]$  in  $H^0(\mathcal{C})$ . From this, we deduce that (for  $-n - 1 \leq j \leq n + 1$ )

$$\begin{aligned}
 H^l(\text{Hom}_n^{-n-1,j}(B_1, B_2)) &\cong \text{Hom}_{H^0(\mathcal{C})}({}^nB_1^{-n-1}[n + 1], {}^nB_2^j[l - j]) \\
 &\cong \text{Hom}_{H^0(\mathcal{C})}\left(\prod_{i < -n} B_1^i[-i], {}^nB_2^j[l - j]\right) \cong \prod_{i < -n} \text{Hom}_{H^0(\mathcal{C})}(B_1^i[-i], {}^nB_2^j[l - j])
 \end{aligned}$$

and similarly (for  $-n - 1 \leq i \leq n + 1$ )

$$\begin{aligned}
 H^l(\text{Hom}_n^{i,n+1}(B_1, B_2)) &\cong \text{Hom}_{H^0(\mathcal{C})}({}^nB_1^i[-i], {}^nB_2^{n+1}[l - n - 1]) \\
 &\cong \text{Hom}_{H^0(\mathcal{C})}\left({}^nB_1^i[-i], \prod_{j > n} B_2^j[l - j]\right) \cong \prod_{j > n} \text{Hom}_{H^0(\mathcal{C})}({}^nB_1^i[-i], B_2^j[l - j]).
 \end{aligned}$$

Thus the first summand in equation (4.5) can be written as

$$\prod_{i=-n-1}^n \prod_{j=-n}^{n+1} H^l(\text{Hom}_n^{i,j}(B_1, B_2)) \cong \prod_{i \leq n} \prod_{j \geq -n} \text{Hom}_{n,l}(B_1^i, B_2^j),$$

where (for  $i \leq n$  and  $j \geq -n$ )

$$\text{Hom}_{n,l}(B_1^i, B_2^j) := \begin{cases} \text{Hom}_{H^0(\mathcal{C})}(B_1^i, B_2^j[l + i - j]) & \text{if } l \leq j - i \text{ or } i < -n \text{ or } j > n \\ 0 & \text{if } l > j - i, i \geq -n \text{ and } j \leq n. \end{cases}$$

Using again the fact that  $\text{Hom}_{H^0(\mathcal{C})}(B_1^i, B_2^j[l + i - j]) = 0$  if  $l < j - i$ , we see that for fixed  $i, j \in \mathbb{Z}$ ,

$$\text{Hom}_{n,l}(B_1^i, B_2^j) \cong \begin{cases} \text{Hom}_{H^0(\mathcal{C})}(B_1^i, B_2^j) & \text{if } l = j - i \\ 0 & \text{if } l \neq j - i \end{cases}$$

when  $n \geq |i|, |j|$ .

Reasoning as before, the last two summands in equation (4.5) can be written as

$$\begin{aligned}
 \prod_{j=-n}^{n+1} H^l(\text{Hom}_n^{n+1,j}(B_1, B_2)) &\cong \prod_{j \geq -n} \text{Hom}_{n,l}(B_1, B_2^j), \\
 \prod_{i=-n-1}^n H^l(\text{Hom}_n^{i,-n-1}(B_1, B_2)) &\cong \prod_{i \leq n} \text{Hom}_{n,l}(B_1^i, B_2),
 \end{aligned}$$

where (for  $j \geq -n$  and  $i \leq n$ )

$$\text{Hom}_{n,l}(B_1, B_2^j) := \begin{cases} \text{Hom}_{H^0(\mathcal{E})}({}^n B_1^{n+1}, B_2^j[l+n+1-j]) & \text{if } l < j-n \text{ or } j > n \\ 0 & \text{if } l \geq j-n \text{ and } j \leq n, \end{cases}$$

$$\text{Hom}_{n,l}(B_1^i, B_2) := \begin{cases} \text{Hom}_{H^0(\mathcal{E})}(B_1^i, {}^n B_2^{-n-1}[l+i+n+1]) & \text{if } l < -n-i \text{ or } i < -n \\ 0 & \text{if } l \geq -n-i \text{ and } i \geq -n. \end{cases}$$

Then given  $j \in \mathbb{Z}$  (respectively,  $i \in \mathbb{Z}$ ),  $\text{Hom}_{n,l}(B_1, B_2^j) = 0$  (respectively,  $\text{Hom}_{n,l}(B_1^i, B_2) = 0$ ) when  $n \geq |j|, j - l$  (respectively  $n \geq |i|, -i - l$ ).

Summing up, we have shown that  $H^l(\text{Hom}_n(B_1, B_2))$  is a product of terms, each of which stabilises as  $n$  goes to infinity. Moreover, the only surviving terms in the limit are  $\text{Hom}_{H^0(\mathcal{E})}(B_1^i, B_2^{i+l})$ , which are precisely those appearing in the right-hand side of equation (4.4). From this, it follows that

$$H^l(\text{Holim Hom}_n(B_1, B_2)) \cong \lim_n H^l(\text{Hom}_n(B_1, B_2))$$

and that the right-hand side can be identified with the left-hand side of equation (4.4). This clearly proves the first statement. As for the second, it is enough to note that, by the same argument,

$$\text{Holim Hom}_n(B_1, B_2)^{\leq k} \cong (\text{Holim Hom}_n(B_1, B_2))^{\leq k}.$$

This concludes the proof. □

Now we are going to see how the problem with compositions mentioned in Remark 4.4 can be overcome.

**Lemma 4.7.** *Given  $B_1, B_2, B_3 \in \text{Ob}(\mathcal{B})$  and  $n, k, l \in \mathbb{Z}$  with  $n \geq k, l, 1$ , the composition map given by equation (4.2) restricts to a morphism of complexes*

$$m_n^{k,l} : \text{Hom}_{2n}(B_1, B_2)^{\leq k} \otimes_{\mathbb{k}} \text{Hom}_{2n}(B_2, B_3)^{\leq l} \longrightarrow \text{Hom}_n(B_1, B_3)^{\leq k+l}.$$

*Proof.* Given  $f \in \text{Hom}_{2n}(B_1, B_2)^{\leq k} \subseteq \text{Hom}_{\mathcal{B}}(B_1, B_2)$  and  $g \in \text{Hom}_{2n}(B_2, B_3)^{\leq l} \subseteq \text{Hom}_{\mathcal{B}}(B_2, B_3)$ , we have to prove that  $g \circ f \in \text{Hom}_{\mathcal{B}}(B_1, B_3)$  actually belongs to  $\text{Hom}_n(B_1, B_3)^{\leq k+l}$ . As obviously  $g \circ f \in \text{Hom}_{\mathcal{B}}(B_1, B_3)^{\leq k+l}$ , it is enough to show that  $g \circ f \in \text{Hom}_n(B_1, B_3)$ . We can clearly assume that there exist  $-2n - 1 \leq h, i, j \leq 2n + 1$  such that

$$f \in \text{Hom}_{2n}^{h,i}(B_1, B_2)^{\leq k} \subseteq \text{Hom}_{\mathcal{E}}({}^{2n} B_1^h[-h], {}^{2n} B_2^i[-i]),$$

$$g \in \text{Hom}_{2n}^{i,j}(B_2, B_3)^{\leq l} \subseteq \text{Hom}_{\mathcal{E}}({}^{2n} B_2^i[-i], {}^{2n} B_3^j[-j]).$$

Using the notation of Remark 4.5 (with  $m = 2n$ ), we know that  $g \circ f \in \text{Hom}_{\mathcal{E}}({}^n B_1^{h'}[-h'], {}^n B_3^{j'}[-j'])$ . Thus, to conclude that  $g \circ f \in \text{Hom}_n^{h',j'}(B_1, B_3)$ , it remains to check that  $g \circ f \in \text{Hom}_{\mathcal{B}}(B_1, B_3)^{\leq j'-h'}$  if  $h' \geq -n$  and  $j' \leq n$ . Now, this is certainly true if  $|i| \leq 2n$ , because in that case  $f \in \text{Hom}_{\mathcal{B}}(B_1, B_2)^{\leq i-h}$ ,  $g \in \text{Hom}_{\mathcal{B}}(B_2, B_3)^{\leq j-i}$  and  $i - h + j - i = j - h \leq j' - h'$ . On the other hand, if  $i = -2n - 1$ , then  $f \in \text{Hom}_{\mathcal{B}}(B_1, B_2)^{\leq -2n-1-h}$ ,  $g \in \text{Hom}_{\mathcal{B}}(B_2, B_3)^{\leq l}$  and  $-2n - 1 - h + l \leq -2n - 1 - h' + n \leq j' - h'$ . Similarly, if  $i = 2n + 1$ , then  $f \in \text{Hom}_{\mathcal{B}}(B_1, B_2)^{\leq k}$ ,  $g \in \text{Hom}_{\mathcal{B}}(B_2, B_3)^{\leq j-2n-1}$  and  $k + j - 2n - 1 \leq n + j' - 2n - 1 \leq j' - h'$ . □

For every  $n > 0$ , the natural projection

$$\begin{aligned} \text{Hom}_n(B_1, B_2) &= \prod_{i=-n-1}^{n+1} \prod_{j=-n-1}^{n+1} \text{Hom}_n^{i,j}(B_1, B_2) \\ &\longrightarrow \prod_{i=-n}^n \prod_{j=-n}^n \text{Hom}_n^{i,j}(B_1, B_2) = \prod_{i=-n}^n \prod_{j=-n}^n \text{Hom}_{\mathcal{E}}(B_1^i[-i], B_2^j[-j])^{\leq j-i} \end{aligned}$$

induces, passing to truncations, a morphism of complexes

$$\text{Hom}_n(B_1, B_2)^{\leq k} \longrightarrow \prod_{i=-n}^n \prod_{j=-n}^n \text{Hom}_{\mathcal{E}}(B_1^i[-i], B_2^j[-j])^{\leq \min(k, j-i)}$$

for every  $k \in \mathbb{Z}$ . Composing it with the product of the natural maps

$$\text{Hom}_{\mathcal{E}}(B_1^i[-i], B_2^j[-j])^{\leq \min(k, j-i)} \longrightarrow (\text{Hom}_{\mathcal{E}}(B_1^i[-i], B_2^j[-j])^{\leq \min(k, j-i)})^{\geq j-i}, \tag{4.6}$$

we obtain a morphism of complexes

$$p_n^k : \text{Hom}_n(B_1, B_2)^{\leq k} \longrightarrow \overline{\text{Hom}}_n^k(B_1, B_2) := \prod_{i=-n}^n \prod_{j=-n}^n (\text{Hom}_{\mathcal{E}}(B_1^i[-i], B_2^j[-j])^{\leq \min(k, j-i)})^{\geq j-i}.$$

Observe that for every  $k, l \in \mathbb{Z}$ , there are also morphisms of complexes

$$\overline{m}_n^{k,l} : \overline{\text{Hom}}_{2n}^k(B_1, B_2) \otimes_{\mathbb{Z}} \overline{\text{Hom}}_{2n}^l(B_2, B_3) \longrightarrow \overline{\text{Hom}}_{2n}^{k+l}(B_1, B_3) \longrightarrow \overline{\text{Hom}}_n^{k+l}(B_1, B_3),$$

where the first map is induced by composition in  $\mathcal{E}$ , and the second one is the natural projection.

**Lemma 4.8.** *The maps  $p_n^k$  are compatible with compositions, meaning for every  $n, k, l \in \mathbb{Z}$  with  $n \geq k, l, 1$ ,*

$$\begin{array}{ccc} \text{Hom}_{2n}(B_1, B_2)^{\leq k} \otimes_{\mathbb{Z}} \text{Hom}_{2n}(B_2, B_3)^{\leq l} & \xrightarrow{m_n^{k,l}} & \text{Hom}_n(B_1, B_3)^{\leq k+l} \\ \downarrow p_{2n}^k \otimes p_{2n}^l & & \downarrow p_n^{k+l} \\ \overline{\text{Hom}}_{2n}^k(B_1, B_2) \otimes_{\mathbb{Z}} \overline{\text{Hom}}_{2n}^l(B_2, B_3) & \xrightarrow{\overline{m}_n^{k,l}} & \overline{\text{Hom}}_n^{k+l}(B_1, B_3) \end{array}$$

is a commutative diagram of complexes.

*Proof.* As in the proof of Lemma 4.7, let  $f \in \text{Hom}_{2n}^{h,i}(B_1, B_2)^{\leq k}$  and  $g \in \text{Hom}_{2n}^{j,l}(B_2, B_3)^{\leq l}$  for some  $-2n - 1 \leq h, i, j \leq 2n + 1$ : we need to prove that

$$p_n^{k+l}(m_n^{k,l}(f \otimes g)) = \overline{m}_n^{k,l}(p_{2n}^k(f) \otimes p_{2n}^l(g)). \tag{4.7}$$

We can restrict to the case  $-n \leq h, j \leq n$ , since otherwise both sides of equation (4.7) are evidently 0. We claim that the same is true if  $|i| = 2n + 1$ . In fact, if  $i = -2n - 1$ , then  $f \in \text{Hom}_{\mathcal{E}}(B_1, B_2)^{\leq -2n-1-h}$ ,  $g \in \text{Hom}_{\mathcal{E}}(B_2, B_3)^{\leq l}$  and  $-2n - 1 - h + l \leq -2n - 1 - h + n < j - h$ . Similarly, if  $i = 2n + 1$ , then  $f \in \text{Hom}_{\mathcal{E}}(B_1, B_2)^{\leq k}$ ,  $g \in \text{Hom}_{\mathcal{E}}(B_2, B_3)^{\leq j-2n-1}$  and  $k + j - 2n - 1 \leq n + j - 2n - 1 < j - h$ . Finally, if  $|i| \leq 2n$ , we can assume that  $f$  and  $g$  are homogeneous, say of degrees  $d$  and  $e$ . We must have  $d \leq i - h$  and  $e \leq j - i$  (whence  $d + e \leq j - h$ ), and again equation (4.7) becomes  $0 = 0$  unless  $d + e = j - h$ . Clearly  $d + e = j - h$  implies  $d = i - h$  and  $e = j - i$ , in which case it is straightforward to see that equation (4.7) is satisfied. □



### 4.3. Homotopy limits of sequences

As the name suggests, homotopy limits are unique only up to homotopy. And there are multiple ways to make enhanced versions of them—we already met this in Section 3.3, where the special case of homotopy pullbacks was discussed in some detail. We remind the reader: in Section 3.3, the approach was to impose a model structure on some ambient category and, with respect to this model structure, do some fibrant replacement. In this section, we want to lay the groundwork for the way we will treat homotopy limits of countable sequences, and the method will be different. It will be based on a (dual) version of Milnor’s mapping telescope.

We start in somewhat greater generality; the sequences will come later. Let  $A_1, A_2, A_3$  be cochain complexes of  $\mathbb{k}$ -modules, and let  $\mu : A_1 \otimes A_2 \rightarrow A_3$  be a cochain map, which we should think of as the composition. Suppose further that for  $i \in \{1, 2, 3\}$ , we are given cochain maps  $\phi_i : A_i \rightarrow A_i$  such that the square below commutes:

$$\begin{array}{ccc} A_1 \otimes A_2 & \xrightarrow{\mu} & A_3 \\ \phi_1 \otimes \phi_2 \downarrow & & \downarrow \phi_3 \\ A_1 \otimes A_2 & \xrightarrow{\mu} & A_3. \end{array}$$

Now, for  $i \in \{1, 2, 3\}$ , we define

$$\tilde{A}_i := \text{Cone}(A_i \xrightarrow{\text{id} - \phi_i} A_i)[-1].$$

And the composition map  $\tilde{\mu} : \tilde{A}_1 \otimes \tilde{A}_2 \rightarrow \tilde{A}_3$  is set to be the composite

$$\begin{array}{c} \text{Cone}\left(A_1 \xrightarrow{\text{id} - \phi_1} A_1\right)[-1] \otimes \text{Cone}\left(A_2 \xrightarrow{\text{id} - \phi_2} A_2\right)[-1] \\ \downarrow \text{truncation} \\ \text{Cone}\left(A_1 \otimes A_2 \xrightarrow{\Psi} (A_1 \otimes A_2) \oplus (A_1 \otimes A_2)\right)[-1] \\ \downarrow \Theta \\ \text{Cone}\left(A_3 \xrightarrow{\text{id} - \phi_3} A_3\right)[-1]. \end{array}$$

The truncation is the obvious map; the tensor product of two mapping cones is the total complex of a complex of three terms, and we truncate the term on the right. The map  $\Psi$  is also obvious, meaning what is left from the tensor product of two mapping cones after truncation: we take the morphisms  $(\text{id} - \phi_1) \otimes \text{id} : A_1 \otimes A_2 \rightarrow A_1 \otimes A_2$  and  $\text{id} \otimes (\text{id} - \phi_2) : A_1 \otimes A_2 \rightarrow A_1 \otimes A_2$  and combine them to form a single map  $\Psi : A_1 \otimes A_2 \rightarrow (A_1 \otimes A_2) \oplus (A_1 \otimes A_2)$ . And finally, the map  $\Theta$  is obtained as the map deduced from the commutative square below by taking the mapping cones of the horizontal maps:

$$\begin{array}{ccc} A_1 \otimes A_2 & \xrightarrow{\Psi} & (A_1 \otimes A_2) \oplus (A_1 \otimes A_2) \\ \mu \downarrow & & \downarrow (\mu, \mu \circ (\phi_1 \otimes \text{id})) \\ A_3 & \xrightarrow{\text{id} - \phi_3} & A_3. \end{array}$$

**Remark 4.9.** Suppose now that  $A_1, A_2$  and  $A_3$  are inverse sequences of cochain complexes of  $\mathbb{k}$ -modules. That is, for any integer  $n > 0$  and for  $i \in \{1, 2, 3\}$ , we are given a cochain complex  $A_{i,n}$ . These come

with multiplication maps  $\mu_n : A_{1,n} \otimes A_{2,n} \rightarrow A_{3,n}$  and sequence maps  $\phi_{i,n} : A_{i,n+1} \rightarrow A_{i,n}$ , and for each  $n$ , the square below commutes:

$$\begin{array}{ccc} A_{1,n+1} \otimes A_{2,n+1} & \xrightarrow{\mu_{n+1}} & A_{3,n+1} \\ \phi_{1,n} \otimes \phi_{2,n} \downarrow & & \downarrow \phi_{3,n} \\ A_{1,n} \otimes A_{2,n} & \xrightarrow{\mu_n} & A_{3,n} \end{array}$$

For  $i \in \{1, 2, 3\}$ , define  $\widehat{A}_i := \prod_{n>0} A_{i,n}$ . The multiplication map  $\widehat{\mu} : \widehat{A}_1 \otimes \widehat{A}_2 \rightarrow \widehat{A}_3$  is the composite

$$\left( \prod_{n>0} A_{1,n} \right) \otimes \left( \prod_{n>0} A_{2,n} \right) \longrightarrow \prod_{n>0} (A_{1,n} \otimes A_{2,n}) \xrightarrow{\prod_{n>0} \mu_n} \prod_{n>0} A_{3,n}.$$

If we let  $\phi_i : \widehat{A}_i \rightarrow \widehat{A}_i$  be the composite

$$\prod_{n>0} A_{i,n} \xrightarrow{\text{projection}} \prod_{n>0} A_{i,n+1} \xrightarrow{\prod_{n>0} \phi_{i,n}} \prod_{n>0} A_{i,n},$$

then the square

$$\begin{array}{ccc} \widehat{A}_1 \otimes \widehat{A}_2 & \xrightarrow{\widehat{\mu}} & \widehat{A}_3 \\ \phi_1 \otimes \phi_2 \downarrow & & \downarrow \phi_3 \\ \widehat{A}_1 \otimes \widehat{A}_2 & \xrightarrow{\widehat{\mu}} & \widehat{A}_3 \end{array}$$

commutes. Then setting

$$\widetilde{A}_i := \text{Cone} \left( \widehat{A}_i \xrightarrow{\text{id} - \phi_i} \widehat{A}_i \right) [-1],$$

the discussion preceding the remark showed us how to construct the composition  $\widetilde{\mu} : \widetilde{A}_1 \otimes \widetilde{A}_2 \rightarrow \widetilde{A}_3$ .

Using this, we can give the following.

**Definition 4.10.** The homotopy limit of a sequence  $A = \{A_n\}$  of complexes of  $\mathbb{k}$ -modules is the complex

$$\text{Holim } A_n := \text{Cone} \left( \prod_{n>0} A_n \xrightarrow{\text{id} - \phi} \prod_{n>0} A_n \right) [-1].$$

Remark 4.9 also showed us how, given three inverse sequences of complexes of  $\mathbb{k}$ -modules  $A_1, A_2$  and  $A_3$  and compatible multiplications  $\mu_n : A_{1,n} \otimes A_{2,n} \rightarrow A_{3,n}$ , we can assemble them to a multiplication map

$$\text{Holim } \mu_n : \text{Holim } A_{1,n} \otimes \text{Holim } A_{2,n} \longrightarrow \text{Holim } A_{3,n}.$$

#### 4.4. Relating $\mathcal{V}$ and $\mathcal{B}$

We have been assembling a sequence of technical lemmas, and the time has come to use them. In this section, we will prove

**Proposition 4.11.** *There exists a morphism  $u: \mathcal{V} \rightarrow \mathcal{B}$  in **Hqe** such that the exact functor  $H^0(u): H^0(\mathcal{V}) \cong \mathbf{V}^2(\mathcal{A}) \rightarrow H^0(\mathcal{B}) \cong \mathbf{B}^2(\mathcal{A})$  (see Remark 4.3) can be identified with the natural functor  $\mathbf{V}^2(\mathcal{A}) \rightarrow \mathbf{B}^2(\mathcal{A})$ .*

The proof will occupy the remainder of this section. We will begin with the dg category  $\mathcal{B}$  and gradually produce a zigzag of dg functors that compose (in the category **Hqe**) to our desired map  $u: \mathcal{V} \rightarrow \mathcal{B}$ .

*Step 1.* The dg category  $\mathcal{B}'$  has the same objects as  $\mathcal{B}$ , and the dg functor  $\mathcal{B} \rightarrow \mathcal{B}'$  is the identity on objects. The Hom-complexes in the dg category  $\mathcal{B}'$ , as well as the dg functor  $\mathcal{B} \rightarrow \mathcal{B}'$ , are specified by giving the cochain map  $\text{Hom}_{\mathcal{B}}(B_1, B_2) \rightarrow \text{Hom}_{\mathcal{B}'}(B_1, B_2)$  for every pair of objects  $B_1, B_2 \in \mathcal{B}$ . We declare this to be the natural cochain map

$$\text{Hom}_{\mathcal{B}}(B_1, B_2) \longrightarrow \text{Holim Hom}_{\mathcal{B}}(B_1, B_2),$$

where on the right, we mean the homotopy limit, in the sense of Section 4.3, of the inverse sequence

$$\cdots \longrightarrow \text{Hom}_{\mathcal{B}}(B_1, B_2) \xrightarrow{\text{id}} \text{Hom}_{\mathcal{B}}(B_1, B_2) \xrightarrow{\text{id}} \text{Hom}_{\mathcal{B}}(B_1, B_2) \cdots$$

The composition law in the category  $\mathcal{B}'$  giving the map

$$\text{Hom}_{\mathcal{B}'}(B_1, B_2) \otimes \text{Hom}_{\mathcal{B}'}(B_2, B_3) \longrightarrow \text{Hom}_{\mathcal{B}'}(B_1, B_3)$$

is as in Remark 4.9 and Definition 4.10.

It is obvious that the dg functor  $\mathcal{B} \rightarrow \mathcal{B}'$  is a quasi-equivalence.

*Step 2.* In this step, we will produce a dg functor  $\mathcal{B}'' \rightarrow \mathcal{B}'$ , with  $\mathcal{B}'$  as in Step 1. Let us start with the following useful definition.

**Definition 4.12.** Let  $\mathcal{S}$  be the set of functions  $f: \mathbb{N} \rightarrow \mathbb{N} \cup \{0\}$  satisfying

- (i)  $f$  is nondecreasing, meaning  $f(n) \leq f(n + 1)$  for all  $n \in \mathbb{N}$ .
- (ii)  $f(n) \rightarrow \infty$  as  $n \rightarrow \infty$ .

We turn  $\mathcal{S}$  into a partially ordered set by setting  $f \leq f'$  if  $f(n) \leq f'(n)$  for all  $n \in \mathbb{N}$ .

Back in Section 4.2, we introduced the subcomplexes  $\text{Hom}_n(B_1, B_2) \subset \text{Hom}_{\mathcal{B}}(B_1, B_2)$  for  $B_1, B_2$  objects of  $\mathcal{B}$  and for  $n \in \mathbb{N}$ . We extend this definition now, allowing  $n = 0$ , by declaring  $\text{Hom}_0(B_1, B_2) := \text{Hom}_{\mathcal{B}}(B_1, B_2)$ .

Given a pair of objects  $B_1, B_2 \in \mathcal{B}$ , a function  $f \in \mathcal{S}$  and an integer  $k > 0$ , we can combine the information recalled in the discussion just prior to Step 2 to form the inverse sequence

$$\cdots \longrightarrow \text{Hom}_{f(3)}(B_1, B_2)^{\leq k} \longrightarrow \text{Hom}_{f(2)}(B_1, B_2)^{\leq k} \longrightarrow \text{Hom}_{f(1)}(B_1, B_2)^{\leq k} \cdots$$

If  $f \leq g$  are elements of  $\mathcal{S}$ , then there is a natural map of inverse sequences

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \text{Hom}_{g(3)}(B_1, B_2)^{\leq k} & \longrightarrow & \text{Hom}_{g(2)}(B_1, B_2)^{\leq k} & \longrightarrow & \text{Hom}_{g(1)}(B_1, B_2)^{\leq k} \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & \text{Hom}_{f(3)}(B_1, B_2)^{\leq k} & \longrightarrow & \text{Hom}_{f(2)}(B_1, B_2)^{\leq k} & \longrightarrow & \text{Hom}_{f(1)}(B_1, B_2)^{\leq k}, \end{array}$$

which allows us to view the construction for fixed  $B_1, B_2, k$  as a functor from  $\mathcal{S}^\circ$  to the category of inverse sequences of cochain complexes. We can now form the dg category  $\mathcal{B}''$ ; the objects are identical to those of  $\mathcal{B}$ . For a pair of objects  $B_1, B_2 \in \mathcal{B}$ , we declare

$$\text{Hom}_{\mathcal{B}''}(B_1, B_2) := \text{colim}_{f \in \mathcal{S}^\circ, k \rightarrow \infty} \left( \text{Holim}_{n \rightarrow \infty} \text{Hom}_{f(n)}(B_1, B_2)^{\leq k} \right).$$

This means for fixed  $f \in \mathcal{S}$  and  $k \in \mathbb{N}$ , we take the homotopy inverse limit of the sequence depicted above. And as this is contravariantly functorial in  $f \in \mathcal{S}$  and covariantly functorial in  $k \in \mathbb{N}$ , we can form the (ordinary) colimit of these cochain complexes. Moreover, for each  $f \in \mathcal{S}$  and  $k \in \mathbb{N}$ , there is an obvious map of inverse systems

$$\begin{array}{ccccc} \cdots & \longrightarrow & \text{Hom}_{f(3)}(B_1, B_2)^{\leq k} & \longrightarrow & \text{Hom}_{f(2)}(B_1, B_2)^{\leq k} & \longrightarrow & \text{Hom}_{f(1)}(B_1, B_2)^{\leq k} \\ & & \downarrow & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & \text{Hom}_{\mathcal{B}}(B_1, B_2) & \xrightarrow{\text{id}} & \text{Hom}_{\mathcal{B}}(B_1, B_2) & \xrightarrow{\text{id}} & \text{Hom}_{\mathcal{B}}(B_1, B_2) \end{array}$$

and, taking homotopy inverse limits, we deduce a map

$$\text{Holim}_{n \rightarrow \infty} \text{Hom}_{f(n)}(B_1, B_2)^{\leq k} \longrightarrow \text{Holim}_{n \rightarrow \infty} \text{Hom}_{\mathcal{B}}(B_1, B_2) = \text{Hom}_{\mathcal{B}'}(B_1, B_2)$$

with  $\mathcal{B}'$  the dg category of Step 1. And as this map is compatible with increasing  $k \in \mathbb{N}$  and decreasing  $f \in \mathcal{S}$ , it gives rise to a cochain map

$$\text{Hom}_{\mathcal{B}''}(B_1, B_2) = \text{colim}_{f \in \mathcal{S}^\circ, k \rightarrow \infty} \left( \text{Holim}_{n \rightarrow \infty} \text{Hom}_{f(n)}(B_1, B_2)^{\leq k} \right) \longrightarrow \text{Hom}_{\mathcal{B}'}(B_1, B_2).$$

This defines for us what the dg functor  $\mathcal{B}'' \rightarrow \mathcal{B}'$  does on Hom-complexes.

It remains to deal with composition. Suppose  $f, g \in \mathcal{S}$  and  $k, l \in \mathbb{N}$  are given. We form  $h \in \mathcal{S}$  as follows

$$h(n) := \begin{cases} 0 & \text{unless } \min(f(n), g(n)) > \max(2k, 2l) \\ \lfloor \frac{\min(f(n), g(n))}{2} \rfloor & \text{otherwise} \end{cases}$$

where the symbol  $\lfloor x \rfloor$  means the integer part of  $x$ ; that is, the function  $\lfloor - \rfloor$  takes a real number  $x$  to the largest integer not larger than  $x$ . With this choice, we have that either  $h(n) = 0$  or  $h(n) \geq \max(k, l, 1)$ , and  $f(n), g(n)$  are both  $\geq 2h(n)$ . Lemma 4.7 tells us that the composition

$$\text{Hom}_{f(n)}(B_1, B_2)^{\leq k} \otimes_{\mathbb{k}} \text{Hom}_{g(n)}(B_2, B_3)^{\leq l} \longrightarrow \text{Hom}_{h(n)}(B_1, B_3)^{\leq k+l}$$

is well-defined, and Remark 4.9 and Definition 4.10 allow us to pass to the homotopy inverse limits, producing a map

$$\begin{array}{c} (\text{Holim}_{n \rightarrow \infty} \text{Hom}_{f(n)}(B_1, B_2)^{\leq k}) \otimes_{\mathbb{k}} (\text{Holim}_{n \rightarrow \infty} \text{Hom}_{g(n)}(B_2, B_3)^{\leq l}) \\ \downarrow \\ \text{Holim}_{n \rightarrow \infty} \text{Hom}_{h(n)}(B_1, B_3)^{\leq k+l}. \end{array}$$

Now, passing to the colimit over  $f, g \in \mathcal{S}$  and  $k, l \in \mathbb{N}$ , we produce the composition map

$$\text{Hom}_{\mathcal{B}''}(B_1, B_2) \otimes \text{Hom}_{\mathcal{B}''}(B_2, B_3) \longrightarrow \text{Hom}_{\mathcal{B}''}(B_1, B_3).$$

*Step 3.* In this step, we will produce a dg functor  $\mathcal{B}'' \rightarrow \overline{\mathcal{B}}$ , with  $\mathcal{B}''$  as in Step 2. On objects, this dg functor is the identity. We need to explain, for every pair of objects  $B_1, B_2 \in \mathcal{B}$ , the map  $\text{Hom}_{\mathcal{B}''}(B_1, B_2) \rightarrow \text{Hom}_{\overline{\mathcal{B}}}(B_1, B_2)$ .

The idea is simple enough. In the paragraphs preceding Lemma 4.8, we introduced the maps  $p_n^k : \text{Hom}_n(B_1, B_2)^{\leq k} \rightarrow \overline{\text{Hom}}_n(B_1, B_2)^{\leq k}$ . Back then, we assumed  $n \geq 1$ . In Step 2, we extended the

definition of  $\text{Hom}_n(B_1, B_2)^{\leq k}$  to  $n = 0$  by setting  $\text{Hom}_0(B_1, B_2)^{\leq k} = \text{Hom}_{\mathcal{B}}(B_1, B_2)^{\leq k}$ , and now we extend the definition of  $\overline{\text{Hom}}_n(B_1, B_2)^{\leq k}$  to  $n = 0$  by setting  $\overline{\text{Hom}}_0(B_1, B_2)^{\leq k} = 0$ . Now let  $B_1, B_2 \in \mathcal{B}$ ,  $f \in \mathcal{S}$  and  $k \in \mathbb{N}$  be given; the maps  $p_{f(i)}^k$  provide morphisms of inverse sequences

$$\begin{array}{ccccc} \cdots & \longrightarrow & \text{Hom}_{f(3)}(B_1, B_2)^{\leq k} & \longrightarrow & \text{Hom}_{f(2)}(B_1, B_2)^{\leq k} & \longrightarrow & \text{Hom}_{f(1)}(B_1, B_2)^{\leq k} \\ & & \downarrow p_{f(3)}^k & & \downarrow p_{f(2)}^k & & \downarrow p_{f(1)}^k \\ \cdots & \longrightarrow & \overline{\text{Hom}}_{f(3)}(B_1, B_2)^{\leq k} & \longrightarrow & \overline{\text{Hom}}_{f(2)}(B_1, B_2)^{\leq k} & \longrightarrow & \overline{\text{Hom}}_{f(1)}(B_1, B_2)^{\leq k}. \end{array}$$

And the map  $\text{Hom}_{\mathcal{B}''}(B_1, B_2) \rightarrow \text{Hom}_{\overline{\mathcal{B}}}(B_1, B_2)$  is obtained by taking first homotopy inverse limits and then colimits, as in

$$\begin{array}{c} \text{colim}_{f \in \mathcal{S}^\circ, k \rightarrow \infty} \left( \text{Holim}_{n \rightarrow \infty} \text{Hom}_{f(n)}(B_1, B_2)^{\leq k} \right) \\ \downarrow \text{colim}_{f \in \mathcal{S}^\circ, k \rightarrow \infty} \left( \text{Holim}_{n \rightarrow \infty} p_{f(n)}^k \right) \\ \text{colim}_{f \in \mathcal{S}^\circ, k \rightarrow \infty} \left( \text{Holim}_{n \rightarrow \infty} \overline{\text{Hom}}_{f(n)}(B_1, B_2)^{\leq k} \right). \end{array}$$

The composition law on the dg category  $\overline{\mathcal{B}}$ , as well as the fact that the map  $\text{Hom}_{\mathcal{B}''}(B_1, B_2) \rightarrow \text{Hom}_{\overline{\mathcal{B}}}(B_1, B_2)$  respects composition, can be deduced by combining Lemma 4.8 with Remark 4.9 and Definition 4.10.

From Lemma 4.6, we learn that the dg functor  $\mathcal{B}'' \rightarrow \overline{\mathcal{B}}$  is a quasi-equivalence.

Step 4. In this step, we will produce a dg functor  $\tilde{\mathcal{B}} \rightarrow \overline{\mathcal{B}}$ . As in the previous steps, the functor is the identity on objects. We define  $\text{Hom}_{\tilde{\mathcal{B}}}(B_1, B_2)$  by the formula

$$\text{Hom}_{\tilde{\mathcal{B}}}(B_1, B_2) := \prod_{i=-\infty}^{\infty} \prod_{j=-\infty}^{\infty} (\text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j])^{\leq j-i})^{\geq j-i},$$

and we need to explain the map  $\text{Hom}_{\tilde{\mathcal{B}}}(B_1, B_2) \rightarrow \text{Hom}_{\overline{\mathcal{B}}}(B_1, B_2)$ .

Now recall that by definition (see just before Lemma 4.8),

$$\overline{\text{Hom}}_n(B_1, B_2)^{\leq k} := \left( \prod_{i=-n}^n \prod_{j=-n}^n (\text{Hom}_{\mathcal{C}}(B_1^i[-i], B_2^j[-j])^{\leq j-i})^{\geq j-i} \right)^{\leq k}.$$

Hence there is an obvious map

$$\text{Hom}_{\tilde{\mathcal{B}}}(B_1, B_2)^{\leq k} \longrightarrow \overline{\text{Hom}}_n(B_1, B_2)^{\leq k},$$

which is just the functor  $(-)^{\leq k}$  applied to the projection from the large product to the smaller one. For any  $f \in \mathcal{S}$ , this induces a map

$$\text{Hom}_{\tilde{\mathcal{B}}}(B_1, B_2)^{\leq k} \longrightarrow \text{Holim}_{n \rightarrow \infty} \overline{\text{Hom}}_{f(n)}(B_1, B_2)^{\leq k}.$$

Now, taking the colimit over  $f \in \mathcal{S}$  and as  $k \rightarrow \infty$  produces a map

$$\text{Hom}_{\tilde{\mathcal{B}}}(B_1, B_2) \longrightarrow \text{colim}_{f \in \mathcal{S}^\circ, k \rightarrow \infty} \left( \text{Holim}_{n \rightarrow \infty} \overline{\text{Hom}}_{f(n)}(B_1, B_2)^{\leq k} \right),$$

which is our definition of the morphism  $\text{Hom}_{\tilde{\mathcal{B}}}(B_1, B_2) \rightarrow \text{Hom}_{\overline{\mathcal{B}}}(B_1, B_2)$ .

It is easy to check that the map is compatible with composition and hence defines a dg functor  $\widetilde{\mathcal{B}} \rightarrow \overline{\mathcal{B}}$ . And it is obvious that this map is a quasi-equivalence.

This concludes the proof of Proposition 4.11, since  $\widetilde{\mathcal{B}}$  is manifestly quasi-equivalent to  $\mathcal{V}$  under the natural dg functor  $\widetilde{\mathcal{B}} \rightarrow \mathcal{V}$ . For later use, let us summarise here the sequence of dg categories and dg functors constructed along the proof

$$\begin{array}{ccccc}
 & \widetilde{\mathcal{B}} & & \mathcal{B}'' & & \mathcal{B} & (4.8) \\
 & \swarrow \bar{F} & & \swarrow F'' & & \swarrow F & \\
 \mathcal{V} & & \overline{\mathcal{B}} & & \mathcal{B}' & & 
 \end{array}$$

where all dg functors but  $F' : \mathcal{B}'' \rightarrow \mathcal{B}'$  are quasi-equivalences.

### 5. Uniqueness for $\mathbf{D}^2(\mathcal{A})$

This section is devoted to the proof of Theorem A (1). It requires some technical observations that are contained in Section 5.2. Some straightforward but interesting applications are discussed in Section 5.4.

#### 5.1. A brief summary of the setting

Let  $\mathcal{A}$  be an abelian category, and let  $(\mathcal{C}, E)$  be an enhancement of  $\mathbf{D}^2(\mathcal{A})$ . In view of Example 3.16, Theorem A (1) will be proved once we show that there is an isomorphism between  $\mathcal{C}$  and  $\mathbf{D}_{\mathbf{dg}}^2(\mathcal{A})$  in **Hqe**.

Let us first define a morphism between these dg categories. One may construct the dg categories  $\mathcal{V} = \mathcal{V}^2(\mathcal{A})$  and  $\mathcal{B} = \mathcal{B}^2(\mathcal{A})$  as in Section 4, which, in view of Remark 4.3, come with isomorphisms in **Hqe**

$$f^K : \text{Perf}(\mathcal{V}) \xrightarrow{\sim} \mathbf{C}_{\mathbf{dg}}^2(\mathcal{A}), \quad f^D : \text{Perf}(\mathcal{B}) \xrightarrow{\sim} \mathcal{C}.$$

Furthermore, by Proposition 4.11, there is a morphism  $u \in \text{Hom}_{\mathbf{Hqe}}(\mathcal{V}, \mathcal{B})$  such that  $H^0(u)$  is the natural functor from  $\mathbf{V}^2(\mathcal{A})$  to  $\mathbf{B}^2(\mathcal{A})$ . Consider then the morphism in **Hqe**

$$f := \text{Ind}(u) : \text{Perf}(\mathcal{V}) \rightarrow \text{Perf}(\mathcal{B}).$$

As a conclusion, we have the exact functors  $F_1, F_2 : \mathbf{K}^2(\mathcal{A}) \rightarrow \mathbf{D}^2(\mathcal{A})$  defined as follows:

$$F_1 := E \circ H^0(f^D \circ f \circ (f^K)^{-1}) \quad F_2 := Q, \tag{5.1}$$

where  $Q$  is the quotient functor in equation (1.1). Set, for later use,

$$g := f^D \circ f \circ (f^K)^{-1} : \mathbf{C}_{\mathbf{dg}}^2(\mathcal{A}) \rightarrow \mathcal{C}. \tag{5.2}$$

The following is clear from the definitions.

**Lemma 5.1.** *In the setting above, there is a natural isomorphism*

$$\theta : F_1|_{\mathbf{V}^2(\mathcal{A})} \xrightarrow{\sim} F_2|_{\mathbf{V}^2(\mathcal{A})}$$

of exact functors.

5.2. Some preliminary results

We discuss a general result that applies nicely to the setting in the previous section.

**Proposition 5.2.** Assume that  $\mathcal{A}$  is an abelian category and that  $G_1, G_2 : \mathbf{K}^?(\mathcal{A}) \rightarrow \mathbf{D}^?(\mathcal{A})$  are exact functors such that

- (i) There is a natural isomorphism  $\theta : G_1|_{\mathbf{V}^?(\mathcal{A})} \xrightarrow{\sim} G_2|_{\mathbf{V}^?(\mathcal{A})}$ .
- (ii) Suppose  $a \leq b$  are integers. If  $V^* \in \text{Ob}(\mathbf{V}^?(\mathcal{A}))$  is such that  $V^i = 0$  for all  $i \notin [a, b]$ , then  $G_1(V^*) \cong G_2(V^*) \in \text{Ob}(\mathbf{D}(\mathcal{A})^{\leq b} \cap \mathbf{D}(\mathcal{A})^{\geq a})$ .

Then for every  $A^* \in \text{Ob}(\mathbf{K}^?(\mathcal{A}))$ , there exists an isomorphism  $\tilde{\theta}_{A^*} : G_1(A^*) \xrightarrow{\sim} G_2(A^*)$  such that the following square commutes in  $\mathbf{D}^?(\mathcal{A})$

$$\begin{array}{ccc} G_1(V^*) & \xrightarrow{G_1(h)} & G_1(A^*) \\ \theta_{V^*} \downarrow & & \downarrow \tilde{\theta}_{A^*} \\ G_2(V^*) & \xrightarrow{G_2(h)} & G_2(A^*) \end{array}$$

for every  $V^* \in \text{Ob}(\mathbf{V}^?(\mathcal{A}))$  and every morphism  $h : V^* \rightarrow A^*$  of  $\mathbf{K}^?(\mathcal{A})$ .

*Proof.* Given  $A^* \in \mathbf{K}^?(\mathcal{A})$ , we denote by  $K^i$  the kernel of the differential  $d^i : A^i \rightarrow A^{i+1}$  and by  $\alpha^i : A^{i-1} \rightarrow K^i$  the natural factorisation of the differential  $d^{i-1} : A^{i-1} \rightarrow A^i$ .

Consider the commutative square

$$\begin{array}{ccc} G_1\left(\bigoplus_{i \in \mathbb{Z}} A^{i-1}[-i]\right) & \xrightarrow{G_1\left(\bigoplus_{i \in \mathbb{Z}} \alpha^i[-i]\right)} & G_1\left(\bigoplus_{i \in \mathbb{Z}} K^i[-i]\right) \\ \theta \downarrow & & \downarrow \theta \\ G_2\left(\bigoplus_{i \in \mathbb{Z}} A^{i-1}[-i]\right) & \xrightarrow{G_2\left(\bigoplus_{i \in \mathbb{Z}} \alpha^i[-i]\right)} & G_2\left(\bigoplus_{i \in \mathbb{Z}} K^i[-i]\right), \end{array}$$

where the vertical maps are isomorphisms by (i). Of course, it could be split into the direct sum of two similar commutative diagrams, where the direct sums are indexed, respectively, over  $i \geq 1$  and  $i \leq 0$ . By (ii), Proposition 1.8 and Proposition 1.7, we obtain two commutative diagrams

$$\begin{array}{ccc} \prod_{i \geq 1} G_1(A^{i-1}[-i]) & \xrightarrow{\prod_{i \geq 1} G_1(\alpha^i[-i])} & \prod_{i \geq 1} G_1(K^i[-i]) \\ \prod_{i \geq 1} \theta \downarrow & & \downarrow \prod_{i \geq 1} \theta \\ \prod_{i \geq 1} G_2(A^{i-1}[-i]) & \xrightarrow{\prod_{i \geq 1} G_2(\alpha^i[-i])} & \prod_{i \geq 1} G_2(K^i[-i]) \end{array}$$

and

$$\begin{array}{ccc} \prod_{i \leq 0} G_1(A^{i-1}[-i]) & \xrightarrow{\prod_{i \leq 0} G_1(\alpha^i[-i])} & \prod_{i \leq 0} G_1(K^i[-i]) \\ \prod_{i \leq 0} \theta \downarrow & & \downarrow \prod_{i \leq 0} \theta \\ \prod_{i \leq 0} G_2(A^{i-1}[-i]) & \xrightarrow{\prod_{i \leq 0} G_2(\alpha^i[-i])} & \prod_{i \leq 0} G_2(K^i[-i]). \end{array}$$

For each  $i \in \mathbb{Z}$ ,  $\theta$  induces by (i) an isomorphism of distinguished triangles

$$\begin{CD} G_1(A^{i-1}[-i]) @>G_1(\alpha^i[-i])>> G_1(K^i[-i]) @>G_1(\varphi^i[-i])>> G_1(\text{Cone}(\alpha^i)[-i]) \\ @V\theta VV @V\theta VV @V\theta'_i VV \\ G_2(A^{i-1}[-i]) @>G_2(\alpha^i[-i])>> G_2(K^i[-i]) @>G_2(\varphi^i[-i])>> G_2(\text{Cone}(\alpha^i)[-i]), \end{CD}$$

where  $\varphi^i$  is the morphism of equation (2.1). By taking products over the integers  $i \geq 1$  and coproducts over  $i \leq 0$ , this produces an isomorphism of distinguished triangles

$$\begin{CD} G_1(\bigoplus_{i \in \mathbb{Z}} A^{i-1}[-i]) @>G_1(\bigoplus_{i \in \mathbb{Z}} \alpha^i[-i])>> G_1(\bigoplus_{i \in \mathbb{Z}} K^i[-i]) @>G_1(\varphi)>> G_1(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]) \\ @V\theta VV @V\theta VV @V\theta' VV \\ G_2(\bigoplus_{i \in \mathbb{Z}} A^{i-1}[-i]) @>G_2(\bigoplus_{i \in \mathbb{Z}} \alpha^i[-i])>> G_2(\bigoplus_{i \in \mathbb{Z}} K^i[-i]) @>G_2(\varphi)>> G_2(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]), \end{CD} \tag{5.3}$$

where  $\varphi := \bigoplus_{i \in \mathbb{Z}} \varphi^i[-i]$  and  $\theta' := \bigoplus_{i \in \mathbb{Z}} \theta'_i$ . Hence the rightmost square in equation (5.3) commutes.

For  $i \in \mathbb{Z}$ , consider now the inclusion  $\rho^i: K^{i-1} \hookrightarrow A^{i-1}$ . The composite  $K^{i-1} \xrightarrow{\rho^i} A^{i-1} \xrightarrow{\alpha^i} K^i$  vanishes in  $\mathcal{A}$ . Thus  $\rho^i$  factors in  $\mathbf{K}^2(\mathcal{A})$  as  $K^{i-1} \xrightarrow{\psi^{i-1}} \text{Cone}(\alpha^i)[-1] \rightarrow A^{i-1}$ , where  $\psi^j$  is the morphism of equation (2.2). From this, we deduce for each  $i \in \mathbb{Z}$  a diagram

$$\begin{CD} G_1(K^{i-1}[-i+1]) @>G_1(\psi^{i-1}[-i+1])>> G_1(\text{Cone}(\alpha^i)[-i]) @>>> G_1(A^{i-1}[-i+1]) \\ @V\theta VV @V\theta'_i VV @V\theta VV \\ G_2(K^{i-1}[-i+1]) @>G_2(\psi^{i-1}[-i+1])>> G_2(\text{Cone}(\alpha^i)[-i]) @>>> G_2(A^{i-1}[-i+1]). \end{CD}$$

Note that if we delete the middle column, the resulting square commutes because of the naturality of the isomorphism  $\theta$ . If we delete the left column, the resulting square commutes by the definition of  $\theta'_i$ . It follows that the difference between the composites in the square on the left is annihilated by the map  $G_2(\text{Cone}(\alpha^i)[-i]) \rightarrow G_2(A^{i-1}[-i+1])$  and hence must factor through  $G_2(K^i[-i])$ . But by (ii), there can be no nonzero map  $G_1(K^{i-1}[-i+1]) \rightarrow G_2(K^i[-i])$ , and hence the square on the left must also commute. Taking the coproduct over  $i \leq 0$  and the product over  $i \geq 1$  and then assembling, we deduce a commutative square

$$\begin{CD} G_1(\bigoplus_{i \in \mathbb{Z}} K^{i-1}[-i+1]) @>G_1(\psi)>> G_1(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]) \\ @V\theta VV @V\theta' VV \\ G_2(\bigoplus_{i \in \mathbb{Z}} K^{i-1}[-i+1]) @>G_2(\psi)>> G_2(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]), \end{CD}$$

where  $\psi := \bigoplus_{i \in \mathbb{Z}} \psi^i[-i]$ .



By putting together this commutative square and rightmost commutative square in equation (5.3), we obtain the commutative square

$$\begin{CD} G_1(\bigoplus_{i \in \mathbb{Z}} K^{i-1}[-i+1]) @>G_1(\varphi+\psi)>> G_1(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]) \\ @V\theta VV @VV\theta' V \\ G_2(\bigoplus_{i \in \mathbb{Z}} K^{i-1}[-i+1]) @>G_2(\varphi+\psi)>> G_2(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]), \end{CD}$$

which can be completed to an isomorphism of distinguished triangles

$$\begin{CD} G_1(\bigoplus_{i \in \mathbb{Z}} K^{i-1}[-i+1]) @>G_1(\varphi+\psi)>> G_1(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]) @>G_1(\sigma)>> G_1(A^*) \\ @V\theta VV @VV\theta' V @VV\tilde{\theta}_{A^*} V \\ G_2(\bigoplus_{i \in \mathbb{Z}} K^{i-1}[-i+1]) @>G_2(\varphi+\psi)>> G_2(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]) @>G_2(\sigma)>> G_2(A^*). \end{CD}$$

Here we use the fact that, as we observed in the proof of Proposition 2.9, the mapping cone of  $\varphi + \psi$  is isomorphic in  $\mathbf{K}(\mathcal{A})$  to  $A^*$ .

Clearly  $\tilde{\theta}_{A^*}$  is an isomorphism; thus, to complete the proof, it remains to prove that the square in the statement commutes.

To this end, observe that both squares in the diagram

$$\begin{CD} G_1(\bigoplus_{i \in \mathbb{Z}} K^{i-1}[-i+1]) @>G_1(\varphi)>> G_1(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]) @>G_1(\sigma)>> G_1(A^*) \\ @V\theta VV @VV\theta' V @VV\tilde{\theta}_{A^*} V \\ G_2(\bigoplus_{i \in \mathbb{Z}} K^{i-1}[-i+1]) @>G_2(\varphi)>> G_2(\bigoplus_{i \in \mathbb{Z}} \text{Cone}(\alpha^i)[-i]) @>G_2(\sigma)>> G_2(A^*) \end{CD} \tag{5.4}$$

commute by construction, and thus the outside square commutes as well. Moreover, any map  $V^* \rightarrow A^*$ , where  $V^* \in \mathbf{V}^2(\mathcal{A})$ , must factor as  $V^* \xrightarrow{\beta} \bigoplus_{i \in \mathbb{Z}} K^i[-i] \xrightarrow{\sigma \circ \varphi} A^*$ , and in the diagram

$$\begin{CD} G_1(V^*) @>G_1(\beta)>> G_1(\bigoplus_{i \in \mathbb{Z}} K^i[-i]) @>G_1(\sigma \circ \varphi)>> G_1(A^*) \\ @V\theta VV @VV\theta V @VV\tilde{\theta}_{A^*} V \\ G_2(V^*) @>G_2(\beta)>> G_2(\bigoplus_{i \in \mathbb{Z}} K^i[-i]) @>G_2(\sigma \circ \varphi)>> G_2(A^*) \end{CD} \tag{5.5}$$

both squares commute: the left-hand square by the naturality of  $\theta$  and the right-hand square (which is the outside square in equation (5.4)) by the observation above. Thus we deduce the commutativity of the outside square in equation (5.5), completing the proof.  $\square$

We now want to apply this result in the setting of Section 5.1.

**Corollary 5.3.** *Let  $\mathcal{A}$  be an abelian category, and let  $F_1, F_2 : \mathbf{K}^2(\mathcal{A}) \rightarrow \mathbf{D}^2(\mathcal{A})$  be as in equation (5.1). Then for every  $A^* \in \text{Ob}(\mathbf{K}^2(\mathcal{A}))$ , there is an isomorphism  $\tilde{\theta}_{A^*} : F_1(A^*) \xrightarrow{\sim} F_2(A^*)$  such that the*

following square commutes in  $\mathbf{D}^?(\mathcal{A})$

$$\begin{array}{ccc} F_1(V^*) & \xrightarrow{F_1(h)} & F_1(A^*) \\ \theta_{V^*} \downarrow & & \downarrow \tilde{\theta}_{A^*} \\ F_2(V^*) & \xrightarrow{F_2(h)} & F_2(A^*) \end{array}$$

for every  $V^* \in \text{Ob}(\mathbf{V}^?(\mathcal{A}))$  and every morphism  $h: V^* \rightarrow A^*$  of  $\mathbf{K}^?(\mathcal{A})$ .

*Proof.* The result is a direct consequence of Proposition 5.2. Indeed, assumption (i) is satisfied thanks to Lemma 5.1. On the other hand, assumption (ii) clearly holds since  $F_2 = Q$ . □

### 5.3. Proof of Theorem A (I)

This part of the argument is very similar to [27, Section 6]. The key differences are our approach to generation for  $\mathbf{D}^?(\mathcal{A})$  and all of its consequences in the previous section. We are in the setting of Section 5.1.

Suppose that  $L^* \in \mathbf{K}^?_{\text{acy}}\mathcal{A}$  is an acyclic object. By Corollary 5.3,  $F_1(L^*) \cong F_2(L^*)$  in  $\mathbf{D}^?(\mathcal{A})$ . Since  $F_2(L^*)$  is zero as  $F_2 = Q$  is the Verdier quotient map, we have  $F_1(L^*) \cong 0$ , whence the functor  $F_1$  must factor through  $Q$ . Let us write this as

$$\mathbf{K}^?(\mathcal{A}) \xrightarrow{Q} \mathbf{D}^?(\mathcal{A}) = \mathbf{K}^?(\mathcal{A})/\mathbf{K}^?_{\text{acy}}(\mathcal{A}) \xrightarrow{F'_1} \mathbf{D}^?(\mathcal{A}).$$

As an immediate consequence, the morphism  $g$  in equation (5.2) must factor through the Drinfeld quotient as follows:

$$\mathbf{C}^?_{\text{dg}}(\mathcal{A}) \longrightarrow \mathbf{D}^?_{\text{dg}}(\mathcal{A}) = \mathbf{C}^?_{\text{dg}}(\mathcal{A})/\mathbf{Acy}^?(\mathcal{A}) \xrightarrow{g'} \mathcal{C}.$$

Note that, by construction, we have

$$F'_1 = E \circ H^0(g'). \tag{5.6}$$

The proof will be complete once we show that  $H^0(g')$  is an equivalence. As  $E$  is an exact equivalence, we are reduced to showing that  $F'_1$  is an equivalence.

Let us prove that  $F'_1$  is fully faithful. By Corollary 2.10, it suffices to show that for any pair of objects  $V_1, V_2 \in \text{Ob}(\mathbf{V}^?(\mathcal{A}))$ , the functor  $F'_1$  induces an isomorphism

$$\text{Hom}_{\mathbf{K}^?(\mathcal{A})/\mathbf{K}^?_{\text{acy}}(\mathcal{A})}(V_1, V_2) \xrightarrow{\sim} \text{Hom}_{\mathbf{D}^?(\mathcal{A})}(F_1(V_1), F_1(V_2)).$$

To prove the injectivity, choose in the category  $\mathbf{K}^?(\mathcal{A})/\mathbf{K}^?_{\text{acy}}(\mathcal{A})$  a morphism  $h: V_1 \rightarrow V_2$  mapping to zero under  $F'_1$ . We may represent  $h$  in  $\mathbf{K}^?(\mathcal{A})$  as

$$\begin{array}{ccc} V_1 & \xrightarrow{a} & A \\ & \searrow & \swarrow \\ & L & \end{array} \quad \begin{array}{ccc} & & V_2 \\ & \swarrow & \searrow \\ & A & \end{array}$$

where  $V_2 \xrightarrow{b} A \rightarrow L$  is a distinguished triangle with  $L \in \mathbf{K}^?_{\text{acy}}(\mathcal{A})$  and where  $h = b^{-1} \circ a$  in  $\mathbf{K}^?(\mathcal{A})/\mathbf{K}^?_{\text{acy}}(\mathcal{A})$ . But then  $0 \cong F'_1(h) = F'_1(b)^{-1} \circ F'_1(a)$ , and we deduce that  $F'_1(a) = F_1(a) \cong 0$ . By

Corollary 5.3, there is an isomorphism  $\tilde{\theta}_A: F_1(A) \xrightarrow{\sim} F_2(A)$  rendering commutative the square

$$\begin{array}{ccc} F_1(V) & \xrightarrow{F_1(a)} & F_1(A) \\ \theta_V \downarrow & & \downarrow \tilde{\theta}_A \\ F_2(V) & \xrightarrow{F_2(a)} & F_2(A). \end{array}$$

The vertical maps are isomorphisms; hence the vanishing of  $F_1(a)$  implies that  $F_2(a) \cong 0$ . Hence  $a$  vanishes in  $\mathbf{K}^2(\mathcal{A})/\mathbf{K}_{\text{acy}}^2(\mathcal{A})$  as  $F_2 = \mathbf{Q}$  is the Verdier quotient. Thus  $h = 0$ .

As for the surjectivity, let  $h: F_1(V_1) \rightarrow F_1(V_2)$  be a morphism in the category  $\mathbf{D}^2(\mathcal{A})$ . By Lemma 5.1,  $\theta$  induces isomorphisms  $F_1(V_i) \cong F_2(V_i)$  for  $i = 1, 2$ . Hence there is a morphism  $h': F_2(V_1) \rightarrow F_2(V_2)$  making the following diagram commutative in  $\mathbf{D}^2(\mathcal{A})$ :

$$\begin{array}{ccc} F_1(V_1) & \xrightarrow{h} & F_1(V_2) \\ \theta_{V_1} \downarrow & & \downarrow \theta_{V_2} \\ F_2(V_1) & \xrightarrow{h'} & F_2(V_2). \end{array}$$

We can represent  $h'$  in  $\mathbf{K}^2(\mathcal{A})$  as

$$\begin{array}{ccc} V_1 & \xrightarrow{a'} & A' \\ & & \swarrow b' \\ & & V_2 \\ & \swarrow & \\ & L' & \end{array}$$

where  $V_2 \xrightarrow{b'} A' \rightarrow L'$  is a distinguished triangle with  $L' \in \mathbf{K}_{\text{acy}}^2(\mathcal{A})$  and where  $h' = (b')^{-1} \circ a'$  in  $\mathbf{D}^2(\mathcal{A})$ . By Corollary 5.3, there is an isomorphism  $\tilde{\theta}_A: F_1(A) \xrightarrow{\sim} F_2(A)$  making the diagram

$$\begin{array}{ccccc} F_1(V_1) & \xrightarrow{F_1(a')} & F_1(A) & \xleftarrow{F_1(b')} & F_1(V_2) \\ \theta_{V_1} \downarrow & & \downarrow \tilde{\theta}_A & & \downarrow \theta_{V_2} \\ F_2(V_1) & \xrightarrow{F_2(a')} & F_2(A) & \xleftarrow{F_2(b')} & F_2(V_2). \end{array}$$

commutative. From this, we deduce that

$$h = \theta_{V_2}^{-1} \circ h' \circ \theta_{V_1} = \theta_{V_2}^{-1} \circ F_2(b')^{-1} \circ F_2(a') \circ \theta_{V_1} = F_1(b')^{-1} \circ F_1(a').$$

Hence, by definition,  $F'_1(h') = h$ .

As for the essential surjectivity of  $F'_1$ , observe that since it is fully faithful, its essential image is a full and thick (because the source category  $\mathbf{D}^2(\mathcal{A})$  is idempotent complete) triangulated subcategory of  $\mathbf{D}^2(\mathcal{A})$  containing  $F_1(\mathbf{V}^2(\mathcal{A}))$ . But  $F_1(\mathbf{V}^2(\mathcal{A})) = F_2(\mathbf{V}^2(\mathcal{A})) = \mathbf{B}^2(\mathcal{A})$  by Lemma 5.1 so, by Corollary 2.10, the essential image of  $F'_1$  coincides with  $\mathbf{D}^2(\mathcal{A})$ .

This concludes the proof of Theorem A (1) as we proved that  $(g')^{-1}$  is an isomorphism between  $\mathcal{E}$  and  $\mathbf{D}_{\text{dg}}^2(\mathcal{A}) = \mathbf{C}_{\text{dg}}^2(\mathcal{A})/\mathbf{Acy}^2(\mathcal{A})$  in **Hqe**.

**Remark 5.4.** It should be noted that our proof shows more:  $\mathbf{D}^2(\mathcal{A})$  has a semi-strongly unique dg enhancement for any abelian category  $\mathcal{A}$ . Indeed as we observed above, by Corollary 5.3, given any object  $A \in \text{Ob}(\mathbf{D}^2(\mathcal{A})) = \text{Ob}(\mathbf{K}^2(\mathcal{A}))$ , there is an isomorphism  $F_1(A) \cong F_2(A)$  in  $\mathbf{D}^2(\mathcal{A})$ . But  $F_1(A) \cong F'_1(A) = E \circ H^0(g')(A)$  by equation (5.6). Moreover,  $F_2(A) = A$  by definition. Hence  $H^0(g')(A) \cong E^{-1}(A)$  for all  $A \in \text{Ob}(\mathbf{D}_{\text{dg}}^2(\mathcal{A}))$ .

**Remark 5.5.** For later use, we can reinterpret the previous proof in terms of the zigzag diagram given by equation (4.8), which describes the morphism  $u: \mathcal{V} \rightarrow \mathcal{B}$  in **Hqe**. First, observe that the morphism  $f = \text{Ind}(u)$  in **Hqe** defined in Section 5.1 is represented by the zigzag diagram

$$\begin{array}{ccccc}
 & \text{Perf}(\widetilde{\mathcal{B}}) & & \text{Perf}(\mathcal{B}'') & & \text{Perf}(\mathcal{B}) & (5.7) \\
 & \swarrow \text{Ind}(\widetilde{F}) & & \swarrow \text{Ind}(F'') & & \swarrow \text{Ind}(F) & \\
 \text{Perf}(\mathcal{V}) & & \text{Perf}(\overline{\mathcal{B}}) & & \text{Perf}(\mathcal{B}') & & \\
 & \searrow \text{Ind}(\overline{F}) & & \searrow \text{Ind}(F') & & \searrow & \\
 & & & & & & 
 \end{array}$$

where all dg functors but  $\text{Ind}(F')$  are quasi-equivalences. By Proposition 3.10, up to taking functorial h-flat resolutions, we can assume without loss of generality that all dg categories in the above diagram are h-flat.

Denoting by  $\mathcal{N}$  the full dg subcategory of  $\text{Perf}(\mathcal{V})$  (which is an enhancement of  $\mathbf{K}^2(\mathcal{A})$  by Remark 4.3) defining an enhancement of  $\mathbf{K}_{\text{acy}}^2(\mathcal{A})$ , our proof above shows that there are full dg subcategories  $\widetilde{\mathcal{S}} \subseteq \text{Perf}(\widetilde{\mathcal{B}})$ ,  $\overline{\mathcal{S}} \subseteq \text{Perf}(\overline{\mathcal{B}})$ ,  $\mathcal{S}'' \subseteq \text{Perf}(\mathcal{B}'')$ ,  $\mathcal{S}' \subseteq \text{Perf}(\mathcal{B}')$  and  $\mathcal{S} \subseteq \text{Perf}(\mathcal{B})$  such that

- (1) we have quasi-equivalences  $\widetilde{\mathcal{S}} \cong \mathcal{N}$ ,  $\widetilde{\mathcal{S}} \cong \overline{\mathcal{S}}$ ,  $\mathcal{S}'' \cong \overline{\mathcal{S}}$  and  $\mathcal{S} \cong \mathcal{S}'$ , which are induced by the corresponding dg functors in equation (5.7);
- (2)  $\mathcal{S}'$  is the smallest full dg subcategory of  $\text{Perf}(\mathcal{B}')$  containing  $\text{Ind}(F')(\mathcal{S}'')$ ;
- (3) we have quasi-equivalences  $\text{Perf}(\widetilde{\mathcal{B}})/\widetilde{\mathcal{S}} \cong \text{Perf}(\mathcal{V})/\mathcal{N}$ ,  $\text{Perf}(\overline{\mathcal{B}})/\overline{\mathcal{S}} \cong \text{Perf}(\mathcal{B}')/\mathcal{S}'$ ,  $\text{Perf}(\mathcal{B}'')/\mathcal{S}'' \cong \text{Perf}(\overline{\mathcal{B}})/\overline{\mathcal{S}}$ ,  $\text{Perf}(\mathcal{B}'')/\mathcal{S}'' \cong \text{Perf}(\mathcal{B}')/\mathcal{S}'$  and  $\text{Perf}(\mathcal{B})/\mathcal{S} \cong \text{Perf}(\mathcal{B}')/\mathcal{S}'$  induced by the dg functors given by equation (5.7) in view of Remark 3.11.

In particular,  $H^0(\mathcal{S}') \cong H^0(\mathcal{S}) \cong 0$ , and we get the commutative diagram of zigzags

$$\begin{array}{ccccccc}
 & \text{Perf}(\widetilde{\mathcal{B}}) & & \text{Perf}(\mathcal{B}'') & & \text{Perf}(\mathcal{B}) & \\
 & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \downarrow \\
 \text{Perf}(\mathcal{V}) & \text{Perf}(\overline{\mathcal{B}})/\overline{\mathcal{S}} & \text{Perf}(\overline{\mathcal{B}}) & \text{Perf}(\mathcal{B}'')/\mathcal{S}'' & \text{Perf}(\mathcal{B}') & \text{Perf}(\mathcal{B})/\mathcal{S} & \\
 \downarrow & \swarrow & \downarrow & \swarrow & \downarrow & \swarrow & \\
 \text{Perf}(\mathcal{V})/\mathcal{N} & & \text{Perf}(\overline{\mathcal{B}})/\overline{\mathcal{S}} & & \text{Perf}(\mathcal{B}')/\mathcal{S}' & & 
 \end{array} \tag{5.8}$$

where the dg functors in the bottom zigzag are quasi-equivalences, the vertical maps are Drinfeld dg quotient functors such that  $\text{Perf}(\mathcal{B}') \rightarrow \text{Perf}(\mathcal{B}')/\mathcal{S}'$  and  $\text{Perf}(\mathcal{B}) \rightarrow \text{Perf}(\mathcal{B})/\mathcal{S}$  are quasi-equivalences.

### 5.4. Applications

The first easy observation is that by Remark 1.3 (and Remark 3.19 (i)), Theorem A (1) immediately implies the following.

**Corollary 5.6.** *If  $\mathcal{A}$  is an abelian category, then  $\mathbf{K}^?( \mathcal{A})$  has a unique enhancement for  $? = b, +, -, \emptyset$ .*

In this light, it would be interesting to investigate further the following problem.

**Question 5.7.** Let  $\mathcal{A}$  be an abelian category. Does  $\mathbf{K}_{\text{acy}}^2(\mathcal{A})$  have a unique enhancement for  $? = b, +, -, \emptyset$ ?

Next, if  $\mathcal{G}$  is a Grothendieck category, then we learnt in Section 2.1 that for  $\alpha$  a sufficiently large cardinal, then  $\mathbf{D}(\mathcal{G})^\alpha \cong \mathbf{D}(\mathcal{G}^\alpha)$ . Hence we get

**Corollary 5.8.** *If  $\mathcal{G}$  is a Grothendieck category and  $\alpha$  is a sufficiently large cardinal, then  $\mathbf{D}(\mathcal{G})^\alpha$  has a unique enhancement.*

When  $\mathcal{G} = \mathbf{Qcoh}(X)$  and  $X$  is an algebraic stack, then this answers the second part of [9, Question 4.7] in the positive. Partial results are in [1].

Continuing the discussion in the geometric setting, let us recall that while quasi-coherent sheaves are defined on any algebraic stack, coherent sheaves seem to be well defined only on locally noetherian algebraic stacks (see, for example, [26, Chapter 15]) or, of course, schemes. But when they are defined, they form an abelian category, and thus we can deduce the following.

**Corollary 5.9.** *If  $X$  is an algebraic stack, then  $\mathbf{D}^?(\mathbf{Qcoh}(X))$  has a unique enhancement for  $? = b, +, -, \emptyset$ . If  $X$  is a scheme or a locally noetherian algebraic stack, then the same is true for  $\mathbf{D}^?(\mathbf{Coh}(X))$ .*

Note that here, apart from Theorem A (1), we are implicitly using Proposition 3.18, as  $\mathbf{Qcoh}(X)$  is not in general small in the given universe. This concludes the proof of the problem posed in [9, Question 4.7].

We conclude this section with a question that we believe to be very natural as it concerns a potentially very interesting generalisation of Theorem A (1).

**Question 5.10.** Let  $\mathcal{A}$  be an abelian category, and let  $\mathcal{B}$  be a Serre subcategory of  $\mathcal{A}$ . Does the full subcategory  $\mathbf{D}_{\mathcal{B}}^?(\mathcal{A})$  of  $\mathbf{D}^?(\mathcal{A})$ , consisting of complexes with cohomology in  $\mathcal{B}$ , have a unique dg enhancement for  $? = b, +, -, \emptyset$ ?

A positive answer to Question 5.10 would immediately imply the uniqueness of dg enhancements for  $\mathbf{D}_{\mathbf{qc}}^?(X)$  in Theorem B. Unfortunately, the techniques developed in this paper seem insufficient to address Question 5.10.

### 5.5. Aside: the ‘realisation’ functor of Beilinson, Bernstein and Deligne

Recall that a *t-structure* on a triangulated category  $\mathcal{T}$  is a pair of full subcategories  $(\mathcal{T}^{\geq 0}, \mathcal{T}^{\leq 0})$  such that

- (i) If  $X \in \text{Ob}(\mathcal{T}^{\leq 0})$  and  $Y \in \text{Ob}(\mathcal{T}^{\geq 0})$ , then  $\text{Hom}_{\mathcal{T}}(X, Y[-1]) = 0$ ;
- (ii)  $\mathcal{T}^{\leq 0}[1] \subseteq \mathcal{T}^{\leq 0}$  and  $\mathcal{T}^{\geq 0}[-1] \subseteq \mathcal{T}^{\geq 0}$  (we set  $\mathcal{T}^{\leq n} := \mathcal{T}^{\leq 0}[-n]$  and  $\mathcal{T}^{\geq n} := \mathcal{T}^{\geq 0}[-n]$ );
- (iii) For any object  $X \in \text{Ob}(\mathcal{T})$  there is a distinguished triangle

$$X^{\leq 0} \longrightarrow X \longrightarrow X^{\geq 1},$$

where  $X^{\leq 0} \in \text{Ob}(\mathcal{T}^{\leq 0})$  and  $X^{\geq 1} \in \text{Ob}(\mathcal{T}^{\geq 1})$ .

The *heart* of a t-structure  $(\mathcal{T}^{\geq 0}, \mathcal{T}^{\leq 0})$  on  $\mathcal{T}$  is the abelian category  $\mathcal{T}^\heartsuit := \mathcal{T}^{\geq 0} \cap \mathcal{T}^{\leq 0}$ . Given an object  $X \in \mathcal{T}$  and an integer  $i$ , we can use the distinguished triangle in (iii) to define the object  $X^{\geq i} \in \text{Ob}(\mathcal{T}^{\geq i})$  as  $(X[i-1]^{\geq 1})[1-i]$ . A t-structure on  $\mathcal{T}$  is *nondegenerate* if the intersections of all the  $\mathcal{T}^{\leq n}$  and of all the  $\mathcal{T}^{\geq n}$  are trivial.

In [4, Proposition 3.1.10], Beilinson, Bernstein and Deligne proved a very interesting result asserting that if  $\mathcal{T}$  is a triangulated category with a t-structure with heart  $\mathcal{T}^\heartsuit$ , under a hypothesis on the existence of filtered derived categories, the natural inclusion  $\mathcal{T}^\heartsuit \subset \mathcal{T}$  extends to an exact functor

$$\text{real}: \mathbf{D}^b(\mathcal{T}^\heartsuit) \longrightarrow \mathcal{T}$$

respecting t-structures. The original proof was relatively complicated, and there has been some literature on this since. But the point here is that the result becomes straightforward by our techniques, which allow for generalisations to  $\mathbf{D}^?(\mathcal{T}^\heartsuit)$ .

As above, let  $\mathcal{T}$  be a triangulated category with a t-structure, and assume we are given an enhancement  $(\mathcal{C}, \mathbf{E})$  of  $\mathcal{T}$ . Let  $\mathbf{V}^?(\mathcal{T}^\heartsuit)$  be as in the opening paragraphs of Section 1.1, and let  $\mathcal{V}^?(\mathcal{T}^\heartsuit)$  be its

enhancement defined at the very beginning of Section 4.1. And finally, assume that there is a morphism  $f \in \text{Hom}_{\mathbf{Hqc}}(\mathcal{V}^?(\mathcal{T}^\heartsuit), \mathcal{E})$  such that the functor  $F := E \circ H^0(f): \mathbf{V}^?(\mathcal{T}^\heartsuit) \rightarrow \mathcal{T}$  satisfies the following properties for every countable collection  $\{T^i : i \in \mathbb{Z}\}$  of objects of  $\mathcal{T}^\heartsuit$ .

- (1)  $F$  takes a finite sum  $\bigoplus_{i=m}^n T^i[-i]$  in  $\mathbf{V}^?(\mathcal{T}^\heartsuit)$  to the object  $\bigoplus_{i=m}^n T^i[-i] \in \mathcal{T}$ , and on morphisms it is the obvious functor.
- (2) If  $? = -, \emptyset$ , then  $F(\bigoplus_{i=-\infty}^n T^i[-i]) \in \mathcal{T}^{\leq n}$ .
- (3) If  $? = +, \emptyset$ , then  $F(\bigoplus_{i=n}^{\infty} T^i[-i]) \in \mathcal{T}^{\geq n}$ .

Consider the morphism  $\tilde{f} := \text{Ind}(f)|_{\text{Perf}(\mathcal{V}^?(\mathcal{T}^\heartsuit))} \in \text{Hom}_{\mathbf{Hqc}}(\text{Perf}(\mathcal{V}^?(\mathcal{T}^\heartsuit)), \text{Perf}(\mathcal{E}))$ . Assuming that  $\mathcal{T}$  is idempotent complete, it induces an exact functor

$$\tilde{F} := E \circ H^0(\tilde{f}): \mathbf{K}^?(\mathcal{T}^\heartsuit) \longrightarrow \mathcal{T},$$

and it is an easy exercise to show that the  $\mathcal{T}^\heartsuit$ -cohomology of  $\tilde{F}(E)$  vanishes for every  $E \in \text{Ob}(\mathbf{K}_{\text{acy}}^?(\mathcal{T}^\heartsuit))$ . Assuming also that the t-structure is nondegenerate, we deduce that  $\tilde{F}(\mathbf{K}_{\text{acy}}^?(\mathcal{T}^\heartsuit)) = 0$ , and the functor  $\tilde{F}$  must factor through

$$\text{real}: \mathbf{D}^?(\mathcal{T}^\heartsuit) \longrightarrow \mathcal{T}.$$

Thus the problem reduces to finding a morphism  $f \in \text{Hom}_{\mathbf{Hqc}}(\mathcal{V}^?(\mathcal{T}^\heartsuit), \mathcal{E})$  with the required properties, and Section 4 is all about methods to do this. The special case  $? = b$  is trivial since the limit arguments disappear. To the best of our knowledge, there is only one other article in the literature that shows how to construct the realisation functor on unbounded or half-bounded derived categories: Virili [48, Sections 5 and 6]. But the hypotheses Virili places on the categories  $\mathcal{T}$  and  $\mathcal{T}^\heartsuit$  are much more restrictive than ours.

### 6. The unseparated and completed derived categories

In this section, we prove Theorem A (2). In Section 6.1, we prove the uniqueness of enhancements for the unseparated derived category of a Grothendieck category, while the proof of the uniqueness of enhancements for the completed derived category of a Grothendieck category is carried out in Section 6.3. This is preceded by a discussion about the basic properties of completed derived categories in Section 6.2. We end this section with some speculations about the uniqueness of dg enhancements for admissible subcategories.

#### 6.1. The unseparated derived category

The uniqueness of dg enhancements for  $\check{\mathbf{D}}(\mathcal{G})$  for  $\mathcal{G}$  a Grothendieck category can be deduced from a general criterion proved in [14].

To state it precisely, let us recall the following general definition.

**Definition 6.1.** Let  $\mathcal{A}$  be a small  $\mathbb{k}$ -linear category considered as a dg category concentrated in degree 0, and let  $\mathcal{T}$  be a triangulated category with small coproducts. An exact functor  $F: \mathcal{D}(\mathcal{A}) \rightarrow \mathcal{T}$  is *right vanishing* if it preserves small coproducts and there exists a full subcategory  $\mathcal{R}$  of  $\mathcal{T}$  with the following properties:

- (R1)  $\mathcal{R}$  is closed under small coproducts.
- (R2)  $\mathcal{R}$  is closed under extensions (meaning if  $X \rightarrow Y \rightarrow Z$  is a distinguished triangle in  $\mathcal{T}$  with  $X, Z \in \mathcal{R}$ , then  $Y \in \mathcal{R}$ , as well).
- (R3)  $F(Y^{\mathcal{A}}(A))[k] \in \mathcal{R}$  for every  $A \in \mathcal{A}$  and every integer  $k < 0$ .
- (R4)  $\text{Hom}_{\mathcal{T}}(F(Y^{\mathcal{A}}(A)), R) = 0$  for every  $A \in \mathcal{A}$  and every  $R \in \mathcal{R}$ .

The following is the revised version of [13, Theorem C], which appeared in the preprint version [14]. It extends [27, Theorem 2.7]. The key notion is well-generation for triangulated categories (see Section 2.1).

**Theorem 6.2.** *Let  $\mathcal{A}$  be a small  $\mathbb{k}$ -linear category considered a dg category concentrated in degree 0, and let  $\mathcal{L}$  be a localising subcategory of  $\mathcal{D}(\mathcal{A})$  such that:*

- (a) *The quotient  $\mathcal{D}(\mathcal{A})/\mathcal{L}$  is a well-generated triangulated category.*
- (b) *The quotient functor  $Q: \mathcal{D}(\mathcal{A}) \rightarrow \mathcal{D}(\mathcal{A})/\mathcal{L}$  is right vanishing.*

*Then  $\mathcal{D}(\mathcal{A})/\mathcal{L}$  has a unique enhancement.*

Let us move to the first part of Theorem A (2). By Theorem 2.6,  $\check{\mathbf{D}}(\mathcal{G})$  is well-generated, so to apply Theorem 6.2, we just need to show that  $\check{\mathbf{D}}(\mathcal{G})$  can be written as a quotient  $\mathcal{D}(\mathcal{A})/\mathcal{L}$  with  $\mathcal{A}$  concentrated in degree 0 and that the quotient functor is right vanishing.

For the given Grothendieck category  $\mathcal{G}$ , take a sufficiently large cardinal  $\alpha$  such that  $\mathcal{G}^\alpha$  is abelian (see Theorem 2.5).

**Remark 6.3.** Even though, by the discussion in Section 3.5, the choice of the universe can be made harmless, it is easy to see that in this case  $\mathcal{G}^\alpha$  is small, since  $\mathcal{G}$  is generated by one object.

By the discussion at the beginning of [23, Section 5], we have an equivalence

$$F: \mathcal{G} \rightarrow \text{Lex}_\alpha((\mathcal{G}^\alpha)^\circ, \text{Mod}(\mathbb{k})) \tag{6.1}$$

given by the assignment  $G \mapsto \text{Hom}_{\mathcal{G}}(-, G)|_{\mathcal{G}^\alpha}$  (see Example 2.3 (ii)). On the other hand, by [23, Proposition 5.4], the natural inclusion  $\text{Lex}_\alpha((\mathcal{G}^\alpha)^\circ, \text{Mod}(\mathbb{k})) \hookrightarrow \text{Mod}(\mathcal{G}^\alpha)$  has a left adjoint

$$Q': \text{Mod}(\mathcal{G}^\alpha) \rightarrow \text{Lex}_\alpha((\mathcal{G}^\alpha)^\circ, \text{Mod}(\mathbb{k})), \tag{6.2}$$

which is an exact functor sending  $Y^{\mathcal{G}^\alpha}(G)$  to itself. Hence, by equations (6.1) and (6.2), we have

$$F^{-1}(Q'(Y^{\mathcal{G}^\alpha}(G))) \cong G \tag{6.3}$$

for all  $G \in \mathcal{G}^\alpha$ .

On the other hand, consider the commutative diagram of natural inclusions

$$\begin{array}{ccc} \mathbf{K}(\mathcal{G}) & \xrightarrow{l_1} & \mathbf{K}(\text{Mod}(\mathcal{G}^\alpha)) \\ \uparrow l_2 & & \uparrow l_3 \\ \check{\mathbf{D}}(\mathcal{G}) & \xrightarrow{l_4} & \mathbf{D}(\text{Mod}(\mathcal{G}^\alpha)). \end{array}$$

By passing to the (left) adjoints, we get the corresponding commutative diagram

$$\begin{array}{ccc} \mathbf{K}(\text{Mod}(\mathcal{G}^\alpha)) & \xrightarrow{Q_1} & \mathbf{K}(\mathcal{G}) \\ \downarrow Q_3 & & \downarrow Q_2 \\ \mathbf{D}(\text{Mod}(\mathcal{G}^\alpha)) & \xrightarrow{Q_4} & \check{\mathbf{D}}(\mathcal{G}). \end{array}$$

If  $K: \mathcal{E}^\alpha \rightarrow \mathbf{K}(\mathcal{E})$  is the natural functor, it can be completed to the following diagram:

$$\begin{array}{ccc}
 \mathcal{E}^\alpha & & \mathbf{K}(\mathcal{E}) \\
 \downarrow \gamma^{\mathcal{E}^\alpha} & \xrightarrow{K} & \downarrow Q_2 \\
 \mathbf{K}(\text{Mod}(\mathcal{E}^\alpha)) & \xrightarrow{Q_1} & \mathbf{K}(\mathcal{E}) \\
 \downarrow Q_3 & & \downarrow Q_2 \\
 \mathbf{D}(\text{Mod}(\mathcal{E}^\alpha)) & \xrightarrow{Q_4} & \check{\mathbf{D}}(\mathcal{E})
 \end{array}
 \tag{6.4}$$

It is easy to see that the diagram is commutative (up to isomorphism). Indeed, the lower triangle is commutative by definition, while the upper one, involving  $K$ , commutes by equation (6.3).

Let us now observe that the triangulated category  $\check{\mathbf{D}}(\mathcal{E})$  fits in the framework of Theorem 6.2. Indeed, by [23, Theorem 5.12], the functor  $Q_4$  is a localisation with localising subcategory  $\mathcal{S} \subseteq \mathbf{D}(\text{Mod}(\mathcal{E}^\alpha))$ . Thus  $Q_4$  induces an equivalence

$$F_1: \mathbf{D}(\text{Mod}(\mathcal{E}^\alpha))/\mathcal{S} \rightarrow \check{\mathbf{D}}(\mathcal{E}).$$

Moreover, by the second main result in [23], the triangulated category  $\check{\mathbf{D}}(\mathcal{E})$  (and thus the localisation  $\mathbf{D}(\text{Mod}(\mathcal{E}^\alpha))/\mathcal{S}$ ) is well-generated.

Therefore, to apply Theorem 6.2 and conclude that  $\check{\mathbf{D}}(\mathcal{E})$  has a unique enhancement, we just need to show that the quotient functor

$$\bar{Q}: \mathbf{D}(\text{Mod}(\mathcal{E}^\alpha)) \rightarrow \mathbf{D}(\text{Mod}(\mathcal{E}^\alpha))/\mathcal{S}$$

is right vanishing. The set  $\mathcal{R}$  of objects of  $\mathbf{D}(\text{Mod}(\mathcal{E}^\alpha))/\mathcal{S}$  with respect to which this is checked consists of the preimage under  $F_1$  of the objects that are isomorphic to complexes of injective objects in  $\mathcal{E}$  concentrated in positive degrees.

It is clear that (R1)–(R3) are satisfied by definition. To prove (R4), consider any  $G \in \mathcal{E}^\alpha$  and  $R \in \mathcal{R}$ . Then

$$\begin{aligned}
 \text{Hom}_{\mathbf{D}(\text{Mod}(\mathcal{E}^\alpha))/\mathcal{S}}(\bar{Q}(Y^{\mathcal{E}^\alpha}(G)), R) &\cong \text{Hom}_{\check{\mathbf{D}}(\mathcal{E})}(F_1(\bar{Q}(Y^{\mathcal{E}^\alpha}(G))), F_1(R)) \\
 &\cong \text{Hom}_{\check{\mathbf{D}}(\mathcal{E})}(Q_4(Y^{\mathcal{E}^\alpha}(G)), F_1(R)) \cong \text{Hom}_{\check{\mathbf{D}}(\mathcal{E})}(Q_2(K(G)), F_1(R)),
 \end{aligned}$$

where the second isomorphism is by the definition of  $F_1$ , and the third is due to the commutativity of equation (6.4). But since the functor  $Q_2$  consists of taking injective resolutions, the complex  $Q_2(K(G))$  has trivial cohomology in positive degrees. Hence

$$\text{Hom}_{\check{\mathbf{D}}(\mathcal{E})}(Q_2(K(G)), F_1(R)) = 0$$

by the definition of  $\mathcal{R}$ .

### 6.2. The completed derived category: basic properties

In this section, we discuss enhancements of the completed derived category. This is interesting to analyse, partly because this is a triangulated category whose definition starts from a dg category—which, of course, turns out to be one of its dg enhancements. In the presentation, we follow [29, Section 1.2.1]. We will not introduce the language of  $\infty$ -categories as we will only use it superficially in this paper.

Let  $\mathcal{E}$  be a Grothendieck category, and fix the enhancement of  $\mathbf{D}(\mathcal{E})$  given by the dg category  $\mathbf{D}_{\text{dg}}(\mathcal{E})$ , which is the Drinfeld quotient  $\mathbf{C}_{\text{dg}}(\mathcal{E})/\mathbf{Acy}(\mathcal{E})$ .



For  $i \in \mathbb{Z}$ , we denote by  $\mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq i}$  the full dg subcategory of  $\mathbf{D}_{\text{dg}}(\mathcal{G})$  such that  $H^0(\mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq i}) = \mathbf{D}(\mathcal{G})^{\geq i}$ . Once we interpret them as  $\infty$ -categories, one obtains functors

$$\tau^{\geq i} : \mathbf{D}_{\text{dg}}(\mathcal{G}) \longrightarrow \mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq i}.$$

From this, one produces a sequence of  $\infty$ -categories and functors

$$\dots \xrightarrow{\tau^{\geq -2}} \mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq -2} \xrightarrow{\tau^{\geq -1}} \mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq -1} \xrightarrow{\tau^{\geq 0}} \mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq 0} \xrightarrow{\tau^{\geq 1}} \mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq 1} \xrightarrow{\tau^{\geq 2}} \dots$$

Denote by  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})$  its homotopy limit.

By [29, Proposition 1.2.1.17],  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})$  is a stable  $\infty$ -category; and thus, by the main result in [15], it is naturally quasi-equivalent to a pretriangulated dg category. So, without loss of generality, we can assume in this paper that  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})$  is a pretriangulated dg category, thus motivating the notation.

**Remark 6.4.** By definition,  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})$  can alternatively be defined as the homotopy limit of the following sequence:

$$\dots \xrightarrow{\tau^{\geq -2}} \mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq -2} \xrightarrow{\tau^{\geq -1}} \mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq -1} \xrightarrow{\tau^{\geq 0}} \mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq 0}. \tag{6.5}$$

More precisely, one defines the left completion of  $\mathbf{D}^+(\mathcal{G})$  by using equation (6.5). By [29, Remark 1.2.1.18], such a completion is naturally quasi-equivalent to  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})$ . It follows that the objects of  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})$  can be identified with homotopy limits of objects in the full dg subcategories  $\mathbf{D}_{\text{dg}}(\mathcal{G})^{\geq i}$  for  $i \leq 0$ .

**Definition 6.5.** The (left) completed derived category of a Grothendieck category  $\mathcal{G}$  is the triangulated category  $\widehat{\mathbf{D}}(\mathcal{G}) := H^0(\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G}))$ .

By construction,  $\widehat{\mathbf{D}}(\mathcal{G})$  has a dg enhancement. The other properties of such a triangulated category that are relevant in this paper are summarised in the following proposition (see Section 5.5 for the definition and basic properties of t-structures).

**Proposition 6.6** ([29], Proposition 1.2.1.17). *Let  $\mathcal{G}$  be a Grothendieck category.*

- (1) *There is a natural exact functor  $\mathbf{D}(\mathcal{G}) \rightarrow \widehat{\mathbf{D}}(\mathcal{G})$ .*
- (2) *The category  $\widehat{\mathbf{D}}(\mathcal{G})$  has a natural t-structure, and the functor in (1) identifies its heart with  $\mathcal{G}$  and induces an equivalence  $\mathbf{D}(\mathcal{G})^{\geq 0} \cong \widehat{\mathbf{D}}(\mathcal{G})^{\geq 0}$ .*
- (3) *The category  $\widehat{\mathbf{D}}(\mathcal{G})$  is left complete (i.e., for any  $X \in \widehat{\mathbf{D}}(\mathcal{G})$ , the natural morphism  $X \rightarrow \text{Holim } X^{\geq i}$  is an isomorphism).*

Part (3) in the proposition above, which essentially follows from the definition, motivates the choice of the name for  $\widehat{\mathbf{D}}(\mathcal{G})$ . As a consequence of (2), we have a natural equivalence

$$\widehat{\mathbf{D}}(\mathcal{G})^+ \cong \mathbf{D}^+(\mathcal{G}), \tag{6.6}$$

where  $\widehat{\mathbf{D}}(\mathcal{G})^+$  denotes the full subcategory of  $\widehat{\mathbf{D}}(\mathcal{G})$  consisting of all the objects  $X \in \text{Ob}(\widehat{\mathbf{D}}(\mathcal{G}))$  such that  $X = X^{\geq i}$  for  $i \ll 0$ . We denote by  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})^+$  the full dg subcategory of  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})$  such that  $H^0(\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{G})^+) = \widehat{\mathbf{D}}(\mathcal{G})^+$ .

**Remark 6.7.** It was proved in [32] that there exist Grothendieck categories  $\mathcal{G}$  such that  $\mathbf{D}(\mathcal{G})$  is not left complete in the sense of Proposition 6.6 (3). The category  $\widehat{\mathbf{D}}(\mathcal{G})$  provides a natural way to complete  $\mathbf{D}(\mathcal{G})$  in view of Proposition 6.6 (1).

### 6.3. The completed derived category: uniqueness of enhancements

Let us prove that  $\widehat{\mathbf{D}}(\mathcal{E})$  has a unique dg enhancement when  $\mathcal{E}$  is a Grothendieck category. To do this, we use Remark 3.19 (ii) and show that  $\widehat{\mathbf{D}}(\mathcal{E})^\circ$  has a unique dg enhancement. Hence we assume that there is an exact equivalence

$$F: \widehat{\mathbf{D}}(\mathcal{E})^\circ \xrightarrow{\sim} H^0(\mathcal{E}),$$

where  $\mathcal{E}$  is a pretriangulated dg category.

**Remark 6.8.** By Proposition 6.6 (2),  $\widehat{\mathbf{D}}(\mathcal{E})$  has a natural t-structure. Hence  $\widehat{\mathbf{D}}(\mathcal{E})^\circ$  and  $H^0(\mathcal{E})$  are endowed with a t-structure  $(H^0(\mathcal{E})^{\geq 0}, H^0(\mathcal{E})^{\leq 0})$  as well. Of course,  $F$  yields equivalences  $H^0(\mathcal{E})^{\geq 0} \cong \widehat{\mathbf{D}}(\mathcal{E})^{\leq 0}$  and  $H^0(\mathcal{E})^{\leq 0} \cong \widehat{\mathbf{D}}(\mathcal{E})^{\geq 0}$ . Moreover, as by Proposition 6.6 (3)  $\widehat{\mathbf{D}}(\mathcal{E})$  is left complete, by passing to the opposite category, we have that the natural morphism

$$\text{Hocolim } X^{\leq i} \longrightarrow X$$

is an isomorphism for any  $X \in \text{Ob}(H^0(\mathcal{E}))$ . More generally, in view of Remark 6.4, the objects of  $\widehat{\mathbf{D}}(\mathcal{E})^\circ$  (respectively,  $H^0(\mathcal{E})$ ) are identified with all the homotopy colimits of sequences of objects in  $(\widehat{\mathbf{D}}(\mathcal{E})^\circ)^{\leq i}$  (respectively,  $H^0(\mathcal{E})^{\leq i}$ ) with eventually stable cohomology.

Clearly, there exists a full pretriangulated dg subcategory  $\mathcal{E}^+ \subseteq \mathcal{E}$  such that  $F^+ := F|_{(\widehat{\mathbf{D}}(\mathcal{E}^+)^\circ)}$  is an exact equivalence

$$F^+: (\widehat{\mathbf{D}}(\mathcal{E}^+)^\circ) \xrightarrow{\sim} H^0(\mathcal{E}^+).$$

In the following, let  $\mathcal{D}$  be either  $\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^\circ$  or  $\mathcal{E}$ , and let  $\mathcal{D}^+$  be either  $(\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E}^+)^\circ)$  or  $\mathcal{E}^+$ . We denote by  $\widehat{Y}_{\text{dg}}^{\mathcal{D}}: \mathcal{D} \rightarrow \text{h-proj}(\mathcal{D}^+)$  the morphism in **Hqe** defined as the composition

$$\mathcal{D} \xrightarrow{Y_{\text{dg}}^{\mathcal{D}}} \text{dgMod}(\mathcal{D}) \xrightarrow{\text{Res}(I)} \text{dgMod}(\mathcal{D}^+) \xrightarrow{P} \text{dgMod}(\mathcal{D}^+)/\text{dgAcy}(\mathcal{D}^+) \xrightarrow{(P \circ J)^{-1}} \text{h-proj}(\mathcal{D}^+),$$

where  $I: \mathcal{D}^+ \hookrightarrow \mathcal{D}$  and  $J: \text{h-proj}(\mathcal{D}^+) \hookrightarrow \text{dgMod}(\mathcal{D}^+)$  are the natural inclusions and  $P$  is the quotient dg functor. Moreover, let  $\mathcal{D}^{\leq i}$  be the full dg subcategory of  $\mathcal{D}$  such that  $H^0(\mathcal{D}^{\leq i}) = H^0(\mathcal{D})^{\leq i}$  (we implicitly refer to the t-structures in Remark 6.8).

**Lemma 6.9.** *The essential image of the exact functor  $H^0(\widehat{Y}_{\text{dg}}^{\mathcal{D}})$  consists of objects isomorphic to homotopy colimits in  $H^0(\text{h-proj}(\mathcal{D}^+))$  of  $Y_{\text{dg}}^{\mathcal{D}^+}(X_i)$ , where  $X_i \in \text{Ob}(\mathcal{D}^{\leq i})$  for  $i \geq 0$ . Moreover,  $\widehat{Y}_{\text{dg}}^{\mathcal{D}}$  is quasi-fully faithful.*

*Proof.* Let  $X \in \text{Ob}(\mathcal{D})$ . By Remark 6.8, we have a (closed, degree 0) morphism in  $\text{dgMod}(\mathcal{D}^+)$

$$\varphi_X: \text{Hocolim } \text{Hom}_{\mathcal{D}}(-, X^{\leq i}) \longrightarrow \text{Hom}_{\mathcal{D}}(-, X),$$

which we claim to be a quasi-isomorphism. To prove this, it is enough to show that the induced morphism  $\widetilde{\varphi}_X(Y): \text{Hocolim } \text{Hom}_{H^0(\mathcal{D})}(Y, X^{\leq i}) \rightarrow \text{Hom}_{H^0(\mathcal{D})}(Y, X)$  is an isomorphism for all  $Y \in \text{Ob}(H^0(\mathcal{D}^+))$ . But in the t-structure on  $H^0(\mathcal{D})$  from Remark 6.8,  $Y$  has bounded above nontrivial cohomologies. That means the natural maps

$$\text{Hom}_{H^0(\mathcal{D})}(-, X^{\leq i}) \longrightarrow \text{Hom}_{H^0(\mathcal{D})}(-, X^{\leq i+1})$$

are isomorphisms for  $i$  sufficiently large. Hence we have natural isomorphisms

$$\text{Hocolim } \text{Hom}_{H^0(\mathcal{D})}(Y, X^{\leq i}) \cong \text{Hom}_{H^0(\mathcal{D})}(Y, \text{Hocolim } X^{\leq i}) \cong \text{Hom}_{H^0(\mathcal{D})}(Y, X)$$

and the composition of these maps is precisely  $\widehat{\varphi}_X(Y)$ . This shows that any object in the essential image of  $H^0(\widehat{Y}_{\text{dg}}^{\mathcal{D}})$  is isomorphic to the homotopy colimit of objects in  $\mathcal{D}^{\leq i}$  for  $i \geq 0$ . To show the other inclusion, note that the same argument proves that the dg module  $\text{Hocolim Hom}_{\mathcal{D}}(-, X_i)$ , where  $X_i \in \text{Ob}(\mathcal{D}^{\leq i})$ , is quasi-isomorphic to  $\text{Hom}_{\mathcal{D}}(-, X)$ , where  $X := \text{Hocolim } X_i$  is in  $\mathcal{D}$  by Remark 6.8.

Let us now prove that  $\widehat{Y}_{\text{dg}}^{\mathcal{D}}$  is quasi-fully faithful. For  $X_1, X_2 \in \text{Ob}(H^0(\mathcal{D}))$ , we have the quasi-isomorphisms

$$\begin{aligned} \text{Hom}_{\text{h-proj}(\mathcal{D}^+) } \left( \widehat{Y}_{\text{dg}}^{\mathcal{D}}(X_1), \widehat{Y}_{\text{dg}}^{\mathcal{D}}(X_2) \right) &\cong \text{Hom}_{\text{h-proj}(\mathcal{D}^+) } \left( \text{Hocolim Hom}_{\mathcal{D}}(-, X_1^{\leq i}), \widehat{Y}_{\text{dg}}^{\mathcal{D}}(X_2) \right) \\ &\cong \text{Holim Hom}_{\text{h-proj}(\mathcal{D}^+) } \left( \text{Hom}_{\mathcal{D}}(-, X_1^{\leq i}), \widehat{Y}_{\text{dg}}^{\mathcal{D}}(X_2) \right) \cong \text{Holim Hom}_{\mathcal{D}}(X_1^{\leq i}, X_2) \\ &\cong \text{Hom}_{\mathcal{D}}(\text{Hocolim } X_1^{\leq i}, X_2) \cong \text{Hom}_{\mathcal{D}}(X_1, X_2), \end{aligned}$$

where the first quasi-isomorphism is a consequence of the fact that  $\varphi_{X_1}$  is a quasi-isomorphism. The third one follows from the Yoneda lemma, while the last one is again Remark 6.8. For the second and the fourth quasi-isomorphisms, we use the fact that the Hom-functor turns colimits in the first argument into limits. Therefore,  $\widehat{Y}_{\text{dg}}^{\mathcal{D}}$  is quasi-fully faithful.  $\square$

By equation (6.6), Theorem A (1) and again Remark 3.19 (ii), there is a morphism

$$g^+ \in \text{Hom}_{\mathbf{Hqe}} \left( (\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^+)^{\circ}, \mathcal{E}^+ \right)$$

such that

- (a)  $G^+ := H^0(g^+)$  is an exact equivalence.
- (b)  $G^+(X) \cong F^+(X)$  for all  $X \in \text{Ob}((\widehat{\mathbf{D}}(\mathcal{E})^+)^{\circ})$  (see Remark 5.4).

The morphism  $g := \text{Ind}(g^+) \in \text{Hom}_{\mathbf{Hqe}}(\text{h-proj}((\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^+)^{\circ}), \text{h-proj}(\mathcal{E}^+))$  is such that  $G := H^0(g)$  is an exact equivalence by (a). Thus we get the diagram

$$\begin{array}{ccccc} \widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^{\circ} & \xleftarrow{\quad} & (\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^+)^{\circ} & \xrightarrow{g^+} & \mathcal{E}^+ \hookrightarrow \mathcal{E} \\ & \searrow \widehat{Y}_{\text{dg}}^{\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^{\circ}} & \downarrow Y_{\text{dg}}^{(\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^+)^{\circ}} & & \downarrow Y_{\text{dg}}^{\mathcal{E}^+} \\ & & \text{h-proj}((\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^+)^{\circ}) & \xrightarrow{g} & \text{h-proj}(\mathcal{E}^+) \end{array} \tag{6.7}$$

which is commutative in  $\mathbf{Hqe}$ . Denote by  $g'$  the composition  $g \circ \widehat{Y}_{\text{dg}}^{\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^{\circ}}$ . It is clear from Lemma 6.9 that  $H^0(g')$  is fully faithful.

Let us prove that the essential image of  $H^0(g')$  coincides with the essential image of  $H^0(\widehat{Y}_{\text{dg}}^{\mathcal{E}})$ . Let  $X \in \text{Ob}(\widehat{\mathbf{D}}(\mathcal{E}))$ . Then by Remark 6.8, equation (6.7) and the fact that  $G$  is an exact equivalence, we get

$$\begin{aligned} H^0(g')(X) &\cong G \left( H^0 \left( \widehat{Y}_{\text{dg}}^{(\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^+)^{\circ}} \right) (\text{Hocolim } X^{\leq i}) \right) \cong G \left( \text{Hocolim } H^0 \left( \widehat{Y}_{\text{dg}}^{(\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^+)^{\circ}} \right) (X^{\leq i}) \right) \\ &\cong \text{Hocolim } G \left( H^0 \left( \widehat{Y}_{\text{dg}}^{(\widehat{\mathbf{D}}_{\text{dg}}(\mathcal{E})^+)^{\circ}} \right) (X^{\leq i}) \right) \cong \text{Hocolim } H^0 \left( Y_{\text{dg}}^{\mathcal{E}^+} \right) (G^+(X^{\leq i})). \end{aligned}$$

By (b),  $G^+(X^{\leq i}) \cong F^+(X^{\leq i}) \in \text{Ob}(\mathcal{E}^{\leq i})$  for all  $i \geq 0$ . By Lemma 6.9, we get that  $H^0(g')(X)$  is in the essential image of  $H^0(\widehat{Y}_{\text{dg}}^{\mathcal{E}})$ . The same argument shows the other inclusion.

In conclusion, let  $g''$  denote the inverse of  $\widehat{Y}_{\text{dg}}^{\mathcal{E}}$  in  $\mathbf{Hqe}$ , when we interpret  $\widehat{Y}_{\text{dg}}^{\mathcal{E}}$  as a morphism between  $\mathcal{E}$  and its essential image. The above argument shows that  $g'' \circ g'$  is an isomorphism in  $\text{Hom}_{\mathbf{Hqe}}(\widehat{\mathbf{D}}(\mathcal{E})^{\circ}, \mathcal{E})$  as we want.

**6.4. Two applications: a shortcut and a negative result**

Let us first note that the methods used in the proof in Section 6.1 (or rather, their specialisation to the subcategory of compact objects) can be used to give a straightforward proof of the uniqueness of dg enhancements of  $\mathbf{D}^b(\mathcal{A})$ , where  $\mathcal{A}$  is a small abelian category.

Indeed, given such an  $\mathcal{A}$ , one takes the Grothendieck category  $\mathcal{G} := \text{Ind}(\mathcal{A})$  (see Example 2.3 (iii)). In that case, by Example 2.7 and [23, Proposition 4.7], we have a sequence of equivalences

$$\mathbf{D}^b(\mathcal{A}) \cong \mathbf{K}(\text{Inj}(\text{Ind}(\mathcal{A})))^c \cong (\mathbf{K}(\text{Inj}(\mathcal{A}))^c / \mathcal{L}')^{ic}, \tag{6.8}$$

where, by abuse of notation, we write  $\text{Inj}(\mathcal{A})$  instead of  $\text{Inj}(\text{Mod}(\mathcal{A}))$  and where  $\mathcal{L}' \subseteq \mathbf{K}(\text{Inj}(\mathcal{A}))^c$  is a full triangulated subcategory. Let  $\mathcal{L}$  be the smallest localising subcategory of  $\mathbf{K}(\text{Inj}(\mathcal{A}))$  containing  $\mathcal{L}'$ . By the main result in [34], we deduce from this the exact equivalence

$$\mathbf{D}^b(\mathcal{A}) \cong (\mathbf{K}(\text{Inj}(\mathcal{A})) / \mathcal{L})^c. \tag{6.9}$$

Since  $\mathbf{K}(\text{Inj}(\mathcal{A})) \cong \mathbf{D}(\text{Mod}(\mathcal{A})) \cong \mathcal{D}(\mathcal{A})$  (see, for example, [23, Lemma 4.8]), the uniqueness of dg enhancements for  $\mathbf{D}^b(\mathcal{A})$  is a consequence of the following result.

**Theorem 6.10** ([27], Theorem 2). *Let  $\mathcal{A}$  be a small category, and let  $\mathcal{L}$  be a localising subcategory of  $\mathcal{D}(\mathcal{A})$  such that:*

- (a)  $\mathcal{L}^c = \mathcal{L} \cap \mathcal{D}(\mathcal{A})^c$  and  $\mathcal{L}^c$  satisfies (G1) in  $\mathcal{L}$ .
- (b)  $\text{Hom}_{\mathcal{D}(\mathcal{A})/\mathcal{L}}(\mathbf{Q}(Y^{\mathcal{A}}(A_1)), \mathbf{Q}(Y^{\mathcal{A}}(A_2))[i]) = 0$  for all  $A_1, A_2 \in \mathcal{A}$  and all integers  $i < 0$ .

Then  $(\mathcal{D}(\mathcal{A})/\mathcal{L})^c$  has a unique enhancement.

Indeed, in our situation,  $\mathcal{L}^c = \mathcal{L}'$  and assumption (b) is clear because, given  $A \in \mathcal{A}$ , the object  $Y^{\mathcal{A}}(A) \in \mathcal{D}(\mathcal{A})$  is mapped to  $A$  by the composition of the equivalences given by equations (6.8) and (6.9) (see the discussion in [13, Section 6.1]).

Let us now pass to the negative examples of this section. The relation between the uniqueness of dg enhancements of triangulated categories and their admissible subcategories turns out to be a delicate and intriguing problem. In [9], the following natural question was raised:

**Question 6.11.** Let  $\mathcal{S}$  be a full triangulated subcategory of a triangulated category  $\mathcal{T}$ . Does  $\mathcal{S}$  have unique enhancement if  $\mathcal{T}$  does? What if  $\mathcal{S}$  is a localising (if  $\mathcal{T}$  has small coproducts) or an admissible subcategory of  $\mathcal{T}$ ?

Recall that a full triangulated subcategory  $\mathcal{S}$  of a triangulated category  $\mathcal{T}$  is *admissible* if the inclusion functor  $\mathcal{S} \hookrightarrow \mathcal{T}$  has left and right adjoints.

As an application of Theorem A (2), we deduce the following result.

**Corollary 6.12.** *Assume that  $\mathbb{k} = \mathbb{Z}$ . Then there exists a triangulated category  $\mathcal{T}$  with small coproducts and a localising and admissible subcategory  $\mathcal{S}$  of  $\mathcal{T}$  such that  $\mathcal{T}$  has a unique enhancement, while  $\mathcal{S}$  does not.*

*Proof.* For  $i = 1, 2$ , consider the triangulated categories  $\mathcal{T}_i$  with localising and admissible subcategories  $\mathcal{S}_i \hookrightarrow \mathcal{T}_i$  in Example 1.4. For the convenience of the reader, let us recall that in the example, we consider rings  $R_1 := \mathbb{Z}/p^2\mathbb{Z}$  and  $R_2 := \mathbb{F}_p[\varepsilon]$  (where  $\varepsilon^2 = 0$ ) and, for  $i = 1, 2$ , we set  $\mathcal{T}_i := \mathbf{D}(\text{Mod}(R_i))$ . Moreover, we denoted by  $\mathcal{S}_i$  the full subcategory of  $\mathcal{T}_i$  consisting of acyclic complexes. By Theorem A,  $\mathcal{T}_i$  has a unique dg enhancement, whereas  $\mathcal{S}_i$  does not, as it is explained in [9, Section 3.3]. Indeed, while  $\mathcal{S}_1$  and  $\mathcal{S}_2$  are both equivalent to the category  $\text{Mod}(\mathbb{F}_p)$  endowed with the triangulated structure defined by  $[1] = \text{id}$  (and, necessarily, distinguished triangles given by triangles inducing long exact sequences), it follows from [17, 42] that the natural enhancements of  $\mathcal{S}_1$  and  $\mathcal{S}_2$  are not isomorphic in  $\mathbf{Hqe}$ .  $\square$

**Remark 6.13.** (i) Keeping the notation of Example 1.4, the triangulated categories  $\mathcal{S}_1$  and  $\mathcal{S}_2$  are  $\mathbb{F}_p$ -linear, and the same is true for the natural enhancement of  $\mathcal{S}_2$ , but not for that of  $\mathcal{S}_1$ . Thus it remains an open problem whether a similar counterexample can be found when  $\mathbb{k}$  is a field.

(ii) In Example 1.4, the fact that  $\mathbf{D}(\text{Mod}(R_i))$  is compactly generated implies that the right adjoint to the quotient functor  $\mathbf{K}(\text{Inj}(\text{Mod}(R_i))) \rightarrow \mathbf{D}(\text{Mod}(R_i))$  has itself a right adjoint. Hence,  $\mathbf{D}(\text{Mod}(R_i))$  is an admissible localising subcategory of  $\mathbf{K}(\text{Inj}(\text{Mod}(R_i)))$  and, moreover, both triangulated categories  $\mathbf{K}(\text{Inj}(\text{Mod}(R_i)))$  and  $\mathbf{D}(\text{Mod}(R_i))$  have unique enhancements (by Theorem A). This, together with Corollary 6.12, shows that the same triangulated category  $\mathbf{K}(\text{Inj}(\text{Mod}(R_i)))$  may have a unique enhancement and, at the same time, contain admissible (localising) subcategories answering both positively (e.g.,  $\mathbf{D}(\text{Mod}(R_i))$ ) and negatively (e.g.,  $\mathcal{S}_i$ ) to Question 6.11. This clarifies that a complete answer to the question above cannot simply rely on the properties of the ambient category.

### 7. Homotopy pullbacks and enhancements

In this part, we investigate the relation of dg enhancement with homotopy pullbacks; this will provide us with the formal machinery needed to prove Theorem B. The framework we will be working with is set up in some generality in Section 7.1, with the examples hinting at the geometric relevance of the formalism. In Section 7.2, we state and prove the main ingredient in the proof of Theorem B, while Section 7.3 provides technical refinements.

#### 7.1. The setting

Let us summarise the abstract setting, where we aim to state and prove the general criterion for the uniqueness of dg enhancements in the presence of homotopy pullbacks.

**Setup 7.1.** Let  $\mathcal{C}$  be a pretriangulated dg category such that  $H^0(\mathcal{C})$  is idempotent complete, and let  $\mathcal{D}_i \subseteq \mathcal{C}$  be a full dg subcategory for  $i = 1, 2$  such that

- (i)  $\mathcal{D}_i$  is closed under shifts for  $i = 1, 2$ .
- (ii)  $\text{Hom}_{H^0(\mathcal{C})}(H^0(\mathcal{D}_i), H^0(\mathcal{D}_j)) = 0$  for  $i \neq j \in \{1, 2\}$ .

Assume further that there is a commutative diagram in **dgCat**

$$\begin{array}{ccc}
 \mathcal{C} & \xrightarrow{Q_1} & \mathcal{C}_{\mathcal{D}_1} \\
 Q_2 \downarrow & & \downarrow \bar{Q}_1 \\
 \mathcal{C}_{\mathcal{D}_2} & \xrightarrow{\bar{Q}_2} & \mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}
 \end{array}$$

of pretriangulated and idempotent complete dg categories and such that

- (iii)  $H^0(Q_i)$  is the idempotent completion of the Verdier quotient functor that sends to zero the thick subcategory  $\mathcal{D}_i \subseteq H^0(\mathcal{C})$  generated by  $H^0(\mathcal{D}_i)$  for  $i = 1, 2$ .
- (iv)  $H^0(\bar{Q}_1) \circ H^0(Q_1) (= H^0(\bar{Q}_2) \circ H^0(Q_2))$  is the idempotent completion of the Verdier quotient functor that sends to zero the thick subcategory  $\mathcal{D}_{1,2} \subseteq H^0(\mathcal{C})$  generated by  $H^0(\mathcal{D}_1) \cup H^0(\mathcal{D}_2)$ .

For the non-expert reader, let us recall that given an exact functor  $F: \mathcal{T}_1 \rightarrow \mathcal{T}_2$  and a thick subcategory  $\mathcal{S} \subseteq \mathcal{T}_1$ , we say that  $F$  is the idempotent completion of the Verdier quotient that sends  $\mathcal{S}$  to zero if we have a factorisation

$$\begin{array}{ccc}
 \mathcal{T}_1 & \xrightarrow{F} & \mathcal{T}_2 \\
 \searrow Q & & \uparrow I \\
 & & \mathcal{T}_1/\mathcal{S}
 \end{array}$$

where  $Q$  is the Verdier quotient functor and  $I$  is the idempotent completion of  $\mathcal{T}_1/\mathcal{S}$ .

Let us now discuss the two geometric examples that are of interest here.

**Example 7.2** (Quasi-coherent sheaves). Let  $X$  be a quasi-compact and quasi-separated scheme. Let  $\mathbf{D}_{\text{qc}}^?(X)$  be the full triangulated subcategory of  $\mathbf{D}^2(\text{Mod}(\mathcal{O}_X))$  consisting of complexes with quasi-coherent cohomologies for  $? = b, +, -, \emptyset$ . Let  $U_1, U_2 \subseteq X$  be quasi-compact open subsets such that  $X = U_1 \cup U_2$ . Denote by  $\iota_i: U_i \hookrightarrow X$  and  $\iota_{1,2}: U_1 \cap U_2 \hookrightarrow X$  the open immersions.

Let  $(\mathcal{C}, F)$  be a dg enhancement of  $\mathbf{D}_{\text{qc}}^?(X)$ , and assume that  $\mathcal{C}$  is h-flat. Set

$$\begin{aligned} \mathcal{D}_i &:= \{C \in \mathcal{C} : \iota_i^* \circ F(C) \cong 0\}, \\ \mathcal{D}_{1,2} &:= \{C \in \mathcal{C} : \iota_{1,2}^* \circ F(C) \cong 0\}, \end{aligned}$$

for  $i = 1, 2$ . Assumption (i) in Setup 7.1 is then verified. Moreover,  $\overline{\mathcal{D}}_i = H^0(\mathcal{D}_i) \cong \mathbf{D}_{X \setminus U_i}^?(X)$  and  $\overline{\mathcal{D}}_{1,2} = H^0(\mathcal{D}_{1,2}) \cong \mathbf{D}_{X \setminus (U_1 \cap U_2)}^?(X)$ , where  $\mathbf{D}_Z^?(X)$  denotes the full (triangulated) subcategory of  $\mathbf{D}_{\text{qc}}^?(X)$  consisting of complexes with cohomology supported on the closed subset  $Z \subseteq X$ . Given this identification, the fact that  $X = U_1 \cup U_2$  clearly implies that  $\text{Hom}_{H^0(\mathcal{C})}(H^0(\mathcal{D}_i), H^0(\mathcal{D}_j)) = 0$ , when  $i \neq j \in \{1, 2\}$ . Thus (ii) in Setup 7.1 holds true.

Consider the (idempotent completion of the) Drinfeld quotients  $\mathcal{C}_{\mathcal{D}_i} := \mathcal{C}/\mathcal{D}_i$  and  $\mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2} := \mathcal{C}/\mathcal{D}_{1,2}$ . Following the discussion in Section 3.2, we get dg quotient functors  $Q_i: \mathcal{C} \rightarrow \mathcal{C}_{\mathcal{D}_i}$  and  $Q_{1,2}: \mathcal{C} \rightarrow \mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}$ . Moreover,  $\mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}$  is, at the same time, the Drinfeld quotient of  $\mathcal{C}_{\mathcal{D}_1}$  and of  $\mathcal{C}_{\mathcal{D}_2}$  by  $Q_1(\mathcal{D}_2)$  and  $Q_2(\mathcal{D}_1)$ , respectively. Hence we also get the dg functors  $\overline{Q}_i: \mathcal{C}_{\mathcal{D}_i} \rightarrow \mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}$  for  $i = 1, 2$ . Since  $\mathcal{C}$  is h-flat, the results in Section 3.2 show that

$$\begin{aligned} H^0(\mathcal{C}_{\mathcal{D}_i}) &\cong H^0(\mathcal{C})/H^0(\mathcal{D}_i) \cong \mathbf{D}_{\text{qc}}^?(U_i), \\ H^0(\mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}) &\cong H^0(\mathcal{C})/H^0(\mathcal{D}_{1,2}) \cong \mathbf{D}_{\text{qc}}^?(U_1 \cap U_2), \end{aligned}$$

for  $i = 1, 2$ . Moreover,  $H^0(Q_i)$  and  $H^0(\overline{Q}_1) \circ H^0(Q_1)$  are the corresponding Verdier quotient functors. Hence assumptions (iii) and (iv) in Setup 7.1 are satisfied as well.

**Example 7.3** (Perfect complexes). As in the previous example, let  $X$  be a quasi-compact and quasi-separated scheme. Let  $\mathbf{Perf}(X)$  be the category of perfect complexes on  $X$  (i.e., complexes in  $\mathbf{D}_{\text{qc}}(X)$ , which are locally quasi-isomorphic to bounded complexes of locally free sheaves of finite type). By [7, Theorem 3.1.1],  $\mathbf{Perf}(X)$  can alternatively be described as the category of compact objects  $\mathbf{D}_{\text{qc}}(X)^c$ .

If  $(\mathcal{C}, F)$  is a dg enhancement of  $\mathbf{Perf}(X)$  with  $\mathcal{C}$  h-flat, then the same construction as in Example 7.2 yields the dg categories  $\mathcal{D}_i, \mathcal{D}_{1,2}, \mathcal{C}_{\mathcal{D}_i}$  and  $\mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}$  with dg functors  $Q_i$  and  $\overline{Q}_i$ . The fact that the assumptions of Setup 7.1 are satisfied can be checked as in Example 7.2. One has to be careful only about (iii) and (iv) as for localisation theory for compact objects in [34, 45], the functor  $H^0(Q_i): H^0(\mathcal{C}) \rightarrow H^0(\mathcal{C}_{\mathcal{D}_i}) \cong \mathbf{Perf}(U_i)$  is the composition

$$H^0(\mathcal{C}) \longrightarrow H^0(\mathcal{C})/H^0(\mathcal{D}_i) \hookrightarrow H^0(\mathcal{C}_{\mathcal{D}_i}),$$

where the first functor is the Verdier quotient functor while the latter inclusion identifies  $H^0(\mathcal{C}_{\mathcal{D}_i})$  to the idempotent completion of  $H^0(\mathcal{C})/H^0(\mathcal{D}_i)$ . The same is true for the composition  $H^0(\overline{Q}_1) \circ H^0(Q_1)$ .

### 7.2. A useful criterion

This section is devoted to the proof of the following result, which will be crucial for our geometric applications.

**Theorem 7.4.** In Setup 7.1, the natural dg functor from  $\mathcal{C}$  to the homotopy pullback  $\mathcal{C}_{\mathcal{D}_1} \times_{\mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}}^h \mathcal{C}_{\mathcal{D}_2}$  of the diagram

$$\mathcal{C}_{\mathcal{D}_1} \xrightarrow{\bar{Q}_1} \mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2} \xleftarrow{\bar{Q}_2} \mathcal{C}_{\mathcal{D}_2}$$

is a quasi-equivalence.

*Proof.* To simplify the notation, we set  $\mathcal{P} := \mathcal{C}_{\mathcal{D}_1} \times_{\mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}}^h \mathcal{C}_{\mathcal{D}_2}$ , and we use its explicit construction discussed in Section 3.3. We denote by  $F: \mathcal{C} \rightarrow \mathcal{P}$  the natural dg functor explicitly described in Remark 3.12. We want to prove that it is a quasi-equivalence.

Let us begin by proving that the functor  $F$  is quasi-fully-faithful. Since the assumptions of Proposition 3.14 are satisfied, any object  $C \in \text{Ob}(\mathcal{C})$  sits in a commutative diagram

$$\begin{array}{ccccc} & & D_2 & \xlongequal{\quad} & D_2 \\ & & \downarrow & & \downarrow \\ D_1 & \longrightarrow & C & \longrightarrow & C_{D_1} \\ \parallel & & \downarrow & & \downarrow \\ D_1 & \longrightarrow & C_{D_2} & \longrightarrow & C_{D_1, D_2} \end{array}$$

where rows and columns yield distinguished triangles in  $H^0(\text{h-proj}(\mathcal{C}))$ ,  $D_i \in \text{Ob}(\text{h-proj}(\mathcal{D}_i))$  and the complexes

$$\text{Hom}_{\text{h-proj}(\mathcal{C})}(\text{h-proj}(\mathcal{D}_i), C_{D_i}) \quad \text{Hom}_{\text{h-proj}(\mathcal{C})}(\text{h-proj}(\mathcal{D}_i), C_{D_1, D_2}) \tag{7.1}$$

are acyclic for  $i = 1, 2$ . It follows first that the square

$$\begin{array}{ccc} \text{Hom}_{\text{h-proj}(\mathcal{C})}(C', C) & \longrightarrow & \text{Hom}_{\text{h-proj}(\mathcal{C})}(C', C_{D_1}) \\ \downarrow & & \downarrow \\ \text{Hom}_{\text{h-proj}(\mathcal{C})}(C', C_{D_2}) & \longrightarrow & \text{Hom}_{\text{h-proj}(\mathcal{C})}(C', C_{D_1, D_2}) \end{array} \tag{7.2}$$

is a homotopy cartesian square in  $\text{dgMod}(\mathbb{k})$  (here we use the same terminology as in [36, Section 1.4]) for all  $C' \in \text{Ob}(\mathcal{C})$ . Now equation (7.1) and assumptions (iii) and (iv) in Setup 7.1 imply that the natural maps

$$\begin{aligned} \text{Hom}_{\text{h-proj}(\mathcal{C})}(C', C_{D_i}) &\longrightarrow \text{Hom}_{\text{h-proj}(\mathcal{D}_i)}(\text{Ind}(Q_i)(C'), \text{Ind}(Q_i)(C_{D_i})) \\ \text{Hom}_{\text{h-proj}(\mathcal{C})}(C', C_{D_1, D_2}) &\longrightarrow \text{Hom}_{\text{h-proj}(\mathcal{D}_1, \mathcal{D}_2)}(\text{Ind}(\bar{Q}_1 \circ Q_1)(C'), \text{Ind}(\bar{Q}_1 \circ Q_1)(C_{D_1, D_2})) \end{aligned}$$

induced, respectively, by  $\text{Ind}(Q_i)$  for  $i = 1, 2$  and by  $\text{Ind}(\bar{Q}_1 \circ Q_1)(= \text{Ind}(\bar{Q}_2 \circ Q_2))$  are all quasi-isomorphisms in  $\text{dgMod}(\mathbb{k})$ . On the other hand, the images under  $\text{Ind}(Q_i)$  and  $\text{Ind}(\bar{Q}_1 \circ Q_1)$  of the natural morphisms  $C \rightarrow C_{D_i}$  and  $C \rightarrow C_{D_1, D_2}$  in  $\text{h-proj}(\mathcal{C})$  become, by construction, quasi-isomorphisms in  $\text{h-proj}(\mathcal{D}_i)$  and  $\text{h-proj}(\mathcal{D}_1, \mathcal{D}_2)$ , respectively.

Hence equation (7.2) is quasi-isomorphic in  $\text{dgMod}(\mathbb{k})$  to the square

$$\begin{CD} \text{Hom}_{\mathcal{E}}(C', C) @>>> \text{Hom}_{\mathcal{E}_{\mathcal{D}_1}}(C', C) \\ @VVV @VVV \\ \text{Hom}_{\mathcal{E}_{\mathcal{D}_2}}(C', C) @>>> \text{Hom}_{\mathcal{E}_{\mathcal{D}_1, \mathcal{D}_2}}(C', C) \end{CD} \tag{7.3}$$

where the arrows are induced by the quotient dg functors  $Q_i$  and  $\bar{Q}_i$  for  $i = 1, 2$ .

In conclusion, equation (7.3) must be homotopy cartesian in  $\text{dgMod}(\mathbb{k})$ , and this implies that the dg functor  $F$  induces a quasi-isomorphism

$$\text{Hom}_{\mathcal{E}}(C', C) \longrightarrow \text{Hom}_{\mathcal{P}}(F(C'), F(C)),$$

for all  $C', C \in \text{Ob}(\mathcal{E})$ , meaning  $F$  is quasi-fully faithful.

It remains to show that  $H^0(F)$  is essentially surjective. To this end, let us recall that an object in  $\mathcal{P}$  is a triple  $(C_1, C_2, f)$ , where  $C_i \in \text{Ob}(\mathcal{E}_{\mathcal{D}_i})$  and  $f: \bar{Q}_1(C_1) \rightarrow \bar{Q}_2(C_2)$  is a homotopy equivalence in  $\mathcal{E}_{\mathcal{D}_1, \mathcal{D}_2}$ . Hence, we need to show that given such a triple  $(C_1, C_2, f)$ , there is  $C \in \text{Ob}(\mathcal{E})$  and a homotopy equivalence  $F(C) \cong (C_1, C_2, f)$  in  $\mathcal{P}$ .

First, we note that by assumption (iii) in Setup 7.1, we may choose  $\tilde{C}_i \in \text{Ob}(\mathcal{E})$  such that  $C_i$  is a direct factor  $Q_i(\tilde{C}_i)$  in  $H^0(\mathcal{E}_{\mathcal{D}_i})$  for  $i = 1, 2$ . We also have triangles in  $\text{h-proj}(\mathcal{E})$

$$D_i \longrightarrow \tilde{C}_i \longrightarrow \tilde{C}_{D_i} \tag{7.4}$$

that become distinguished in  $H^0(\text{h-proj}(\mathcal{E}))$ , such that  $\text{Hom}_{\text{h-proj}(\mathcal{E})}(\text{h-proj}(\mathcal{D}_i), \tilde{C}_{D_i})$  is acyclic and with  $D_i \in \text{h-proj}(\mathcal{D}_i)$  for  $i = 1, 2$  (see Proposition 3.14). If we apply  $\text{Ind}(Q_i)$  to equation (7.4), we get quasi-isomorphisms in  $\text{h-proj}(\mathcal{E}_{\mathcal{D}_i})$

$$Q_i(\tilde{C}_i) \xrightarrow{\sim} \text{Ind}(Q_i)(\tilde{C}_{D_i}), \tag{7.5}$$

for  $i = 1, 2$ . As the map induced by  $\text{Ind}(Q_i)$

$$\text{Hom}_{\text{h-proj}(\mathcal{E})}(\tilde{C}_{D_i}, \tilde{C}_{D_i}) \longrightarrow \text{Hom}_{\text{h-proj}(\mathcal{E}_{\mathcal{D}_i})}(Q_i(\tilde{C}_{D_i}), Q_i(\tilde{C}_{D_i}))$$

is a quasi-isomorphism for  $i = 1, 2$ , the idempotent  $e'_i$  realising  $C_i$  as a direct summand of  $Q_i(\tilde{C}_i) \cong \text{Ind}(Q_i)(\tilde{C}_{D_i})$  in  $H^0(\text{h-proj}(\mathcal{E}_{\mathcal{D}_i}))$  can be lifted to a closed degree-0 morphism  $e_i: \tilde{C}_{D_i} \rightarrow \tilde{C}_{D_i}$  in  $\text{h-proj}(\mathcal{E})$ . Therefore, we can set

$$\widehat{C}_i := \text{Hocolim}\left(\tilde{C}_{D_i} \xrightarrow{e_i} \tilde{C}_{D_i} \xrightarrow{e_i} \dots\right) \in \text{Ob}(\text{h-proj}(\mathcal{E}))$$

for  $i = 1, 2$ . We claim that the complex  $\text{Hom}_{\text{h-proj}(\mathcal{E})}(\text{h-proj}(\mathcal{D}_i), \widehat{C}_i)$  is acyclic. Indeed, for all  $D \in \mathcal{D}_i$ , we have natural isomorphisms

$$\text{Hom}_{H^0(\text{h-proj}(\mathcal{E}))}(D, \widehat{C}_i) \cong \text{colim Hom}_{H^0(\text{h-proj}(\mathcal{E}))}(D, C_{D_i})$$

and  $\text{Hom}_{H^0(\text{h-proj}(\mathcal{E}))}(D, C_{D_i})$  is trivial by the definition of  $C_{D_i}$ . For the isomorphism, we used that  $D \in \mathcal{D}_i \subseteq \mathcal{E}$  is a compact object in  $H^0(\text{h-proj}(\mathcal{E}))$ . Since  $H^0(\text{h-proj}(\mathcal{D}_i))$  is compactly generated by the objects in  $\mathcal{D}_i$ , the claim follows.

Moreover, there is an isomorphism in  $H^0(\text{h-proj}(\mathcal{E}_{\mathcal{D}_i}))$

$$\text{Ind}(Q_i)(\widehat{C}_i) \cong C_i. \tag{7.6}$$



Indeed, using the fact that  $\text{Ind}(Q_i)$  commutes with coproducts in  $Z^0(\text{h-proj}(\mathcal{E}))$ , we get

$$\begin{aligned} \text{Ind}(Q_i)(\widehat{C}_i) &\cong \text{Hocolim}(\text{Ind}(Q_i)(\widetilde{C}_{D_i}) \rightarrow \text{Ind}(Q_i)(\widetilde{C}_{D_i}) \rightarrow \dots) \\ &\cong \text{Hocolim}(Q_i(\widetilde{C}_i) \xrightarrow{e'_i} Q_i(\widetilde{C}_i) \xrightarrow{e'_i} \dots) \cong C_i. \end{aligned}$$

For the penultimate isomorphism, we used equation (7.5).

Now we can go further and invoke Proposition 3.14 again to get a triangle

$$D \longrightarrow \widehat{C}_2 \longrightarrow C_D, \tag{7.7}$$

with  $D \in \text{Ob}(\text{h-proj}(\mathcal{D}_1))$  and such that it becomes distinguished in  $H^0(\text{h-proj}(\mathcal{E}))$  and the complex  $\text{Hom}_{\text{h-proj}(\mathcal{E})}(\text{h-proj}(\mathcal{D}_i), C_D)$  is acyclic for  $i = 1, 2$ .

Hence we have a sequence of homotopy equivalences

$$\overline{Q}_1(\text{Ind}(Q_1)(\widehat{C}_1)) \cong \overline{Q}_1(C_1) \cong \overline{Q}_2(C_2) \cong \overline{Q}_2(\text{Ind}(Q_2)(\widehat{C}_2)) \cong \overline{Q}_2(\text{Ind}(Q_2)(C_D)). \tag{7.8}$$

The first and the third are those in equation (7.6). The second one is just  $f$  in the given triple  $(C_1, C_2, f)$ . For the last one, observe that

$$H := H^0(\overline{Q}_2) \circ H^0(\text{Ind}(Q_2)) = H^0(\overline{Q}_1) \circ H^0(\text{Ind}(Q_1)).$$

Thus, if we apply  $H$  to the distinguished triangle given by equation (7.7) in  $H^0(\text{h-proj}(\mathcal{E}))$ , we get an isomorphism  $H(\widehat{C}_2) \cong H(C_D)$  because  $H(D) = H^0(\overline{Q}_1)(H^0(\text{Ind}(Q_1))(D)) = 0$ .

In conclusion, we have morphisms

$$\widehat{C}_1 \xrightarrow{f_1} C_D \xleftarrow{f_2} \widehat{C}_2 \tag{7.9}$$

in  $Z^0(\text{h-proj}(\mathcal{E}))$ . We claim that this can be completed to a square

$$\begin{array}{ccc} C & \xrightarrow{g_2} & \widehat{C}_2 \\ g_1 \downarrow & & \downarrow f_2 \\ \widehat{C}_1 & \xrightarrow{f_1} & C_D \end{array}$$

in  $Z^0(\text{h-proj}(\mathcal{E}))$ , which becomes commutative in  $H^0(\text{h-proj}(\mathcal{E}))$ . Indeed, one proceeds as in [36, Section 1.4] and considers the morphism  $\varphi := f_1 + f_2: \widehat{C}_1 \oplus \widehat{C}_2 \rightarrow C_D$  in  $Z^0(\text{h-proj}(\mathcal{E}))$ . We set  $C := \text{Cone}(\varphi)[-1]$  so that we get a triangle

$$C \xrightarrow{\psi} \widehat{C}_1 \oplus \widehat{C}_2 \xrightarrow{\varphi} C_D \tag{7.10}$$

which becomes distinguished in  $H^0(\text{h-proj}(\mathcal{E}))$ . One sets  $g_i$  to be the composition of  $\psi$  with the natural projection  $\widehat{C}_1 \oplus \widehat{C}_2 \rightarrow \widehat{C}_i$ . It is clear from the defining triangle given by equation (7.10) that there are isomorphisms in  $H^0(\text{h-proj}(\mathcal{E}))$

$$\text{Ind}(Q_i)(C) \cong \text{Ind}(Q_i)(\widehat{C}_i) \cong C_i, \tag{7.11}$$

where the last one is equation (7.6).

The object  $C$  is actually compact in  $H^0(\text{h-proj}(\mathcal{E}))$  (and thus it is contained in  $H^0(\mathcal{E}) \cong H^0(\text{Perf}(\mathcal{E}))$ ). Indeed, by equation (7.11),  $\text{Lnd}(\mathcal{Q}_i)(C) \in \mathcal{E}_{\mathcal{D}_i}$  is compact. By (ii) in Setup 7.1, we immediately have that

$$\text{Hom}_{H^0(\text{h-proj}(\mathcal{E}))} \left( H^0(\text{h-proj}(\mathcal{D}_i)), H^0(\text{h-proj}(\mathcal{D}_j)) \right) = 0$$

for  $i \neq j \in \{1, 2\}$ , and thus  $H^0(\text{h-proj}(\mathcal{D}_1))$  and  $H^0(\text{h-proj}(\mathcal{D}_2))$  are localising completely orthogonal full triangulated subcategories of  $H^0(\text{h-proj}(\mathcal{E}))$ . Thus they provide Bousfield localisations. Since  $H^0(\text{h-proj}(\mathcal{D}_i))$  is compactly generated by the objects in  $\mathcal{D}_i$ , which are compact in  $H^0(\text{h-proj}(\mathcal{E}))$ , [41, Corollary 5.12] implies that  $C$  is compact.

In conclusion,  $C \in \mathcal{E}$ ,  $\mathcal{Q}_i(C) \cong C_i$  by equation (7.11); and then, by equation (7.8), we conclude that there is a quasi-isomorphism  $F(C) \cong (C_1, C_2, f)$ . □

### 7.3. Refining the geometric application

We want to extend the discussion further in Example 7.2 and Example 7.3. Assume  $X$  to be a quasi-compact and quasi-separated scheme, and let  $U_1, \dots, U_n \subset X$  be a finite collection of quasi-compact open subsets such that  $X = U_1 \cup \dots \cup U_n$ .

For  $I \subseteq N := \{1, \dots, n\}$ , we set  $U_I := \cap_{i \in I} U_i$ , and we denote by  $\iota_I : U_I \hookrightarrow X$  the corresponding open immersion. Clearly  $U_\emptyset = X$  and  $\iota_\emptyset = \text{id}$ .

**Remark 7.5.** It is clear that  $\cup_{i \in I} U_i$  is quasi-compact and quasi-separated for every  $I \subseteq N$ . Also, the open subsets  $U_I$  must be quasi-compact because they are the intersections of the quasi-compact open subsets  $U_i$  in the quasi-separated  $X$ . Now assume that the open subsets  $U_i \subset X$  are all affine; in this case, for  $I \neq \emptyset$ , the open subset  $U_I$  is also separated, being an open subset of an affine (hence separated) scheme. Therefore, by [5, Corollary 5.5],  $\mathbf{D}_{\text{qc}}^?(U_I) \cong \mathbf{D}^?(U_I)$  for  $I \neq \emptyset$ . Here and in what follows we use the shorthand  $\mathbf{D}^?(U_I) := \mathbf{D}^?(\mathbf{Qcoh}(U_I))$ .

Let  $(\mathcal{E}, E)$  be a dg enhancement either of  $\mathbf{D}_{\text{qc}}^?(X)$  or of  $\mathbf{Perf}(X)$ , and let us assume that  $\mathcal{E}$  is h-flat. We denote by  $\mathcal{D}_I$  the full dg subcategory of  $\mathcal{E}$  defined by

$$\text{Ob}(\mathcal{D}_I) := \{C \in \mathcal{E} : \iota_I^* \circ E(C) \cong 0\}$$

and set  $\mathcal{E}_I := \text{Perf}(\mathcal{E}/\mathcal{D}_I)$ . We have an exact equivalence given by the composition

$$E_I : H^0(\mathcal{E}_I) \xrightarrow{\sim} H^0(\mathcal{E})/H^0(\mathcal{D}_I) \xrightarrow{\sim} \mathbf{D}_{\text{qc}}^?(U_I) \quad (\text{respectively } E_I : H^0(\mathcal{E}_I) \cong \mathbf{Perf}(U_I)).$$

Moreover, the dg quotient functors  $\mathcal{Q}_I : \mathcal{E} \rightarrow \mathcal{E}_I$  are such that the diagram of triangulated categories and exact functors

$$\begin{CD} H^0(\mathcal{E}) @>H^0(\mathcal{Q}_I)>> H^0(\mathcal{E}_I) \\ @V E VV @VV E_I V \\ \mathbf{D}_{\text{qc}}^?(X) @>\iota_I^*>> \mathbf{D}^?(U_I) \end{CD} \tag{7.12}$$

is commutative.

By construction, whenever  $I \subseteq I' \subseteq N$ , there is a natural dg functor  $\mathcal{E}_I \rightarrow \mathcal{E}_{I'}$ , and these functors compose nicely. In particular, as in Remark 3.13, we can form the homotopy limit

$$\mathcal{E}^{\text{hl}} := \text{Holim}_{\emptyset \neq I \subseteq N} \mathcal{E}_I.$$

**Remark 7.6.** In analogy with the usual homotopy limit of categories, we can inductively compute  $\mathcal{C}^{\text{hl}}$  up to quasi-equivalence as follows due to the universal property in Remark 3.13.

Let  $N' := \{1, \dots, n - 1\}$ , and consider the homotopy limits

$$\mathcal{C}' := \text{Holim}_{\emptyset \neq I \subseteq N'} \mathcal{C}_I \quad \mathcal{C}'' := \text{Holim}_{\{n\} \subseteq I \subseteq N} \mathcal{C}_I.$$

By construction, we have natural dg functors

$$\bar{Q}'_1: \mathcal{C}' \longrightarrow \mathcal{C}'' \quad \bar{Q}'_2: \mathcal{C}_{\{n\}} \longrightarrow \mathcal{C}''.$$

Then  $\mathcal{C}^{\text{hl}}$  is the homotopy pullback of the diagram

$$\mathcal{C}' \xrightarrow{\bar{Q}'_1} \mathcal{C}'' \xleftarrow{\bar{Q}'_2} \mathcal{C}_{\{n\}}.$$

For  $n = 2$ , we have the following easy result.

**Lemma 7.7.** *Let  $X$  be a quasi-compact and quasi-separated scheme, and let  $U_1, U_2 \subseteq X$  be quasi-compact open subschemes such that  $X = U_1 \cup U_2$ . Assume that  $\mathcal{C}$  is an  $h$ -flat dg enhancement either of  $\mathbf{D}_{\text{qc}}^?(X)$  or of  $\mathbf{Perf}(X)$  for  $? = b, +, -, \emptyset$ . Then there is a quasi-equivalence  $\mathcal{C} \rightarrow \mathcal{C}^{\text{hl}}$ .*

*Proof.* By Remark 7.6,  $\mathcal{C}^{\text{hl}}$  is just the pullback of the diagram

$$\mathcal{C}_{\{1\}} \longrightarrow \mathcal{C}_{\{1,2\}} \longleftarrow \mathcal{C}_{\{2\}}.$$

Since, by Example 7.2 and Example 7.3, the assumptions in Setup 7.1 are verified (with  $\mathcal{C}_{\{i\}} = \mathcal{C}_{\mathcal{D}_i}$  for  $i = 1, 2$ , and  $\mathcal{C}_{\{1,2\}} = \mathcal{C}_{\mathcal{D}_{1,2}}$ ), the result follows from Theorem 7.4.  $\square$

The following is the natural generalisation of Lemma 7.7.

**Proposition 7.8.** *Let  $X$  be a quasi-compact and quasi-separated scheme, and let  $\mathcal{C}$  be an  $h$ -flat dg enhancement either of  $\mathbf{D}_{\text{qc}}^?(X)$  or of  $\mathbf{Perf}(X)$  for  $? = b, +, -, \emptyset$ . Then there is a quasi-equivalence  $\mathcal{C} \rightarrow \mathcal{C}^{\text{hl}}$ .*

*Proof.* The argument is by induction on  $n$ , the number of quasi-compact open subsets in the covering  $\{U_1, \dots, U_n\}$  of  $X$ . Clearly, if  $n = 1$ , there is nothing to prove. Thus we can assume  $n \geq 2$  and set

$$V_1 := U_1 \cup \dots \cup U_{n-1} \quad V_2 := U_n.$$

Clearly,  $V_i$  is quasi-compact for  $i = 1, 2$ . Define  $\mathcal{C}_{\mathcal{D}_i}$ ,  $\mathcal{C}_{\mathcal{D}_{1,2}}$ ,  $\mathbf{Q}_i$  and  $\bar{\mathbf{Q}}_i$  as in Example 7.2 or Example 7.3. In particular,  $\mathcal{C}_{\mathcal{D}_i}$  and  $\mathcal{C}_{\mathcal{D}_{1,2}}$  are dg enhancements of  $\mathbf{D}_{\text{qc}}^?(V_i)$  and  $\mathbf{D}_{\text{qc}}^?(V_1 \cap V_2)$  (or of  $\mathbf{Perf}(V_i)$  and  $\mathbf{Perf}(V_1 \cap V_2)$ ) for  $i = 1, 2$ .

Now, consider the homotopy limits  $\mathcal{C}'$  and  $\mathcal{C}''$  in Remark 7.6. By induction, based on Remark 7.6 and Lemma 7.7, it is easy to see that  $\mathcal{C}'$  is a dg enhancement of  $\mathbf{D}_{\text{qc}}^?(V_1)$  and  $\mathcal{C}''$  is a dg enhancement of  $\mathbf{D}_{\text{qc}}^?(V_1 \cap V_2)$  (respectively,  $\mathbf{Perf}(V_1)$  and  $\mathbf{Perf}(V_1 \cap V_2)$ ). The induction gives more precise information: there are quasi-equivalences  $F': \mathcal{C}_{\mathcal{D}_1} \rightarrow \mathcal{C}'$  and  $F'': \mathcal{C}_{\mathcal{D}_{1,2}} \rightarrow \mathcal{C}''$ , making the diagrams

$$\begin{array}{ccccc} \mathcal{C}_{\mathcal{D}_1} & \xrightarrow{\bar{\mathbf{Q}}_1} & \mathcal{C}_{\mathcal{D}_{1,2}} & \xleftarrow{\bar{\mathbf{Q}}_2} & \mathcal{C}_{\mathcal{D}_2} \\ F' \downarrow & & \downarrow F'' & & \parallel \\ \mathcal{C}' & \xrightarrow{\bar{\mathbf{Q}}'_1} & \mathcal{C}'' & \xleftarrow{\bar{\mathbf{Q}}'_2} & \mathcal{C}_{\{n\}} \end{array} \tag{7.13}$$

commutative in  $\mathbf{dgCat}$ . Note that the induction applies to  $\mathcal{C}''$  as well, since  $V_1 \cap V_2 = (U_1 \cap U_n) \cup \dots \cup (U_{n-1} \cap U_n)$  is the union of  $n - 1$  quasi-compact open subsets.

By Lemma 7.7, there is a quasi-equivalence  $F_1 : \mathcal{C} \rightarrow \mathcal{C}_{\mathcal{D}_1} \times_{\mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}}^h \mathcal{C}_{\mathcal{D}_2}$ . By equation (7.13) and [20, Proposition 13.3.9], we get a quasi-equivalence

$$F_2 : \mathcal{C}_{\mathcal{D}_1} \times_{\mathcal{C}_{\mathcal{D}_1, \mathcal{D}_2}}^h \mathcal{C}_{\mathcal{D}_2} \longrightarrow \mathcal{C}' \times_{\mathcal{C}''}^h \mathcal{C}_{\{n\}},$$

where the latter dg category is  $\mathcal{C}^{hl}$  by Remark 7.6. Hence  $F := F_2 \circ F_1 : \mathcal{C} \rightarrow \mathcal{C}^{hl}$  is the quasi-equivalence we are looking for.  $\square$

In Proposition 7.8 (and in Lemma 7.7), we always assumed that the dg enhancement  $\mathcal{C}$  is h-flat. If not, we can take the quasi-equivalence  $l_{\mathcal{C}}^{hf} : \mathcal{C}^{hf} \rightarrow \mathcal{C}$  in Proposition 3.10. By Proposition 7.8, we get a quasi-equivalence

$$F : \mathcal{C}^{hf} \longrightarrow (\mathcal{C}^{hf})^{hl}.$$

Then the composition  $[F] \circ [l_{\mathcal{C}}^{hf}]^{-1}$  yields an isomorphism between  $\mathcal{C}$  and  $(\mathcal{C}^{hf})^{hl}$  in  $\mathbf{Hqe}$ .

In conclusion, we have the following.

**Corollary 7.9.** *Let  $X$  be a quasi-compact and quasi-separated scheme, and let  $\mathcal{C}$  be a dg enhancement either of  $\mathbf{D}_{qc}^?(X)$  or of  $\mathbf{Perf}(X)$  for  $? = b, +, -, \emptyset$ . Then there is an isomorphism  $\mathcal{C} \cong (\mathcal{C}^{hf})^{hl}$  in  $\mathbf{Hqe}$ .*

### 8. Uniqueness of enhancements for geometric categories

In this section, we prove Theorem B; more precisely, the proof is in Section 8.1 and Section 8.2. In Section 8.3, we show how a weaker version of Theorem B, for the category of perfect complexes, can be obtained by ‘simpler’ means.

#### 8.1. Proof of Theorem B: the case of $\mathbf{D}_{qc}^?(X)$

Write  $X$  as a finite union  $X = U_1 \cup \dots \cup U_n$  of affine open subschemes. The argument is based on induction on  $n$ .

When  $X$  is affine (or, more generally, quasi-compact and separated),  $\mathbf{D}_{qc}^?(X) \cong \mathbf{D}^?(Qcoh(X))$  by [5, Corollary 5.5], whence  $\mathbf{D}_{qc}^?(X)$  has a unique enhancement by Theorem A (1). Thus we may assume  $n \geq 2$ .

Suppose that  $(\mathcal{C}, E)$  is a dg enhancement of  $\mathbf{D}_{qc}^?(X)$ . Up to replacing  $\mathcal{C}$  with the quasi-equivalent dg category  $\mathbf{Perf}(\mathcal{C})$ , we can assume that  $\mathcal{C}$  satisfies equation (4.1). Now we construct the dg category  $\widehat{\mathcal{B}} := \widehat{\mathcal{B}}_{\mathcal{C}, E}(\mathbf{Mod}(X), Qcoh(X))$ , together with a fully faithful dg functor  $\widehat{\mathcal{B}} \rightarrow \mathcal{C}$ . In view of Proposition 4.1, up to replacing  $\mathcal{C}$  with the quasi-equivalent dg category  $\mathcal{C}^{hf}$ , we can assume that  $\mathcal{C}$  is h-flat.

Now we can define the dg categories  $\mathcal{C}_I$  and the dg functors  $Q_I : \mathcal{C} \rightarrow \mathcal{C}_I$  as in Section 7.3, and we keep the same notation as in that section. For every subset  $\emptyset \neq I \subseteq N = \{1, \dots, n\}$ , as  $\mathcal{C}_I$  has the same objects as  $\mathcal{C}$ , we can construct a dg category  $\mathcal{B}_I$  with the same objects as  $\widehat{\mathcal{B}}$  and with

$$\mathrm{Hom}_{\mathcal{B}_I}(B_1, B_2) := \mathrm{Hom}_{\mathcal{C}_I}(B_1^- \oplus B_1^+, B_2^- \oplus B_2^+).$$

Moreover, by construction, the dg functor  $Q_I$  induces a natural dg functor

$$\widehat{Q}_I : \widehat{\mathcal{B}} \longrightarrow \mathcal{B}_I.$$

**Remark 8.1.** Arguing as in Remark 4.3, we see that the image of  $H^0(\widehat{\mathcal{B}})$  under the composition

$$H^0(\mathcal{C}) \xrightarrow{H^0(Q_I)} H^0(\mathcal{C}_I) \xrightarrow{E_I} \mathbf{D}_{qc}^?(U_I) \cong \mathbf{D}^?(U_I) := \mathbf{D}^?(Qcoh(U_I))$$

(see Remark 7.5 for the last equivalence) coincides with  $\mathbf{B}^2(\mathbf{Qcoh}(U_I))$ . This simple observation follows from the definition of  $\widehat{\mathcal{B}}$  together with the fact that a quasi-coherent sheaf on  $U_I$  extends to a quasi-coherent sheaf on  $X$ , and the functor

$$H^0(\widehat{\mathcal{B}}) \rightarrow H^0(\mathcal{C}) \xrightarrow{\mathbb{E}} \mathbf{D}_{\mathbf{qc}}^2(X)$$

surjects on complexes of quasi-coherent sheaves with zero differentials. Hence,  $H^0(\mathbf{Q}_I)(H^0(\widehat{\mathcal{B}})) \cong H^0(\mathcal{B}_I) \cong \mathbf{B}^2(\mathbf{Qcoh}(U_I))$  and the dg functor  $\mathcal{B}_I \rightarrow \mathcal{C}_I$  induces a quasi-equivalence  $\text{Perf}(\mathcal{B}_I) \cong \mathcal{C}_I$  because of Corollary 2.10. Moreover, if  $\emptyset \neq I \subseteq I' \subseteq N$ , we have dg functors

$$\widetilde{\mathbf{Q}}_{I,I'}: \text{Perf}(\mathcal{B}_I) \longrightarrow \text{Perf}(\mathcal{B}_{I'})$$

corresponding to the restriction  $\mathbf{D}^2(U_I) \rightarrow \mathbf{D}^2(U_{I'})$ . In particular, these dg functors compose nicely.

By the invariance of homotopy limits under quasi-equivalences (see Remark 3.13), we have a quasi-equivalence


$$\text{Holim Perf}(\mathcal{B}_I) \longrightarrow \mathcal{C}^{\text{hl}}.$$

On the other hand, by Proposition 7.8 (see also Corollary 7.9), we have a quasi-equivalence  $\mathcal{C} \longrightarrow \mathcal{C}^{\text{hl}}$ . Putting it all together, we get an isomorphism

$$\mathcal{C} \cong \text{Holim Perf}(\mathcal{B}_I) \tag{8.1}$$

in **Hqe**.

By Remark 5.5, given  $I$ , as a consequence of our proof of Theorem A (1) for the uniqueness of dg enhancements of  $\mathbf{D}^2(U_I)$ , we get full dg subcategories  $\mathcal{N}_I \subseteq \text{Perf}(\mathcal{V}_I)$ ,  $\widetilde{\mathcal{S}}_I \subseteq \text{Perf}(\widetilde{\mathcal{B}}_I)$ ,  $\overline{\mathcal{S}}_I \subseteq \text{Perf}(\overline{\mathcal{B}}_I)$ ,  $\mathcal{S}'_I \subseteq \text{Perf}(\mathcal{B}'_I)$ ,  $\mathcal{S}_I \subseteq \text{Perf}(\mathcal{B}_I)$  and  $\mathcal{S}_I \subseteq \text{Perf}(\mathcal{B}_I)$  and the commutative diagram of zigzags



where the dg functors in the bottom zigzag are quasi-equivalences, the vertical maps are Drinfeld dg quotient functors such that  $\text{Perf}(\mathcal{B}'_I) \rightarrow \text{Perf}(\mathcal{B}'_I)/\mathcal{S}'_I$  and  $\text{Perf}(\mathcal{B}_I) \rightarrow \text{Perf}(\mathcal{B}_I)/\mathcal{S}_I$  are quasi-equivalences. Most importantly, the lower zigzag remains functorial in  $I$  (by Remark 3.11).

Thus, by Remark 3.13, we get the following chain of isomorphisms in **Hqe**:

$$\begin{aligned} \text{Holim Perf}(\mathcal{V}_I)/\mathcal{N}_I &\cong \text{Holim Perf}(\widetilde{\mathcal{B}}_I)/\widetilde{\mathcal{S}}_I \cong \text{Holim Perf}(\overline{\mathcal{B}}_I)/\overline{\mathcal{S}}_I \cong \text{Holim Perf}(\mathcal{B}'_I)/\mathcal{S}'_I \\ &\cong \text{Holim Perf}(\mathcal{B}'_I)/\mathcal{S}'_I \cong \text{Holim Perf}(\mathcal{B}_I)/\mathcal{S}_I \cong \text{Holim Perf}(\mathcal{B}_I). \end{aligned}$$

Together with equation (8.1), we get an isomorphism  $\text{Holim Perf}(\mathcal{V}_I)/\mathcal{N}_I \cong \mathcal{C}$  in **Hqe**. But the construction of  $\mathcal{N}_I$  and of the dg functors  $\mathcal{V}_I \rightarrow \mathcal{V}_{I'}$  (and thus of the induced ones  $\text{Perf}(\mathcal{V}_I)/\mathcal{N}_I \rightarrow \text{Perf}(\mathcal{V}_{I'})/\mathcal{N}_{I'}$ ) are independent of the given dg enhancement  $(\mathcal{C}, \mathbb{E})$ . Indeed,  $\mathcal{N}_I$  is an enhancement of  $\mathbf{K}_{\text{acy}}^2(U_I)$ , and the dg functors are induced by the open immersions  $\iota_{I,I'}$  above.

In conclusion, all dg enhancements of  $\mathbf{D}_{\mathbf{qc}}^2(X)$  are isomorphic in **Hqe** to  $\text{Holim Perf}(\mathcal{V}_I)/\mathcal{N}_I$ , and thus  $\mathbf{D}_{\mathbf{qc}}^2(X)$  has a unique dg enhancement.

8.2. Proof of Theorem B: the case of  $\mathbf{Perf}(X)$

The case of perfect complexes is treated with a strategy that is essentially identical to the one in Section 8.1: we write  $X$  as a finite union  $X = U_1 \cup \dots \cup U_n$  of affine open subschemes, we proceed by induction on  $n$  and finish by using Section 7.3 in an essential way. The main difference is that instead of following the construction of  $\widetilde{\mathcal{B}}$  and  $\mathcal{V}$  in Section 4.1 and Section 4.2, we will partially use the strategy used in [27] to prove Theorem 6.10.

If  $n = 1$ , then  $X$  is affine and the result follows from [27, Proposition 2.6]. Thus we can assume  $n \geq 2$  and start proving the following simple result.

**Lemma 8.2.** *Let  $X$  be an affine scheme and  $\iota: U \hookrightarrow X$  be an open subscheme. Then  $\mathbf{D}_{\mathbf{qc}}(U)$  is compactly generated by the compact object  $\mathcal{O}_U = \iota^* \mathcal{O}_X$  and  $\mathbf{Perf}(U) = \langle \mathcal{O}_U \rangle_{\infty}$ .*

*Proof.* Let  $Z := X \setminus U$ , and denote by  $\mathbf{D}_Z(X)$  the full triangulated subcategory of  $\mathbf{D}_{\mathbf{qc}}(X)$  consisting of complexes with (topological) support contained in the closed subset  $Z$ . Consider the short exact sequence of triangulated categories

$$0 \longrightarrow \mathbf{D}_Z(X) \longrightarrow \mathbf{D}_{\mathbf{qc}}(X) \xrightarrow{\iota^*} \mathbf{D}_{\mathbf{qc}}(U) \longrightarrow 0.$$

By [41, Theorem 6.8],  $\mathbf{D}_Z(X)$  is compactly generated by objects that are compact in  $\mathbf{D}_{\mathbf{qc}}(X)$ . Hence, by [34, Theorem 2.1],  $\iota^*$  sends the compact generator  $\mathcal{O}_X$  to the compact generator  $\mathcal{O}_U$  and  $\mathbf{D}_{\mathbf{qc}}(U)^c = \langle \mathcal{O}_U \rangle_{\infty}$ . Since  $\mathbf{Perf}(U) = \mathbf{D}_{\mathbf{qc}}(U)^c$  (see Example 7.3), this concludes the proof.  $\square$

Hence, in our situation,  $\mathbf{Perf}(U_I) = \langle \mathcal{O}_{U_I} \rangle_{\infty}$  for all  $\emptyset \neq I \subseteq N$ . This implies that given a dg enhancement  $(\mathcal{C}, E)$  of  $\mathbf{Perf}(X)$  inducing equivalences  $E_I: H^0(\mathcal{C}_I) \xrightarrow{\sim} \mathbf{Perf}(U_I)$ , we can take the dg category  $\mathcal{N}_I \subseteq \mathcal{C}_I$  with only one object  $O_I$ , which is the lift of  $\mathcal{O}_{U_I}$  along  $E_I$ . By the above result, the inclusion of  $\mathcal{N}_I$  in  $\mathcal{C}_I$  induces a quasi-equivalence  $\mathbf{Perf}(\mathcal{N}_I) \cong \mathcal{C}_I$ . Moreover, as in the previous section, the quotient dg functors  $\mathcal{C}_I \rightarrow \mathcal{C}_{I'}$  for  $\emptyset \neq I \subseteq I' \subset N$ , induce dg functors  $\mathcal{N}_I \rightarrow \mathcal{N}_{I'}$  and thus dg functors

$$\widetilde{Q}_{I,I'}: \mathbf{Perf}(\mathcal{N}_I) \longrightarrow \mathbf{Perf}(\mathcal{N}_{I'})$$

with the same properties as in Remark 8.1.

On the other hand,  $\mathcal{O}_{U_I}$  as an object of  $\mathbf{Perf}(U_I)$  generates a dg category  $\mathcal{A}_I$  all sitting in degree zero. The open inclusion  $\iota_{I,I'}: U_{I'} \hookrightarrow U_I$  induces a dg functor

$$\widetilde{Q}'_{I,I'}: \mathcal{A}_I \longrightarrow \mathcal{A}_{I'}.$$

The argument in [27, Section 6] applies in this case, and thus:

- (a) There is a morphism  $\mathbf{Perf}(\mathcal{A}_I) \rightarrow \mathbf{Perf}(\mathcal{N}_I)$  in  $\mathbf{Hqe}$  that can be represented by a roof

$$\begin{array}{ccc} & \mathbf{Perf}(\widetilde{\mathcal{N}}_I) & \\ \swarrow & & \searrow \\ \mathbf{Perf}(\mathcal{A}_I) & & \mathbf{Perf}(\mathcal{N}_I), \end{array}$$

which is functorial in  $I$ .<sup>2</sup>

- (b) A full dg subcategory  $\mathcal{L}_I \subseteq \mathbf{Perf}(\mathcal{A}_I)$  such that the idempotent completion

$$\widehat{\mathcal{A}}_I := (\mathbf{Perf}(\mathcal{A}_I) / \mathcal{L}_I)^{\mathrm{ic}}$$

<sup>2</sup>The argument in [27] actually provides a composition of roofs  $\mathcal{A}_I \rightarrow H^0(\mathcal{N}_I)$  and  $H^0(\mathcal{N}_I) \hookrightarrow \tau_{\leq 0}(\mathcal{N}_I) \rightarrow \mathcal{N}_I$ . But this composition can be easily rearranged in the form above.

of the Drinfeld quotient is a dg enhancement of  $\mathbf{Perf}(U_I)$ . Moreover, the definition of  $\widehat{\mathcal{A}}_I$  is functorial in  $I$ , and both its definition and that of the natural dg functors  $\widehat{\mathcal{A}}_I \rightarrow \widehat{\mathcal{A}}_{I'}$  induced by  $\widetilde{\mathcal{Q}}_{I,I'}$  are independent of the dg enhancement  $(\mathcal{C}, \mathbf{E})$ .

Here, as in the previous section, we freely use the possibility of taking functorial h-flat resolutions. As a result of Lemma 6.2 and Lemma 6.3 in [27], we get a roof of quasi-equivalences

$$\begin{array}{ccc} & \mathbf{Perf}(\widetilde{\mathcal{N}}_I)/\mathcal{L}'_I & \\ \swarrow & & \searrow \\ \widehat{\mathcal{A}}_I & & \mathbf{Perf}(\mathcal{N}_I), \end{array}$$

where  $\mathcal{L}'_I$  is a full pretriangulated dg subcategory of  $\mathbf{Perf}(\widetilde{\mathcal{N}}_I)$  identified with  $\mathcal{L}_I$  under the quasi-equivalence  $\mathbf{Perf}(\widetilde{\mathcal{N}}_I) \rightarrow \widehat{\mathcal{A}}_I$  and the diagram is functorial in  $I$ .

In conclusion, as in the previous section, we have a chain of isomorphism in **Hqe**

$$\mathcal{C} \cong \mathcal{C}^{\text{hl}} \cong \text{Holim } \mathbf{Perf}(\mathcal{N}_I) \cong \text{Holim } \widehat{\mathcal{A}}_I$$

due to Proposition 7.8 and the invariance of homotopy limits under quasi-equivalences. Thus  $\mathbf{Perf}(X)$  has a unique enhancement since the latter homotopy limit does not depend on  $(\mathcal{C}, \mathbf{E})$ . Therefore, the proof of Theorem B is complete.

**Remark 8.3.** As we observed in Example 7.3, the triangulated category  $\mathbf{Perf}(X)$  identifies with  $\mathbf{D}_{\text{qc}}(X)^c$  when  $X$  is a quasi-compact and quasi-separated scheme. It is then easy (see, for example, the argument in [9, Remark 5.5]) to deduce the uniqueness of enhancement of  $\mathbf{D}_{\text{qc}}(X)$  from that of  $\mathbf{Perf}(X)$ . The complications in Section 8.1 emerge while dealing with  $\mathbf{D}_{\text{qc}}^?(X)$ , when  $? = b, +, -$ .

### 8.3. An aside

It is worth pointing out that a weak version of Theorem B for  $\mathbf{Perf}(X)$  can be proved using techniques very similar to the ones in [13].

The simple observation is that assumption (2) in [13, Theorem B] can be replaced by the weaker (ii) below to get the following slightly more general result.

**Theorem 8.4.** *Let  $\mathcal{G}$  be a Grothendieck category with a small set  $\mathcal{A}$  of generators such that*

- (i)  $\mathcal{A}$  is closed under finite coproducts;
- (ii) If  $f: \coprod_{i \in I} C_i \rightarrow C$  (with  $I$  a small set) is a morphism in  $\mathcal{G}$  and  $A \in \text{Ob}(\mathcal{A})$  is a subobject of  $C$  such that  $A \subseteq \text{im } f$ , then there exists a finite subset  $I'$  of  $I$  such that  $A \subseteq f(\coprod_{i \in I'} C_i)$ ;
- (iii) If  $f: A' \twoheadrightarrow A$  is an epimorphism of  $\mathcal{G}$  with  $A, A' \in \mathcal{A}$ , then  $\ker f \in \mathcal{A}$ ;
- (iv) For every  $A \in \mathcal{A}$ , there exists  $N(A) > 0$  such that  $\text{Hom}_{\mathbf{D}(\mathcal{G})}(A, A'[N(A)]) = 0$  for every  $A' \in \text{Ob}(\mathcal{A})$ .

Then  $\mathbf{D}(\mathcal{G})^c$  has a unique enhancement.

Actually, it is very easy to see that the argument used in [13] to prove the theorem still works with the new hypothesis and proves Theorem 8.4. Indeed, in [13, Remark 6.3], it is proved, in particular, that the old assumption (2) implies (ii). Moreover, in the other parts of the proof of [13, Theorem B], where (2) was used (in the proofs of Lemma 6.4 and Proposition 6.5, where Remark 6.3 is invoked), precisely (ii) is needed.

As an application, we obtain the following improvement of [13, Proposition 6.10] in the case of schemes. Recall that a scheme  $X$  has enough perfect coherent sheaves if  $\mathbf{Qcoh}(X)$  is generated, as a Grothendieck category, by a small set of objects in  $\mathbf{Coh}(X) \cap \mathbf{Perf}(X)$ .

**Proposition 8.5.** *Let  $X$  be a quasi-compact and semi-separated scheme with enough perfect coherent sheaves. Then  $\mathbf{Perf}(X)$  has a unique enhancement.*

*Proof.* The argument of the proof of [13, Proposition 6.10] (with the same  $\mathcal{A}$ ) shows that conditions (i), (iii) and (iv) of Theorem B are satisfied. As for (ii), one is reduced to proving the following statement: if  $\mathcal{F} = \bigcup_{i \in I} \mathcal{F}_i \in \mathbf{Coh}(X)$ , with  $\mathcal{F}_i$  a quasi-coherent subsheaf of  $\mathcal{F}$  for every  $i \in I$ , then there exists a finite subset  $I'$  of  $I$  such that  $\mathcal{F} = \bigcup_{i \in I'} \mathcal{F}_i$ . Since  $X$  (being quasi-compact) has a finite affine open cover, we can also assume that  $X$  is affine, in which case the statement follows directly from the fact that  $\mathcal{F}$  is of finite type.  $\square$

Proposition 8.5 is a strictly stronger version than [13, Proposition 6.10], as is demonstrated by the following example.

**Example 8.6.** If  $X$  is an affine scheme such that  $\mathcal{O}_X$  is coherent, then the hypotheses of Proposition 8.5 are satisfied, since  $\mathcal{O}_X \in \mathbf{Coh}(X) \cap \mathbf{Perf}(X)$  is a generator of  $\mathbf{Qcoh}(X)$ . On the other hand, if  $X$  is not noetherian, then the hypotheses of [13, Proposition 6.10] are not satisfied. An explicit example is given by  $X = \text{Spec}(A)$  with  $A$  the polynomial ring in infinitely many variables over a field.

We conclude our presentation by recalling that in [1, Corollary 9], the author proved that  $\mathbf{Perf}(X)$  has a unique enhancement when  $X$  is a quasi-compact, quasi-separated and 0-complicial scheme. We refer to [1] for the precise definition of 0-complicial that is not needed here. On the other hand, it is not complicated to see that a quasi-compact and semi-separated scheme with enough perfect coherent sheaves is 0-complicial and thus [1, Corollary 9] should be more general than Proposition 8.5. It would be interesting to find an example of a scheme satisfying the assumptions in [1, Corollary 9] but not the assumptions in the above proposition.

**Acknowledgements.** It is our great pleasure to thank Benjamin Antieau, Pieter Belmans, Henning Krause and Valerio Melani for insightful conversations about several problems related to the contents of this paper. We are also very grateful to Michel Van den Bergh, Alexander Kuznetsov, Valery Lunts and Bregje Pauwels for comments on a preliminary version of this paper.

**Conflicts of Interest.** The authors have no conflicts of interest to declare.

**Financial support.** A. C. was partially supported by the research project PRIN 2017 ‘Moduli and Lie Theory’. A. N. was partially supported by grants DP150102313 and DP200102537 from the Australian Research Council. P. S. was partially supported by the ERC Consolidator Grant ERC-2017-CoG-771507-StabCondEn, by the research project PRIN 2017 ‘Moduli and Lie Theory’ and by the research project FARE 2018 HighCaSt (grant number R18YA3ESPJ).

## References

- [1] B. Antieau, ‘On the uniqueness of infinity-categorical enhancements of triangulated categories’, [arXiv:1812.01526](https://arxiv.org/abs/1812.01526).
- [2] B. Antieau, D. Gepner, J. Heller, ‘ $K$ -theoretic obstructions to bounded  $t$ -structures’, *Invent. Math.* **216** (2019), 241–300.
- [3] P. Balmer, M. Schlichting, ‘Idempotent Completion of Triangulated Categories’, *J. Algebra* **236** (2001), 819–834.
- [4] A. A. Beilinson, J. Bernstein, P. Deligne, ‘Faisceaux pervers’, *Astérisque* **100** (1982), 5–171.
- [5] M. Bökstedt, A. Neeman, ‘Homotopy limits in triangulated categories’, *Comp. Math.* **86** (1993), 209–234.
- [6] A. Bondal, M. Larsen, V. Lunts, ‘Grothendieck ring of pretriangulated categories’, *Int. Math. Res. Not.* **29** (2004), 1461–1495.
- [7] A. Bondal, M. Van den Bergh, ‘Generators and representability of functors in commutative and noncommutative geometry’, *Moscow Math. J.* **3** (2003), 1–36.
- [8] A. Canonaco, M. Orsnaghi, P. Stellari, ‘Localizations of the category of  $A_\infty$ -categories and internal Homs’, *Doc. Math.* **24** (2019), 2463–2492.
- [9] A. Canonaco, P. Stellari, ‘A tour about existence and uniqueness of dg enhancements and lifts’, *J. Geom. Phys.* **122** (2017), 28–52.
- [10] A. Canonaco, P. Stellari, *Fourier–Mukai functors: a survey*, EMS Ser. Congr. Rep., Eur. Math. Soc. (2013), 27–60.
- [11] A. Canonaco, P. Stellari, ‘Internal Homs via extensions of dg functors’, *Adv. Math.* **277** (2015), 100–123.
- [12] A. Canonaco, P. Stellari, ‘Non-uniqueness of Fourier–Mukai kernels’, *Math. Z.* **272** (2012), 577–588.
- [13] A. Canonaco, P. Stellari, ‘Uniqueness of dg enhancements for the derived category of a Grothendieck category’, *J. Eur. Math. Soc.* **20** (2018), 2607–2641.
- [14] A. Canonaco, P. Stellari, Uniqueness of dg enhancements for the derived category of a Grothendieck category, [arXiv:1507.05509v5](https://arxiv.org/abs/1507.05509v5).
- [15] L. Cohn, Differential graded categories are  $k$ -linear stable infinity categories, [arXiv:1308.2587](https://arxiv.org/abs/1308.2587).
- [16] V. Drinfeld, ‘DG quotients of DG categories’, *J. Algebra* **272** (2004), 643–691.



- [17] D. Dugger, B. Shipley, 'A curious example of triangulated-equivalent model categories which are not Quillen equivalent, Algebr'. *Geom. Topol.* **9** (2009), 135–166.
- [18] F. Genovese, Quasi-functors as lifts of Fourier-Mukai functors: the uniqueness problem, PhD thesis, available at: [https://fgenovese1987.github.io/documents/thesis\\_phd.pdf](https://fgenovese1987.github.io/documents/thesis_phd.pdf).
- [19] A. Grothendieck, J.-L. Verdier, 'Préfaïceaux', in *SGA 4, Théorie des Topos et Cohomologie Etale des Schémas, Tome 1. Théorie des Topos, Lect. Notes in Math.* **269**, Springer, Heidelberg, 1972–1973, pp. 1–184.
- [20] P. Hirschhorn, *Model Categories and Their Localizations, Math. Surveys and Monographs Series* **99**, Am. Math. Soc. (2003), xvi+457.
- [21] M. Hovey, *Model Categories, Mathematical Surveys and Monographs* **63**, Am. Math. Soc. (1998), xii+209.
- [22] G. M. Kelly, 'Chain maps inducing zero homology maps', *Proc. Cambridge Philos. Soc.* **61** (1965), 847–854.
- [23] H. Krause, 'Deriving Auslander's formula', *Documenta Math.* **20** (2015), 669–688.
- [24] H. Krause, 'On Neeman's well generated triangulated categories', *Documenta Math.* **6** (2001), 119–125.
- [25] H. Krause, 'Localization theory for triangulated categories', in: *Triangulated Categories, London Math. Soc. Lecture Note Ser.* **375**, Cambridge Univ. Press (2010), 161–235.
- [26] G. Laumon, L. Moret-Bailly, *Champs algébriques, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* **39**, Springer-Verlag (2000), xii+208.
- [27] V. Lunts, D. Orlov, 'Uniqueness of enhancements for triangulated categories', *J. Amer. Math. Soc.* **23** (2010), 853–908.
- [28] V. Lunts, O.M. Schnürer, 'New enhancements of derived categories of coherent sheaves and applications', *J. Algebra* **446** (2016), 203–274.
- [29] J. Lurie, Higher algebra, <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [30] J. Lurie, Spectral algebraic geometry, <https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>.
- [31] A. Neeman, 'A counterexample to vanishing conjectures for negative K-theory', *Invent. Math.* **225** (2021), 427–452.
- [32] A. Neeman, 'Non-left-complete derived categories', *Math. Res. Lett.* **18** (2011), 827–832.
- [33] A. Neeman, 'On the derived category of sheaves on a manifold', *Documenta Math.* **6** (2001), 483–488.
- [34] A. Neeman, 'The connection between the K-theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel', *Ann. Sci. École Norm. Sup.* **25** (1992), 547–566.
- [35] A. Neeman, 'The homotopy category of injectives', *Algebra Number Theory* **8** (2014), 429–456.
- [36] A. Neeman, *Triangulated Categories*, Annals of Mathematics Studies **148**, Princeton University Press (2001), viii+449.
- [37] A. Neeman, 'Approximable triangulated categories', in: *Representations of Algebras, Geometry and Physics, Contemp. Math.* **769**, Amer. Math. Soc. (2021), 111–155.
- [38] A. Rizzardo, M. Van den Bergh, A. Neeman, 'An example of a non-Fourier–Mukai functor between derived categories of coherent sheaves', *Invent. Math.* **216** (2019), 927–1004.
- [39] A. Rizzardo, M. Van den Bergh, 'A note on non-unique enhancements', *Proc. Amer. Math. Soc.* **147** (2019), 451–453.
- [40] A. Rizzardo, M. Van den Bergh, 'A  $k$ -linear triangulated category without a model', *Ann. Math.* **191** (2020), 393–437.
- [41] R. Rouquier, 'Dimensions of triangulated categories', *J. K-theory* **1** (2008), 193–258.
- [42] M. Schlichting, 'A note on K-theory and triangulated categories', *Invent. Math.* **150** (2002), 111–116.
- [43] M. Schlichting, 'Negative K-theory of derived categories', *Math. Z.* **253** (2006), 97–134.
- [44] G. Tabuada, 'Une structure de catégorie de modèles de Quillen sur la catégorie des dg-catégories', *C. R. Math. Acad. Sci. Paris* **340** (2005), 15–19.
- [45] R. W. Thomason, T. Trobaugh, *Higher Algebraic K-Theory of Schemes and of Derived Categories, The Grothendieck Festschrift III*, Birkhäuser, Boston, Basel, Berlin, (1990), 247–436.
- [46] B. Toën, 'The homotopy theory of dg-categories and derived Morita theory', *Invent. Math.* **167** (2007), 615–667.
- [47] J.-L. Verdier, 'Des catégories dérivées des catégories abéliennes', *Astérisque* **239** (1996), xii+253 pp.
- [48] S. Virili, 'Morita theory for stable derivators', [arXiv:1807.01505](https://arxiv.org/abs/1807.01505).