



ANNALES DE L'INSTITUT FOURIER

Luca F. DI CERBO & Luigi LOMBARDI

**L^2 -Betti Numbers and Convergence of Normalized Hodge
Numbers via the Weak Generic Nakano Vanishing Theorem**

Article à paraître, mis en ligne le 3 juillet 2023, 27 p.

Article mis à disposition par ses auteurs selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE



<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les *Annales de l'Institut Fourier* sont membres du
Centre Mersenne pour l'édition scientifique ouverte

www.centre-mersenne.org

e-ISSN : 1777-5310

L^2 -BETTI NUMBERS AND CONVERGENCE OF NORMALIZED HODGE NUMBERS VIA THE WEAK GENERIC NAKANO VANISHING THEOREM

by **Luca F. DI CERBO & Luigi LOMBARDI** (*)

ABSTRACT. — We study the rate of growth of normalized Hodge numbers along a tower of abelian covers of a smooth projective variety with semismall Albanese map. These bounds are in some cases optimal. Moreover, we compute the L^2 -Betti numbers of irregular varieties that satisfy the weak generic Nakano vanishing theorem (e.g., varieties with semismall Albanese map). Finally, we study the convergence of normalized plurigenera along towers of abelian covers of any irregular variety. As an application, we extend a result of Kollár concerning the multiplicativity of higher plurigenera of a smooth projective variety of general type, to a wider class of varieties. In the Appendix, we study irregular varieties for which the first Betti number diverges along a tower of abelian covers induced by the Albanese variety.

RÉSUMÉ. — Nous étudions le taux de croissance des nombres de Hodge normalisés le long d'une tour de revêtement abéliennes d'une variété projective lisse avec l'application d'Albanese semi-petite. Ces bornes sont dans certains cas optimales. De plus, nous calculons les nombres de Betti L^2 des variétés irrégulières qui satisfont le théorème d'annulation générique faible de Nakano (e.g., variétés avec l'application d'Albanese semi-petite). Enfin, nous étudions la convergence de plurigenres normalisés le long de tours de revêtement abéliennes de toute variété irrégulière. On applique ça à l'extension d'un résultat de Kollár concernant la multiplicativité des plurigenres supérieurs d'une variété projective lisse de type général, à une classe plus large de variétés. En annexe, nous étudions les variétés irrégulières pour lesquelles le premier nombre de Betti diverge le long d'une tour de revêtement abéliennes induite par la variété d'Albanese.

Keywords: L^2 -Betti Numbers, Normalized Hodge Numbers, Irregular Varieties.
2020 Mathematics Subject Classification: 14F06, 32J25, 32L20.

(*) The first author is partially supported by NSF grant DMS-2104662.

The second author is partially supported by the Rita Levi-Montalcini Program, Simons Foundation, SIR 2014: AnHyC: “Analytic aspects in complex and hypercomplex geometry” (code RBSI14DYEB), and Grant 261756 of the Research Councils of Norway.

1. Introduction and Main Results

In this paper, we study the asymptotic behavior of Hodge and Betti numbers of sequences of coverings of complex projective varieties with semismall Albanese map. Similar problems have attracted considerable interest over the last four decades, and they have been extensively studied in a variety of different geometric contexts. For instance, in [9] and [10], DeGeorge and Wallach study the asymptotic behavior of limit multiplicities of representations in L^2 of discrete co-compact lattices of isometries of symmetric varieties. In cohomological terms, they study the asymptotic behavior of Betti numbers on regular coverings of compact locally symmetric spaces of non-compact type (e.g., compact real and complex hyperbolic manifolds). The problem addressed in [9, 10] is natural for researchers interested in the cohomology of locally symmetric varieties, and it can be easily described. In what follows, we rephrase the main result in [9, 10] in terms of *normalized* Betti numbers. We refer to page 714 in the introduction of [2], or to Chapter 5 in the book [31], for more details concerning the connections between the representation theoretic results of DeGeorge and Wallach, and the asymptotic properties of the cohomology of compact locally symmetric varieties.

Given a torsion free lattice Γ acting co-compactly on a symmetric space of non-compact type, say G/K , a sequence of nested, normal, finite index subgroups $\{\Gamma_i\}$ of Γ is a cofinal filtration of Γ if $\cap_i \Gamma_i$ is the identity element. Define $\pi_i: X_i \rightarrow X$ as the finite index regular cover of $X \stackrel{\text{def}}{=} \Gamma \backslash (G/K)$ associated to Γ_i . The main result of [9] implies that

$$\lim_{i \rightarrow \infty} \frac{b_k(X_i)}{\deg \pi_i} = 0 \quad \text{for any } k \neq \frac{1}{2} \dim(G/K),$$

where $b_k(X_i)$ denotes the k -th Betti number of X_i . We refer to the ratio $b_k(X_i)/\deg \pi_i$ as the *normalized* k -Betti number of the cover $\pi_i: X_i \rightarrow X$. Thus, for k different from the middle dimension, the growth of Betti numbers in a tower of coverings associated to a cofinal filtration has sub-degree (or sub-volume) growth, and the normalized Betti numbers converge to zero.

The study of Betti numbers in a sequence of coverings continues to fascinate many mathematicians; see for example the recent work of Abert et al. [2]. In this remarkable paper, the authors extend the results of DeGeorge–Wallach to sequences of compact locally symmetric varieties which Benjamini–Schramm converge to their universal covers. We refer to [2] for the precise definition of this notion of convergence; here we

simply remark that a tower of coverings associated to a cofinal filtration does indeed Benjamini–Schramm converge. The techniques employed both in [9, 10] and [2] are based on representation theory, and they do not immediately generalize to non-symmetric varieties. Nevertheless, there is a large and growing literature concerning these kind of problems outside the locally symmetric context; see for example [1], [12], [42] and the bibliography therein. These papers employ geometric analysis techniques, and they extend much of the DeGeorge–Wallach theory to negatively curved compact Riemannian manifolds which are *not* locally symmetric.

Here we contribute to this circle of ideas by studying the cohomology of complex projective varieties with *semismall* Albanese map, a further instance of varieties of non-locally symmetric type. Our approach is based on tools of Algebraic Geometry and Hodge Theory, and it employs sheaf-theoretic techniques specific to this class of varieties. As an important ingredient, we employ the generic vanishing theory of bundles of holomorphic p -forms developed by Popa and Schnell in [34] via Saito’s theory of mixed Hodge modules.

We now turn to details and present our main results. Let X be an irregular smooth projective complex variety of dimension n , and let $a_X : X \rightarrow \text{Alb}(X)$ be its Albanese map. The Albanese torus $\text{Alb}(X)$ is an abelian variety of dimension $g = h^{1,0}(X)$ (we recall that the variety X is irregular if $g > 0$). We say that the Albanese map a_X is *semismall* if for every integer $k > 0$ the following inequalities hold

$$(1.1) \quad \text{codim}\{x \in a_X(X) \mid \dim(a_X^{-1}(x)) \geq k\} \geq 2k.$$

In particular, if a_X is semismall, then a_X is generically finite onto its image, but the converse does not hold in general. For instance, the Albanese map of the blow-up of an abelian variety along a smooth subvariety of codimension c is semismall if and only if $c \leq 2$. Next, let

$$\mu_d : \text{Alb}(X) \rightarrow \text{Alb}(X), \quad \mu_d(x) = dx = \overbrace{x + \cdots + x}^{d\text{-times}}, \quad d \geq 1$$

be the multiplication maps on $\text{Alb}(X)$, and define the varieties X_d via the fiber product diagrams

$$(1.2) \quad \begin{array}{ccc} X_d & \xrightarrow{a_d} & \text{Alb}(X) \\ \downarrow \varphi_d & & \downarrow \mu_d \\ X & \xrightarrow{a_X} & \text{Alb}(X). \end{array}$$

Our first result controls the rate of growth of the Hodge numbers of X_d with respect to the degrees of the covers $\varphi_d: X_d \rightarrow X$. We refer to the ratios $h^{p,q}(X_d)/\deg \varphi_d$ as the *normalized* (p, q) -Hodge numbers. The following theorem provides an *effective* estimate for the rate of convergence of the normalized Hodge numbers, and it also yields the optimal rate of convergence of one of them.

THEOREM 1.1. — *Let X be a smooth projective variety of complex dimension n , and let $\varphi_d: X_d \rightarrow X$ be the étale covers defined in (1.2). If the Albanese map a_X is semismall, then for any pair of integers $(p, q) \in [0, n]^2$ there exists a positive constant $B(p, q)$ such that*

$$(1.3) \quad \frac{h^{p,q}(X_d)}{\deg \varphi_d} \leq B(p, q) d^{-2|n-p-q|} \quad \text{for all } d \geq 1.$$

Moreover, we have

$$(1.4) \quad \lim_{d \rightarrow \infty} \frac{h^{p,q}(X_d)}{\deg \varphi_d} = (-1)^q \chi(\Omega_X^p) \quad \text{if } p + q = n.$$

Conversely, if X is a smooth projective variety of dimension n that satisfies both $\dim \text{Alb}(X) > n$ and the bounds in (1.3) for all pairs of indexes $(p, q) \in [0, n]^2$, then the Albanese map a_X is semismall.

In order to prove the previous theorem, in Section 3 we develop a general machinery that establishes the convergence of the normalized cohomology ranks $h^q(X_d, \varphi_d^* \mathcal{F})/\deg \varphi_d$ of a coherent sheaf \mathcal{F} on X subject to certain cohomological conditions (cf. Theorem 3.6). In particular, Theorem 1.1 corresponds to the case of bundles of holomorphic p -forms $\mathcal{F} = \Omega_X^p$. In Section 5, we apply this machinery to the case of pluricanonical bundles $\mathcal{F} = \omega_X^{\otimes m}$ for $m \geq 1$. We refer to Section 4 for the details of the proof of Theorem 1.1, and to a generalization that takes into account all values of the defect of semismallness of the Albanese map (cf. Theorem 4.3 and (2.1)). Finally, Theorem 1.1 implies the following statement regarding the normalized Betti numbers.

COROLLARY 1.2. — *Let X be a smooth projective variety of dimension n such that the Albanese map a_X is semismall. Then for any integer $k \neq n$ there exists a positive constant $C(k)$ such that*

$$\frac{b_k(X_d)}{\deg \varphi_d} \leq C(k) d^{-2|n-k|} \quad \text{for all } d \geq 1.$$

Furthermore, we have

$$\lim_{d \rightarrow \infty} \frac{b_n(X_d)}{\deg \varphi_d} = (-1)^n \chi_{\text{top}}(X).$$

When combined with Lück’s Approximation Theorem (cf. [30, Main Theorem]), Theorem 1.1 can be used to compute the L^2 -Betti numbers of the Albanese universal cover $\bar{\pi}: \bar{X} \rightarrow X$, when the Albanese map of X is semismall. Throughout the paper, the *Albanese universal cover* is defined as the pullback of a_X via the universal topological cover π of $\text{Alb}(X)$:

$$(1.5) \quad \begin{array}{ccc} \bar{X} & \xrightarrow{\bar{a}} & \mathbb{C}^g \\ \downarrow \bar{\pi} & & \downarrow \pi \\ X & \xrightarrow{a_X} & \text{Alb}(X), \end{array}$$

where $g = h^{1,0}(X) = \dim \text{Alb}(X) \neq 0$. Notice that, up to a finite cover, the Albanese universal cover coincides with the universal abelian cover. Indeed, these infinite covers are equal if and only if $H_1(X, \mathbb{Z})$ is torsion free. We refer to Section 6 for the formal definition of L^2 -Betti numbers of any infinite G -covering map $X' \rightarrow X'/G$. It turns out that our calculation of L^2 -Betti numbers holds for a more general class of smooth irregular projective varieties, which we now define. We say that X satisfies the *weak generic Nakano vanishing theorem* if for any pair of integers $(p, q) \in [0, n]^2$ such that $p + q \neq n$ we have

$$H^q(X, \Omega_X^p \otimes \alpha_{p,q}) = 0$$

for at least one topologically trivial line bundle $\alpha_{p,q} \in \text{Pic}^0(X)$. Instances of varieties that satisfy this property are varieties with semismall Albanese map (cf. Theorem 2.6), and varieties that admit one holomorphic 1-form such that its zero-set is either finite or empty (cf. Theorem 2.7). We refer to [13] and [27, Introduction, Sections 3.1 and 3.2] for examples and basic properties of this class of varieties.

THEOREM 1.3. — *Let X be a smooth projective variety of complex dimension n and let \bar{X} be the universal Albanese cover. If X satisfies the weak generic Nakano vanishing theorem, then the L^2 -Betti numbers of \bar{X} are:*

$$b_k^{(2)}(\bar{X}) = \begin{cases} (-1)^n \chi_{\text{top}}(X) & \text{if } k = n \\ 0 & \text{if } k \neq n. \end{cases}$$

It is tantalizing to compare Theorem 1.3 with an old conjecture of Singer concerning the L^2 -Betti numbers of the universal covering space of an aspherical manifold.

CONJECTURE 1.4 (Singer Conjecture). — *If X is a closed aspherical manifold of real dimension $2n$, then*

$$b_k^{(2)}(\tilde{X}) = \begin{cases} (-1)^n \chi_{\text{top}}(X) & \text{if } k = n \\ 0 & \text{if } k \neq n, \end{cases}$$

where $\pi: \tilde{X} \rightarrow X$ is the topological universal cover of X .

Interestingly, Theorem 1.3 provides a vanishing theorem analogous to Singer's conjecture when the L^2 -Betti numbers are computed with respect to the Albanese universal cover. It seems worth asking whether Theorem 1.3 holds when the L^2 -Betti numbers are computed with respect to the topological universal cover, and more generally, if Singer's conjecture can be extended meaningfully outside the class of aspherical manifolds, at least within the class of projective varieties.

We point out that in [20, Theorem 3(i)] Jost and Zuo prove, among other things, a special case of Theorem 1.3. More specifically, they prove Theorem 1.3 in the case of smooth projective varieties whose Albanese map $a_X: X \rightarrow \text{Alb}(X)$ is an immersion. The techniques used by Jost and Zuo rely on analytical arguments introduced by Gromov in [16], where the author confirms Singer's conjecture for *Kähler hyperbolic* manifolds. These manifolds include Kähler manifolds with negative and pinched sectional curvature, and do *not* contain any rational curve. On the contrary, varieties with Albanese map semismall may contain rational curves. Finally, we remark that in [25] the semismallness condition of the Albanese map is studied by means of topological generic vanishing theory. Via the general strategy of [26, Theorem 2.28], it is possible that the techniques of [25, Theorem 1.2] suffice to give an alternative proof of Theorem 1.3 in the case of varieties with semismall Albanese map; however we do not pursue this direction in this paper. Finally, we also point out the related work of Budur [6], where the author shows polynomial periodicity of the Hodge numbers of congruence covers.

In Section 5, we apply the techniques of Section 3 to prove a version of Theorem 1.1 for pluricanonical bundles $\omega_X^{\otimes m}$ with $m \geq 2$. More precisely, we compute the following limits

$$(1.6) \quad \tilde{P}_m(X) \stackrel{\text{def}}{=} \lim_{d \rightarrow \infty} \frac{P_m(X_d)}{\deg \varphi_d} = \lim_{d \rightarrow \infty} \frac{h^0(X_d, \omega_{X_d}^{\otimes m})}{\deg \varphi_d}, \quad m \geq 2$$

of normalized plurigenera (whenever they exist). Let $I: X \rightarrow Z$ be a smooth representative of the Iitaka fibration, and let $q(I) = q(X) - q(Z)$ be the difference of the irregularities. In Proposition 5.2, we prove that the

limits in (1.6) exist and are computed by:

$$(1.7) \quad \tilde{P}_m(X) = \begin{cases} P_m(X) & \text{if } q(I) = 0, \\ 0 & \text{if } q(I) > 0. \end{cases}$$

We recall that if X is of general type (hence satisfying $q(I) = 0$), then a classical result of Kollár [22, Proposition 9.4] (cf. also [24, Theorem 11.2.23]) ensures that its higher plurigenera are multiplicative with respect to *any* étale cover. As suggested by (1.7), we extend this property to smooth projective varieties satisfying $q(I) = 0$, when the étale covers are induced by the Albanese variety via base change. Also, as a by-product, we show that [22, Proposition 9.4] cannot be extended to varieties with $q(I) > 0$. We refer to Section 5 for the proof of Theorem 5.4, and examples of varieties with $q(I) = 0$ that are not of general type.

In the Appendix (Section A), we discuss the irregular varieties for which the first Betti number b_1 goes to infinity along the unramified covers induced by the multiplication maps on the Albanese variety (regardless of the semismallness of the Albanese map). Building upon results of Beauville [3], we prove that if this is the case, then the base variety must be fibered over a curve having either genus at least two, or genus equal to one and the fibration admits two multiple fibers whose multiplicities are not coprime. Moreover, if the group $H^2(X, \mathbb{Z})$ is torsion free, then the converse of this result holds as well (cf. Theorem A.1). We remark that, in many interesting cases, the converse can be used to deduce that the first Betti number is indeed uniformly bounded on these abelian covers. Very recently, Stover [40] and Vidussi [41] study the boundedness of the first Betti number of abelian covers of the Cartwright–Steger surface [8]. While our analysis does not fully recover their theorems, it has the advantage to put in perspective their results in the framework of higher-dimensional varieties.

Acknowledgments

The authors thank Rob Lazarsfeld and Christian Schnell for their interest and suggestions, Stefano Vidussi and Botong Wang for email correspondence, and the anonymous referee for pertinent and constructive comments on a preliminary version of this work. They thank the Mathematics Department of Stony Brook University for the ideal research environment they enjoyed at the beginning of this project. The first named author thanks the University of Florida where most of these ideas crystallized, and where some of the results were presented in the Fall 2018 Topology Seminar. During the

preparation of this work, the second named author visited both the University of Florida and the Max Planck Institute for Mathematics in Bonn. He thanks these institutions for their hospitality and support. Finally, he thanks Daniele Angella of the University of Florence for financial support.

2. Weak GV -Sheaves

In this section, we recall a few basic results from generic vanishing theory. The following presentation is tailored to our purposes; we refer to [13, 14, 33, 38] for a comprehensive introduction.

Let X be a smooth projective complex variety of dimension n , and let $f: X \rightarrow A$ be a morphism to an abelian variety of dimension g . The *non-vanishing loci* attached to a coherent sheaf \mathcal{F} on X relative to $f: X \rightarrow A$ are defined as

$$V^i(\mathcal{F}) = \{ \alpha \in \widehat{A} \mid H^i(X, \mathcal{F} \otimes f^*\alpha) \neq 0 \} \quad (i \geq 0)$$

(in the notation $V^i(\mathcal{F})$ we omit the reference to the morphism f). Here $\widehat{A} \simeq \text{Pic}^0(A)$ denotes the dual torus of A , which parameterizes isomorphism classes of holomorphic line bundles with trivial first Chern class. By the Semicontinuity Theorem [19, Theorem III.12.8], the loci $V^i(\mathcal{F})$ are algebraic closed subsets of \widehat{A} .

DEFINITION 2.1. — *The sheaf \mathcal{F} satisfies GV (or the generic vanishing property) if $\text{codim}_{\widehat{A}} V^i(\mathcal{F}) \geq i$ for all $i > 0$.*

A fundamental result of Green and Lazarsfeld proves that if the Albanese map $a_X: X \rightarrow \text{Alb}(X)$ is generically finite onto its image, then the canonical bundle ω_X satisfies GV . Moreover, the loci $V^i(\omega_X)$ are *torsion linear* varieties for all $i \geq 0$ regardless the Albanese dimension of X , i.e. every irreducible component $T \subset V^i(\omega_X)$ is of type $\beta + T_0$ where $\beta \in \text{Pic}^0(\text{Alb}(X)) \simeq \text{Pic}^0(X)$ is an element of finite order, and $T_0 \subset \text{Pic}^0(X)$ is a subtorus (cf. [14, Theorem 0.1], [39] and [38, Corollary 19.2]). For the purposes of this paper, we will consider the following weaker notion of generic vanishing.

DEFINITION 2.2. — *The sheaf \mathcal{F} satisfies weak GV with index p if $V^i(\mathcal{F}) \subsetneq \widehat{A}$ for all $i \neq p$.*

Obviously, GV -sheaves satisfy weak GV with index 0. We conclude this subsection with a useful result which we will use in Section 5. The Euler characteristic of a sheaf \mathcal{F} is defined as $\chi(\mathcal{F}) = \sum_{i \geq 0} (-1)^i h^i(X, \mathcal{F})$.

LEMMA 2.3. — *If \mathcal{F} is a weak GV-sheaf with index p , then*

$$\chi(\mathcal{F}) = (-1)^p h^p(X, \mathcal{F} \otimes f^* \alpha)$$

for a generic element $\alpha \in \widehat{A}$. In particular, $\chi(\mathcal{F}) = 0$ if \mathcal{F} is a weak GV-sheaf with respect to two distinct indexes.

Proof. — If $\alpha \in \widehat{A}$ is generic, then the cohomology groups $H^i(X, \mathcal{F} \otimes f^* \alpha)$ vanish for all $i \neq p$. Since $\chi(\mathcal{F})$ is invariant under twists with line bundles in $\text{Pic}^0(X)$, we find $\chi(\mathcal{F}) = \chi(\mathcal{F} \otimes f^* \alpha) = (-1)^p h^p(X, \mathcal{F} \otimes f^* \alpha)$. Moreover, if \mathcal{F} is a weak GV-sheaf with respect to two distinct indexes, then all the loci $V^i(\mathcal{F})$ are proper subset of \widehat{A} , hence $\chi(\mathcal{F}) = 0$. \square

2.1. (Weak) Generic Nakano Vanishing Theorem

Let X be a smooth projective variety of dimension n , and let $a_X : X \rightarrow \text{Alb}(X)$ be its Albanese map. Moreover, denote by $\Omega_X^p \stackrel{\text{def}}{=} \wedge^p \Omega_X$ the bundle of holomorphic p -forms on X . Following [34, Definition 12.1], we say that X satisfies the *generic Nakano vanishing theorem* if $\text{codim}_{\text{Pic}^0(X)} V^q(\Omega_X^p) \geq |p + q - n|$ for all indexes p and q . In this paper, we consider varieties that satisfy a weaker vanishing condition.

DEFINITION 2.4. — *The variety X satisfies the weak generic Nakano vanishing theorem if Ω_X^p is a weak GV-sheaf with index $n - p$ for all $p = 0, \dots, n$.*

It turns out that X satisfies the generic Nakano vanishing theorem if and only if it satisfies a condition on the dimension of the fibers of the Albanese map. This goes as follows. Set $V_l \stackrel{\text{def}}{=} \{y \in \text{Alb}(X) \mid \dim a_X^{-1}(y) \geq l\}$ and define the *defect of semismallness* of a_X as:

$$(2.1) \quad \delta(a_X) = \max_{l \in \mathbb{N}} \{2l - n + \dim V_l\}.$$

DEFINITION 2.5. — *We say that a_X is semismall if $\delta(a_X) = 0$. Equivalently, a_X is semismall if the inequalities of (1.1) are satisfied for all $k \geq 1$.*

THEOREM 2.6 (Popa–Schnell). — *If X is a smooth projective variety of dimension n , then*

$$\text{codim}_{\text{Pic}^0(X)} V^q(\Omega_X^p) \geq |p + q - n| - \delta(a_X)$$

for all $p \geq 0$ and $q \geq 0$. Moreover, there exists a pair (p, q) for which the equality is attained. In particular, if a_X is semismall, then X satisfies the generic Nakano vanishing theorem.

The previous theorem appears in [34, Theorem 3.2], and it is proved by means of Saito's theory of mixed Hodge modules and the Fourier–Mukai transform. Besides varieties with semismall Albanese map, another class of varieties that satisfies Definition 2.4 is provided by the following result of Green and Lazarsfeld [13, Theorem 3.1].

THEOREM 2.7 (Green–Lazarsfeld). — *Let X be a smooth projective variety. If X carries a holomorphic 1-form such that its zero-set is either finite or empty, then X satisfies the weak generic Nakano vanishing theorem.*

The previous theorem relies on the deformation theory of the derivative complexes associated to Ω_X^p . A natural question is the characterization of varieties that satisfy Definition 2.4. Here we note that a variety that satisfies the weak generic Nakano vanishing theorem does not necessarily carry a holomorphic 1-form whose zero-set is either finite or empty. For instance, consider a smooth projective variety Y of general type such that its Albanese map is an immersion and $\text{codim}_{\text{Alb}(Y)} Y = 2$ (for instance a genus 3 curve in its Jacobian). Then the blow-up Z of $\text{Alb}(Y)$ along Y is the counterexample we are looking for. In fact, by [35, Theorem 2.1], any holomorphic 1-form on Y has at least one zero, and its pull-back to Z vanishes along some curves in the exceptional divisor. Moreover, all 1-forms of Z are obtained in this way as $H^0(Z, \Omega_Z) = H^0(\text{Alb}(Y), \Omega_{\text{Alb}(Y)}) = H^0(Y, \Omega_Y)$. On the other hand, the Albanese map of Z is semismall so that Z satisfies the generic Nakano vanishing theorem.

3. Limits of Normalized Cohomology Ranks

Let X be a smooth projective variety of complex dimension n , and $f: X \rightarrow A$ be a morphism to an abelian variety, as in Section 2. Given any integer $d \geq 1$, we denote by $\mu_d: A \rightarrow A$ the multiplication map $\mu_d(x) = dx$. Furthermore, by means of the fiber product construction, we define the varieties X_d as follows:

$$(3.1) \quad \begin{array}{ccc} X_d & \xrightarrow{f_d} & A \\ \downarrow \varphi_d & & \downarrow \mu_d \\ X & \xrightarrow{f} & A. \end{array}$$

In general the varieties X_d may be disconnected, but if $f = a_X$ is the Albanese map they are irreducible. Finally, we set $\mathcal{F}_d \stackrel{\text{def}}{=} \varphi_d^* \mathcal{F}$ if \mathcal{F} is

a coherent sheaf on X . In this section we aim to calculate the following limits of *normalized cohomology ranks*:

$$\liminf_{d \rightarrow \infty} \frac{h^p(X_d, \mathcal{F}_d)}{\deg \varphi_d} \quad \text{and} \quad \limsup_{d \rightarrow \infty} \frac{h^p(X_d, \mathcal{F}_d)}{\deg \varphi_d}.$$

To this end, we introduce first some notation. We denote by r_i the number of irreducible components of $V^i(\mathcal{F})$, and by v_i the maximum dimension of an irreducible component of $V^i(\mathcal{F})$. Moreover, we set:

$$M_i = \max \{ h^i(X, \mathcal{F} \otimes f^* \alpha) \mid \alpha \in V^i(\mathcal{F}) \}$$

$$m_i = \min \{ h^i(X, \mathcal{F} \otimes f^* \alpha) \mid \alpha \in V^i(\mathcal{F}) \}.$$

Finally, we denote by T_d^i the set of d -torsion points of $V^i(\mathcal{F})$, and by $\tau_d^i = |T_d^i|$ its cardinality. We use the following lemma in order to bound τ_d^i .

LEMMA 3.1. — *Let V be a complex torus and $S = p_0 + B$ be a translate of a subtorus $B \subset V$, and let $d \geq 1$ be an integer. If the set of d -torsion points of S is not empty, then it consists of exactly $d^{2 \dim B}$ elements.*

Proof. — Denote by $\nu_d(x) = dx$ the multiplication map on B . We notice that if $y = p_0 + x \in S = p_0 + B$ is a d -torsion point, then $dx = -dp_0$. Hence x is an element of the fiber $\nu_d^{-1}(-dp_0)$ which consists of exactly $d^{2 \dim B}$ elements. Conversely, if $x \in \nu_d^{-1}(-dp_0)$, then $y = p_0 + x$ belongs to S and it is trivially a d -torsion point. \square

DEFINITION 3.2. — *The locus $V^i(\mathcal{F})$ is said linear (resp. torsion linear) if it consists of a finite union of translates (resp. torsion translates) of subtori of \widehat{A} .*

PROPOSITION 3.3. — *If $V^i(\mathcal{F})$ is linear, then for all $d \geq 1$ the following inequalities hold:*

- (i) $\tau_d^i \leq r_i d^{2v_i}$,
- (ii) $\sum_{\alpha \in T_d^i} h^i(X, \mathcal{F} \otimes f^* \alpha) \leq M_i r_i d^{2v_i}$.

Proof. — The proposition follows by Lemma 3.1 and the following inequalities

$$(3.2) \quad m_i \tau_d^i \leq \sum_{\alpha \in T_d^i} h^i(X, \mathcal{F} \otimes f^* \alpha) \leq M_i \tau_d^i. \quad \square$$

PROPOSITION 3.4. — *If $V^i(\mathcal{F})$ is torsion linear and $v_i > 0$, then $\tau_d^i \geq d^{2v_i}$ and $\sum_{\alpha \in T_d^i} h^i(X, \mathcal{F} \otimes f^* \alpha) \geq m_i d_i^{2v_i}$ for infinitely many $d \geq 1$.*

Proof. — If S is a component of dimension v_i , then it contains d -torsion points for infinitely many $d \geq 1$. The result follows by Lemma 3.1 and (3.2). \square

There exist upper bounds on the cardinalities τ_d^i even if $V^i(\mathcal{F})$ is not linear.

PROPOSITION 3.5. — *There are positive constants a_1, a_2 such that for all $d \geq 1$ we have:*

- (i) $\tau_d^i \leq a_1 d^{2v_i}$,
- (ii) $\sum_{\alpha \in T_d^i} h^i(X, \mathcal{F} \otimes f^* \alpha) \leq a_2 d^{2v_i}$.

Proof. — We employ the following theorem of Raynaud [37, p. 327]. Let Y be a closed integral subscheme of a complex abelian variety V , and let $T \subset V$ be the set of torsion points. If $T \cap Y$ is dense in Y with respect to the Zariski topology, then Y is a translate of an abelian subvariety by a point of finite order.

Take now the Zariski closure of all the torsion points in $V^i(\mathcal{F})$. This is a finite union of irreducible closed subvarieties where in each component the torsion points are dense. Hence, by Raynaud's Theorem, each component is a translate of an abelian subvariety of dimension at most v_i by a torsion point. \square

The following theorem is the main result of this section. The equation (3.3) is a generalization of [43, Theorem 4.1] in which the author studies the particular case of the structure sheaf of a smooth projective variety with respect to the Albanese map.

THEOREM 3.6. — *If $V^i(\mathcal{F})$ is a proper subset of \widehat{A} , then we have*

$$(3.3) \quad \lim_{d \rightarrow \infty} \frac{h^i(X_d, \mathcal{F}_d)}{\deg \varphi_d} = 0.$$

Moreover, if \mathcal{F} satisfies weak GV with index p , then we have

$$\lim_{d \rightarrow \infty} \frac{h^p(X_d, \mathcal{F}_d)}{\deg \varphi_d} = (-1)^p \chi(\mathcal{F}).$$

Proof. — Denote by S_d the set of all d -torsion points of \widehat{A} so that

$$\mu_{d*} \mathcal{O}_A \simeq \bigoplus_{\alpha \in S_d} \alpha$$

(cf. [43, Proof of Theorem 4.1]). As both μ_d and φ_d are étale morphisms, there are isomorphisms of complexes $\mathbf{R}\mu_{d*} \mathcal{O}_A \simeq \mu_{d*} \mathcal{O}_A$ and $\mathbf{R}\varphi_{d*} \mathcal{O}_{X_d} \simeq \varphi_{d*} \mathcal{O}_{X_d}$. Hence, by performing the base change of [5, Lemma 1.3] along $f: X \rightarrow A$, we obtain a further decomposition:

$$\varphi_{d*} \mathcal{O}_{X_d} \simeq \bigoplus_{\alpha \in S_d} f^* \alpha.$$

Finally, by the projection formula of [19, Example 8.3], we obtain the following isomorphisms

$$\varphi_{d*}\mathcal{F}_d \simeq \mathcal{F} \otimes \varphi_{d*}\mathcal{O}_{X_d} \simeq \bigoplus_{\alpha \in S_d} (\mathcal{F} \otimes f^*\alpha),$$

so that

$$(3.4) \quad h^i(X_d, \mathcal{F}_d) = h^i(X, \varphi_{d*}\mathcal{F}_d) = \sum_{\alpha \in S_d} h^i(X, \mathcal{F} \otimes f^*\alpha).$$

Hence, if $V^i(\mathcal{F}) = \emptyset$, then all summands in the right hand side of (3.4) are equal to zero. On the other hand, if $V^i(\mathcal{F}) \neq \emptyset$, then Proposition 3.5 yields

$$\sum_{\alpha \in S_d} h^i(X, \mathcal{F} \otimes f^*\alpha) \leq a_i d^{2v_i}$$

for some positive constants a_i which are independent of d . This proves the first claim as $\deg \varphi_d = \deg \mu_d = d^{2g}$ and $v_i < g$.

In order to prove the second claim, we recall that the Euler characteristic $\chi(\mathcal{F})$ is multiplicative under étale covers [23, Proposition 1.1.28], i.e., $\chi(\mathcal{F}_d) = (\deg \varphi_d) \chi(\mathcal{F})$. Therefore an application of (3.3) gives

$$\chi(\mathcal{F}) = \lim_{d \rightarrow \infty} \frac{\chi(\mathcal{F}_d)}{\deg \varphi_d} = \lim_{d \rightarrow \infty} (-1)^p \frac{h^p(X_d, \mathcal{F}_d)}{\deg \varphi_d}. \quad \square$$

Remark 3.7. — By Corollary 2.3, the Euler characteristic of a weak GV -sheaf with index p satisfies $\chi(\mathcal{F}) = (-1)^p h^p(X, \mathcal{F} \otimes f^*\alpha)$, for some line bundle α generic in \widehat{A} . Therefore, if $h^p(X, \mathcal{F})$ assumes the least (or generic) value in the set $\{h^p(X, \mathcal{F} \otimes f^*\alpha) \mid \alpha \in \widehat{A}\}$, then the computation of $\chi(\mathcal{F})$ simplifies to

$$\chi(\mathcal{F}) = (-1)^p h^p(X, \mathcal{F}).$$

This is the case if the sheaf \mathcal{F} satisfies the *Index Theorem with index p* (or *I.T.* for short), namely that $V^i(\mathcal{F}) = \emptyset$ for all $i \neq p$. In fact, by the invariance of the Euler characteristic, it follows that $h^p(X, \mathcal{F} \otimes f^*\alpha)$ is independent on α and $V^p(\mathcal{F}) = \widehat{A}$.

Example 3.8. — By Mumford’s Index Theorem [32, Section 16], any non-degenerate line bundle L on A satisfies the Index Theorem with index p , for some $p \in [0, g = \dim A]$ (see Remark 3.7; moreover note that $p = 0$ if and only if L is ample). Therefore, by taking $f = \text{id}_A$, we have

$$\lim_{d \rightarrow \infty} \frac{h^p(A, L_d)}{\deg \mu_d} = (-1)^p \chi(L) = (-1)^p \frac{(L^g)}{g!}.$$

There are examples of higher rank vector bundles that satisfy the *I.T.* condition as well, for instance, the class of non-degenerate simple semi-homogeneous vector bundles on an abelian variety (cf. [15, Proposition 2.1]).

4. Limits of Normalized Hodge and Betti Numbers

We denote by

$$h^{p,q}(X) = \dim_{\mathbb{C}} H^q(X, \Omega_X^p)$$

the Hodge numbers of a smooth projective variety X , and by

$$(4.1) \quad b_k(X) = \sum_{p+q=k} h^{p,q}(X)$$

its Betti numbers.

PROPOSITION 4.1. — *Let X be a smooth projective variety of dimension n that satisfies the weak generic Nakano vanishing theorem. Then we have*

$$\lim_{d \rightarrow \infty} \frac{h^{p,q}(X_d)}{\deg \varphi_d} = \begin{cases} (-1)^q \chi(\Omega_X^p) & \text{if } p+q = n \\ 0 & \text{if } p+q \neq n \end{cases}$$

and

$$\lim_{d \rightarrow \infty} \frac{b_k(X_d)}{\deg \varphi_d} = \begin{cases} (-1)^n \chi_{\text{top}}(X) & \text{if } k = n \\ 0 & \text{if } k \neq n. \end{cases}$$

Proof. — Since $h^{p,q}(X_d) = h^q(X_d, \Omega_{X_d}^p) = h^q(X_d, \varphi_d^* \Omega_X^p)$, the first statement is an application of Theorem 3.6. For $k \neq n$, the second statement follows by the first statement and the equations (4.1). For $k = n$, we further observe that

$$\sum_{p=0}^n (-1)^{n-p} \chi(\Omega_X^p) = (-1)^n \chi_{\text{top}}(X). \quad \square$$

Remark 4.2. — Proposition 4.1 may fail if the Albanese map is only generically finite onto its image, but not semismall (cf. [20, Remark on p. 6]). A counterexample is provided by the construction in [13, Section 3] (or [38, Example 9.1]) which we here briefly recall. Let A be an abelian variety of dimension four and let $a_X: X \rightarrow A$ be the blowup of A along a smooth curve $C \subset A$ of genus $g(C) \geq 2$ with exceptional divisor E . Hence a_X is the Albanese map of X and $\delta(a_X) = 1$ (see [34, Example 12.3]). By means of the exact sequence $0 \rightarrow a_X^* \Omega_A \rightarrow \Omega_X \rightarrow \Omega_{E/C} \rightarrow 0$ and the Leray spectral sequence, we deduce that

$$V^i(\Omega_X) = \{\mathcal{O}_X\}, \quad V^2(\Omega_X) = \widehat{A}, \quad V^3(\Omega_X) \subseteq \{\mathcal{O}_X\}, \quad i = 0, 1, 4.$$

Hence Ω_X satisfies weak GV with index 2, and, by Theorem 3.6, we find

$$\lim_{d \rightarrow \infty} \frac{h^{1,2}(X_d)}{\deg \varphi_d} = \chi(\Omega_X) = \chi(\Omega_{E/C}) = g(C) - 1 \neq 0.$$

Moreover, as the loci $V^3(\mathcal{O}_X) \simeq V^0(\Omega_X^3) \simeq V^1(\omega_X)$ are of codimension at least one (see [13, Theorem 1]), by Theorem 3.6 we have that also the following limit

$$\lim_{d \rightarrow \infty} \frac{b_3(X_d)}{\deg \varphi_d} = \lim_{d \rightarrow \infty} \frac{\sum_{p+q=3} h^{p,q}(X_d)}{\deg \varphi_d} = 2 \lim_{d \rightarrow \infty} \frac{h^{1,2}(X_d)}{\deg \varphi_d} = 2g(C) - 2$$

is non-zero.

Now we prove Theorem 1.1 of the Introduction. The theorem is a special case of the following more general result, where all the values of the defect of semismallness of the Albanese map $\delta(a_X)$ are taken in consideration (see (2.1)). Theorem 1.1 is the case $\delta(a_X) = 0$. First of all, we note that the limits (1.4) are peculiar to the case $\delta(a_X) = 0$, and they have been essentially proved in Proposition 4.1.

THEOREM 4.3. — *Let X be a smooth projective variety of complex dimension n , and let $\varphi_d: X_d \rightarrow X$ be the étale covers defined in (1.2). If the defect of semismallness of the Albanese map satisfies $\delta(a_X) \leq N$, then for any pair of integers $(p, q) \in [0, n]^2$ there exists a positive constant $B(p, q)$ such that*

$$(4.2) \quad \frac{h^{p,q}(X_d)}{\deg \varphi_d} \leq B(p, q) d^{-2(|n-p-q|-N)} \quad \text{for all } d \geq 1.$$

Conversely, if $N \geq 0$ is an integer and X is a smooth projective variety of dimension n that satisfies both $\dim \text{Alb}(X) > n$ and the bounds in (4.2) for all pairs of indexes $(p, q) \in [0, n]^2$, then the defect of semismallness satisfies $\delta(a_X) \leq N$.

Proof. — Let S_d denote the set of d -torsion points on $\text{Alb}(X)$. As in (3.4), we have for all p and q the following equalities

$$h^{p,q}(X_d) = \sum_{\alpha \in S_d} h^q(X, \Omega_X^p \otimes \alpha).$$

By Proposition 3.3, there exist positive constants $B = B(p, q)$ such that

$$\frac{h^{p,q}(X_d)}{\deg \varphi_d} \leq B d^{2(\dim V^q(\Omega_X^p) - g)}$$

where $g = \dim \text{Alb}(X)$. Moreover, by Theorem 2.6, we have $\dim V^q(\Omega_X^p) \leq g - |p + q - n| + \delta(a_X)$ and

$$\frac{h^{p,q}(X_d)}{\deg \varphi_d} \leq B d^{-2(|p+q-n| - \delta(a_X))}$$

for all $d \geq 1$. This shows one implication.

Assume now that $\dim \text{Alb}(X) > n$ and that the bounds (4.2) hold. Moreover, assume by contradiction that $\delta(a_X) \geq N + 1$. By Theorem 2.6, there exists a pair $(p_0, q_0) \in [0, n]^2$ such that $\text{codim } V^{q_0}(\Omega_X^{p_0}) = |n - p_0 - q_0| - \delta(a_X)$. Then $\dim V^{q_0}(\Omega_X^{p_0}) = \dim \text{Alb}(X) - |n - p_0 - q_0| + \delta(a_X) > 0$, and by Proposition 3.4 we have

$$\frac{h^{p_0, q_0}(X_d)}{\deg \varphi_d} \geq A d^{-2(|n - p_0 - q_0| - \delta(a_X))} \quad \text{for infinitely many } d \geq 1$$

for some positive constant A independent of d (note that the loci $V^q(\Omega_X^p)$ are torsion linear by [38, Corollary 19.2]). For $d \gg 0$, this contradicts the bounds (4.2) when $(p, q) = (p_0, q_0)$. \square

Proof of Corollary 1.2. The first statement of the corollary is an application of Theorem 1.1 and (4.1). On the other hand, the second point is again Proposition 4.1. \square

5. Limits of Normalized Plurigenera

In this subsection, we apply Theorem 3.6 to the pluricanonical bundles $\omega_X^{\otimes m}$ ($m \geq 1$) of a smooth projective variety X . We set $p_g(X) = P_1(X) = h^0(X, \omega_X)$ for the geometric genus of X , and

$$P_m(X) = h^0(X, \omega_X^{\otimes m}), \quad m \geq 2,$$

for the plurigenera.

In the following proposition we fix a morphism $f: X \rightarrow A$ to an abelian variety.

PROPOSITION 5.1. — *Let X_d be the fiber product between $f: X \rightarrow A$ and μ_d as in the commutative diagram (3.1). Then for any integer $m \geq 1$ we have*

$$(5.1) \quad \lim_{d \rightarrow \infty} \frac{P_m(X_d)}{\deg \varphi_d} = \chi(f_* \omega_X^{\otimes m}).$$

Moreover, if $f: X \rightarrow A$ is generically finite onto its image, then

$$\lim_{d \rightarrow \infty} \frac{p_g(X_d)}{\deg \varphi_d} = \chi(\omega_X).$$

Proof. — By [36, Theorem 1.10] the sheaves $f_* \omega_X^{\otimes m}$ satisfy *GV* for all $m \geq 1$. Hence, by Theorem 3.6, we have

$$\lim_{d \rightarrow \infty} \frac{h^0(A, \mu_d^* f_* \omega_X^{\otimes m})}{\deg \varphi_d} = \chi(f_* \omega_X^{\otimes m}).$$

We observe that by base change, together with the fact that $h^0(A, f_{d*}\mathcal{G}) = h^0(X_d, \mathcal{G})$ for any coherent sheaf \mathcal{G} on X_d , we have the equalities

$$h^0(X_d, \omega_{X_d}^{\otimes m}) = h^0(X_d, \varphi_d^* \omega_X^{\otimes m}) = h^0(A, f_{d*} \varphi_d^* \omega_X^{\otimes m}) = h^0(A, \mu_d^* f_* \omega_X^{\otimes m}).$$

The second claimed limit follows by the Grauert–Riemenschneider Vanishing [23, Theorem 4.3.9], which yields $\chi(f_* \omega_X) = \chi(\omega_X)$ if f is generically finite onto its image. \square

For $m \geq 2$ there are two cases where one can improve the results of Proposition 5.1. The first is the case of smooth projective varieties of general type. Indeed, Kollár in [22, Proposition 9.4] shows the multiplicativity of the higher plurigenera under any étale map, so that $\frac{P_m(X_d)}{\deg \varphi_d}$ are constants and trivially

$$\lim_{d \rightarrow \infty} \frac{P_m(X_d)}{\deg \varphi_d} = P_m(X) \quad \text{for all } m \geq 2.$$

The second is the case of the Albanese map $f = a_X : X \rightarrow \text{Alb}(X)$. With a slight abuse of notation, we denote by $I : X \rightarrow Z$ a non-singular representative of the Iitaka fibration of X . Moreover, we set

$$q(I) = q(X) - q(Z) = h^0(X, \Omega_X) - h^0(Z, \Omega_Z)$$

for the difference of the irregularities.

PROPOSITION 5.2. — *Let X_d be the fiber product between a_X and μ_d as in (1.2), and fix an integer $m \geq 2$. Then there exists a positive constant M such that*

$$\frac{P_m(X_d)}{\deg \varphi_d} \leq M d^{-2q(I)} \quad \text{for all } d \geq 1.$$

Moreover we have

$$\lim_{d \rightarrow \infty} \frac{P_m(X_d)}{\deg \varphi_d} = \begin{cases} P_m(X) & \text{if } q(I) = 0 \\ 0 & \text{if } q(I) > 0. \end{cases}$$

Proof. — By [18, Theorem 11.2(b)], for each $m \geq 2$ there exist line bundles $\alpha_1, \dots, \alpha_t \in \text{Pic}^0(X)$ of finite order such that

$$V^0(\omega_X^{\otimes m}) = \bigcup_{j=1}^t (\alpha_j + \text{Pic}^0(Z)).$$

Hence we have $\dim V^0(\omega_X^{\otimes m}) = q(Z)$. Moreover, as

$$P_m(X_d) = \sum_{\alpha \in S_d} h^0(X, \omega_X^{\otimes m} \otimes \alpha)$$

(where S_d is the set of d -torsion points on $\text{Alb}(X)$), by Proposition 3.3 there exists a positive constant $M > 0$ such that the first claim holds. This also shows that the limit $\lim_{d \rightarrow \infty} \frac{P_m(X_d)}{\deg \varphi_d}$ vanishes for $q(I) > 0$. In order to complete the proof, thanks to Proposition 5.1, we only need to calculate the Euler characteristic $\chi(a_{X*}\omega_X^{\otimes m})$. As $a_{X*}\omega_X^{\otimes m}$ satisfies *GV*, by Lemma 2.3 we find that

$$(5.2) \quad \chi(a_{X*}\omega_X^{\otimes m}) = h^0(\text{Alb}(X), a_{X*}\omega_X^{\otimes m} \otimes \alpha) = h^0(X, \omega_X^{\otimes m} \otimes \alpha)$$

for a generic element $\alpha \in \text{Pic}^0(\text{Alb}(X)) \simeq \text{Pic}^0(X)$. However, if $q(I) = 0$, then by [18, Theorem 11.2(a)] we have $V^0(\omega_X^{\otimes m}) = \text{Pic}^0(X)$ and moreover the quantities $h^0(X, \omega_X^{\otimes m} \otimes \alpha)$ are independent of α . \square

Remark 5.3. — Smooth projective varieties of general type fall within the class $q(I) = 0$. Instances of varieties with $q(I) = 0$, but which are not of general type, are provided by non-isotrivial elliptic surfaces fibered over smooth projective curves Σ_g of genus $g \geq 2$. Indeed, given an elliptic surface $p: X \rightarrow \Sigma_g$, one can show that the corresponding morphism $P: \text{Alb}(X) \rightarrow \text{Alb}(\Sigma_g)$ is an isomorphism if and only if the elliptic fibration is *not* isotrivial. For more details, we refer to [4, Chapter IX]. Higher-dimensional examples may be constructed in the same fashion.

In analogy to Kollár's result [22, Proposition 9.4], the previous proposition suggests that the higher plurigenera ought to be multiplicative under étale morphisms also in the more general case $q(I) = 0$. We confirm this expectation for the étale covers induced via base change by the isogenies of $\text{Alb}(X)$. In [29, Corollary 12.2], the reader may notice a further property that shows how varieties with $q(I) = 0$ behave like varieties of general type. Indeed, the sheaves $a_{X*}\omega_X^{\otimes m}$ satisfy *I.T.* with index 0 for all $m \geq 2$ as soon as $q(I) = 0$ (cf. Remark 3.7).

THEOREM 5.4. — *Let X be a smooth projective variety with $q(I) = 0$, and let Y be the fiber product between a_X and an isogeny $\mu: B \rightarrow \text{Alb}(X)$, as in the following cartesian diagram:*

$$\begin{array}{ccc} Y & \xrightarrow{\tilde{a}} & B \\ \downarrow \varphi & & \downarrow \mu \\ X & \xrightarrow{a_X} & \text{Alb}(X). \end{array}$$

Then for all $d \geq 1$ and $m \geq 2$ we have $P_m(Y) = (\deg \varphi)P_m(X)$.

Proof. — The proof follows the general strategy of [18, Theorem 11.2] and [24, Theorem 11.2.23]. This goes as follows. As in the proof of Proposition 5.2, the Iitaka fibration I induces a surjective morphism $a_I: \text{Alb}(X) \rightarrow \text{Alb}(Z)$ with connected fibers such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{a_X} & \text{Alb}(X) \\ \downarrow I & & \downarrow a_I \\ Z & \xrightarrow{a_Z} & \text{Alb}(Z) \end{array}$$

commutes (cf. [17, Proposition 2.1]). Therefore a_I is an isomorphism as $q(I) = 0$. Fix now an integer $m \geq 2$ and let $\mathcal{I}(\|\omega_X^{\otimes(m-1)}\|)$ be the asymptotic multiplier ideal sheaf as defined in [24, Definition 11.1.2]. Moreover set $g = a_I \circ a_X$ and define the sheaf

$$\mathcal{H} = g_* \left(\omega_X^{\otimes m} \otimes \mathcal{I}(\|\omega_X^{\otimes(m-1)}\|) \right).$$

Since g factors through I , we have a linear equivalence relation $tK_X \sim g^*H + E$ where H is an ample divisor on $\text{Alb}(Z)$, E is an effective divisor, and $t \gg 0$ is a sufficiently large integer. This implies, as proved in the course of the proof of [18, Theorem 11.2], that

$$H^0(\text{Alb}(Z), \mathcal{H}) = H^0(X, \omega_X^{\otimes m}) \quad \text{and} \quad H^i(\text{Alb}(Z), \mathcal{H}) = 0 \text{ for all } i > 0.$$

Thus we have $P_m(X) = \chi(\mathcal{H})$.

Now, consider the Stein factorization $s: Y \rightarrow S$ of the composition $I \circ \varphi: Y \rightarrow Z$. As the general fiber of s has Kodaira dimension equal to zero, by [23, Remark 2.1.35] s factors through a non-singular representative of the Iitaka fibration of Y , which, with a slight abuse of notation, we denote it by $I_Y: Y \rightarrow W$. Define $\tilde{g} = a_I \circ \mu \circ \tilde{a}$. As \tilde{g} factors through I_Y , we can write

$$(5.3) \quad \tilde{t}K_Y \sim \tilde{g}^* \tilde{H} + \tilde{E}$$

for some ample line bundle \tilde{H} on $\text{Alb}(Z)$, effective divisor \tilde{E} on Y , and large integer $\tilde{t} \gg 0$. By defining the sheaves

$$\tilde{\mathcal{G}} = \tilde{g}_* \left(\omega_Y^{\otimes m} \otimes \mathcal{I}(\|\omega_Y^{\otimes(m-1)}\|) \right), \quad \tilde{\mathcal{H}} = \tilde{a}_* \left(\omega_Y^{\otimes m} \otimes \mathcal{I}(\|\omega_Y^{\otimes(m-1)}\|) \right),$$

the relation (5.3) ensures that

$$(5.4) \quad H^0(\text{Alb}(Z), \tilde{\mathcal{G}}) = H^0(Y, \omega_Y^{\otimes m}) \quad \text{and} \quad H^i(\text{Alb}(Z), \tilde{\mathcal{G}}) = 0 \text{ for } i > 0,$$

again as shown in the argument of the proof of [18, Theorem 11.2].⁽¹⁾ We conclude that $P_m(Y) = \chi(\tilde{\mathcal{G}}) = \chi(\tilde{\mathcal{H}})$ as in addition there are isomorphisms $H^i(B, \tilde{\mathcal{H}}) \simeq H^i(\text{Alb}(Z), \tilde{\mathcal{G}})$ for all $i \geq 0$ (recall that μ is étale). Finally, we note that $\chi(\tilde{\mathcal{H}}) = \chi(\mu^*(a_I^{-1})_* \mathcal{H})$. Indeed, by base change and [24, Theorem 11.2.16], we obtain the following isomorphisms

$$\tilde{\mathcal{H}} \simeq \tilde{a}_{*} \varphi^* \left(\omega_X^{\otimes m} \otimes \mathcal{I}(\|\omega_X^{\otimes(m-1)}\|) \right) \simeq \mu^*(a_I^{-1})_* \mathcal{H}.$$

To conclude, we note that

$$\chi\left(\mu^*(a_I^{-1})_* \mathcal{H}\right) = (\deg \mu) \chi\left((a_I^{-1})_* \mathcal{H}\right) = (\deg \mu) \chi(\mathcal{H})$$

as $a_I: \text{Alb}(X) \rightarrow \text{Alb}(Z)$ is an isomorphism. \square

Remark 5.5 (Higher direct images and multiplier ideal sheaves). — One can apply Theorem 3.6 to other classes of sheaves that satisfy the generic vanishing condition of Definition 2.1. In this direction, the paper [33] contains several examples of GV -sheaves. As an example, by keeping the notation of (3.1), Theorem 3.6 and [33, Theorem 5.8] give

$$\lim_{d \rightarrow \infty} \frac{h^0(A, R^i f_{d*} \omega_{X_d})}{\deg \mu_d} = \chi(R^i f_* \omega_X) \quad \text{for any } i \geq 0.$$

Moreover, Theorem 3.6 in combination with [33, Corollary 5.2] give the following statement. Suppose that the Albanese map $a_X: X \rightarrow \text{Alb}(X)$ is generically finite onto its image, and let L be a line bundle with non-negative Kodaira dimension. With notation as in (1.2), we have

$$\lim_{d \rightarrow \infty} \frac{h^0(X_d, \omega_{X_d} \otimes L_d \otimes \mathcal{I}(\|L_d\|))}{\deg \varphi_d} = \chi(\omega_X \otimes L \otimes \mathcal{I}(\|L\|)).$$

6. Applications to L^2 -Cohomology

In order to define L^2 -Betti numbers we follow the reference [31]. Let G be a discrete group, and let M be a co-compact free proper G -manifold without boundary endowed with a G -invariant Riemannian metric. Define the space of smooth L^2 -integrable harmonic k -forms

$$\mathcal{H}_{(2)}^k(M) = \left\{ \omega \in \Omega^k(M) \mid \Delta_d \omega = 0, \int_M \omega \wedge * \omega < \infty \right\}$$

⁽¹⁾Even if not explicitly stated, the proof of [18, Theorem 11.2] actually proves the isomorphism and vanishings in (5.4) for any morphism from Y to an abelian variety that factors through the Iitaka fibration of Y .

where $*$ is the Hodge star operator and $\Delta_d = dd^* + d^*d$ is the Hodge-Laplacian operator. By [31, Section 1.3.2], the spaces $\mathcal{H}_{(2)}^k(M)$ are finitely generated Hilbert modules over the von Neumann algebra $\mathcal{N}(G)$ of G . We define the L^2 -Betti numbers $b_k^{(2)}(M; \mathcal{N}(G))$ of (M, G) as the von Neumann dimension of the $\mathcal{N}(G)$ -modules $\mathcal{H}_{(2)}^k(M)$:

$$b_k^{(2)}(M; \mathcal{N}(G)) \stackrel{\text{def}}{=} \dim_{\mathcal{N}(G)} \mathcal{H}_{(2)}^k(M).$$

The L^2 -Betti numbers assume values in the extended interval $[0, \infty]$ of the real numbers, and $b_k^{(2)}(M, \mathcal{N}(G)) \in [0, \infty)$ if the action of G is co-compact.

Finally, in order to define the L^2 -Hodge numbers $h_{p,q}^{(2)}(M; \mathcal{N}(G))$ of (M, G) , we define

$$\mathcal{H}_{(2)}^{p,q}(M) = \left\{ \omega \in \Omega^{p,q}(M) \mid \Delta_{\bar{\partial}} \omega = 0, \int_M \omega \wedge * \omega < \infty \right\},$$

where $\Delta_{\bar{\partial}} = \bar{\partial} \bar{\partial}^* + \bar{\partial}^* \bar{\partial}$ is the $\bar{\partial}$ -Laplacian, and set

$$h_{p,q}^{(2)}(M; \mathcal{N}(G)) \stackrel{\text{def}}{=} \dim_{\mathcal{N}(G)} \mathcal{H}_{(2)}^{p,q}(M).$$

By [31, Chapter 11], there is a L^2 -Hodge decomposition which gives

$$(6.1) \quad b_k^{(2)}(M; \mathcal{N}(G)) = \sum_{p+q=k} h_{p,q}^{(2)}(M; \mathcal{N}(G)).$$

6.1. (Non-)Vanishing of L^2 -Betti numbers

Let X be a smooth projective variety of dimension n , and let $a_X: X \rightarrow \text{Alb}(X)$ be the Albanese map. Moreover set $g = \dim \text{Alb}(X)$. The *universal Albanese cover* $\bar{\pi}: \bar{X} \rightarrow X$ is defined as the pullback of a_X via the topological universal cover $\mathbb{C}^g \rightarrow \text{Alb}(X)$ (cf. [11, Section 3.2]). We set

$$\Gamma = \pi_1(X), \quad \bar{\Gamma} = \pi_1(\bar{X}), \quad G = \Gamma/\bar{\Gamma} \quad \text{and} \quad A = \text{Alb}(X).$$

THEOREM 6.1. — *If X satisfies the weak generic Nakano vanishing theorem, then the L^2 -Betti numbers of \bar{X} are*

$$b_k^{(2)}(\bar{X}; \mathcal{N}(G)) = \begin{cases} (-1)^n \chi_{\text{top}}(X) & \text{if } k = n \\ 0 & \text{if } k \neq n. \end{cases}$$

In particular, we have $\mathcal{H}_{(2)}^{p,q}(\bar{X}) = 0$ if $p + q \neq n$.

Proof. — Consider the following cartesian diagram induced inductively by the multiplication maps μ_d via the base change:

$$\begin{array}{ccccccccccc}
 \cdots & \longrightarrow & Y_{d+1} & \longrightarrow & Y_d & \longrightarrow & Y_{d-1} & \longrightarrow & \cdots & \longrightarrow & Y_2 & \longrightarrow & X \\
 & & \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow a_X \\
 \cdots & \longrightarrow & A & \xrightarrow{\mu_{d+1}} & A & \xrightarrow{\mu_d} & A & \xrightarrow{\mu_{d-1}} & \cdots & \longrightarrow & A & \xrightarrow{\mu_2} & A.
 \end{array}$$

By using the notation of the commutative diagram (3.1), we immediately realize that $Y_d \simeq X_d$. We set $\Gamma_d = \pi_1(Y_d)$. By [11, Section 3.1], together with the proof of [11, Lemma 3.2], the homomorphism $a_{X\#} : \pi_1(X) \rightarrow \pi_1(A)$ is surjective and the varieties Y_d satisfy

$$\ker(a_{X\#} : \pi_1(X) \rightarrow \pi_1(A)) = \bigcap_{d=1}^{\infty} \Gamma_d$$

(i.e., in the terminology of [11, Lemma 3.2], the isogenies r_i can be chosen as the multiplication maps μ_d). Moreover, the universal Albanese cover $\bar{\pi} : \bar{X} \rightarrow X$ is identified to the regular cover associated to the normal separable subgroup $\ker(a_{X\#})$. Therefore $\bar{\Gamma} = \ker(a_{X\#})$ and there are isomorphisms

$$Y_d \simeq \bar{X} / G_d \quad \text{where} \quad G_d \stackrel{\text{def}}{=} \Gamma_d / \bar{\Gamma}.$$

As the sequence $\{G_d\}_{d \geq 1}$ is an inverse system of normal subgroups such that $\bigcap_{d \geq 1} G_d = \{1\}$, Lück’s Approximation Theorem [31, Theorem 13.3] and [31, Example 1.32] yield

$$b_k^{(2)}(\bar{X}; \mathcal{N}(G)) = \lim_{d \rightarrow \infty} b_k^{(2)}(Y_d; \mathcal{N}(G/G_d)) = \lim_{d \rightarrow \infty} \frac{b_k(Y_d)}{[\Gamma : \Gamma_d]} = \lim_{d \rightarrow \infty} \frac{b_k(X_d)}{\deg \mu_d}.$$

At this point, the first statement of the theorem is an application of Proposition 4.1. On the other hand, the second follows by the fact that the von Neumann dimension of a Hilbert module is zero if and only if the module itself is trivial (see [31, Theorem 1.12(1)]), and the L^2 -Hodge decomposition (6.1). □

The following non-vanishing result was proved by Gromov in the case of topological universal covers of Kähler hyperbolic manifolds (cf. [16, Section 2] and [31, Theorem 11.35]).

COROLLARY 6.2. — *Let X be as in Theorem 6.1. If $\chi(\Omega_X^p) \neq 0$, then we have*

$$\mathcal{H}_{(2)}^{p, n-p}(\bar{X}) \neq 0.$$

Moreover, if $\chi(\omega_X) \neq 0$, then there exists a nontrivial holomorphic L^2 -integrable n -form on the universal Albanese cover \bar{X} .

Proof. — We use the notation of the previous proof. First of all we observe that by Proposition 4.1 we have the inequalities

$$\limsup_{d \rightarrow \infty} \frac{h^{p,n-p}(Y_d)}{\deg \mu_d!} \geq \lim_{d \rightarrow \infty} \frac{h^{p,n-p}(X_{d!})}{\deg \mu_d!} = (-1)^{n-p} \chi(\Omega_X^p) > 0$$

Thus the result follows by Kazhdan’s inequality [21, Theorem 2] (cf. also [20, p. 6–7]):

$$h_{p,n-p}^{(2)}(\bar{X}, \mathcal{N}(G)) \geq \limsup_{d \rightarrow \infty} \frac{h^{p,n-p}(Y_d)}{\deg \mu_d!}.$$

The second statement is proved as in [31, Corollary 11.36]. In other words, if we have a non-zero form $\omega \in \mathcal{H}_{(2)}^{n,0}(\bar{X})$, then $\Delta_{\bar{\partial}}\omega = 0$ and $\bar{\partial}\omega = 0$. This means that ω is holomorphic. \square

Appendix A. Coverings of Varieties with Unbounded Irregularity

Let Y be a smooth projective variety satisfying $q(Y) = h^{1,0}(Y) = \dim \text{Alb}(Y) > 0$ and $H^2(Y, \mathbb{Z})_{\text{tor}} = 0$. We provide sufficient and necessary conditions for the irregularities $Q = \{q(Y_i)\}_{i=1}^{\infty}$ of a series of coverings $\pi_i: Y_i \rightarrow Y$ induced by the multiplication maps on $\text{Alb}(Y)$ to diverge as $\deg \pi_i \rightarrow \infty$. This problem has been already addressed in the literature. For instance, by the recent work of Vidussi [41, Lemma 1.3] and Stover [40, Theorem 3], the irregularity of any unramified abelian cover of the Cartwright–Steger surface⁽²⁾ is equal to one. On the other hand, it is very easy to construct towers of coverings with unbounded irregularities.

Turning to details, let X be a smooth projective variety of dimension n and $a_X: X \rightarrow \text{Alb}(X)$ be the Albanese map. The multiplication maps $\mu_d: \text{Alb}(X) \rightarrow \text{Alb}(X)$, $\mu_d(x) = dx$ induce via base-change unramified covers $a_d: X_d \rightarrow X$. We use the term *fibration* to mean a surjective morphism of varieties with connected fibers. The following result builds upon [3, Corollaire 2.3].

THEOREM A.1. — *Suppose that $\limsup_{d \rightarrow \infty} q(X_d) = \infty$. Then X admits a fibration $p: X \rightarrow C$ onto a smooth curve of genus g such that either $g \geq 2$, or $g = 1$ and the fibration admits two multiple fibers whose multiplicities are not coprime. If in addition $H^2(X, \mathbb{Z})_{\text{tor}} = 0$, then the converse holds.*

⁽²⁾The Cartwright–Steger surface S is a complex hyperbolic surface with *minimal* Euler characteristic $\chi(S) = 3$, and non-trivial first Betti number. It was computationally discovered in [8] during the classification of fake projective planes. We refer to [7] for an in depth study of its geometry.

Proof. — Let S_d be the set of d -torsion points of $\text{Pic}^0(X)$. The irregularity $q(X_d) = h^1(X_d, \mathcal{O}_{X_d})$ can be computed with the techniques of Theorem 3.6:

$$(A.1) \quad q(X_d) = q(X) + \sum_{\alpha \in S_d, \alpha \neq \mathcal{O}_X} h^1(X, \alpha).$$

First of all we prove that $\limsup_{d \rightarrow \infty} q(X_d) = \infty$ if and only if there exists a positive-dimensional component of the Green–Lazarsfeld locus

$$V^{n-1}(\omega_X) \simeq V^1(\mathcal{O}_X) \stackrel{\text{def}}{=} \{\alpha \in \text{Pic}^0(X) \mid h^1(X, \alpha) > 0\}.$$

In fact, if $\limsup_{d \rightarrow \infty} q(X_d) = \infty$, then by (A.1) there are infinitely many distinct elements of $V^1(\mathcal{O}_X)$. As $V^1(\mathcal{O}_X)$ is an algebraic variety, these elements must form one irreducible component. On the other hand, if $v_1 = \dim V^1(\mathcal{O}_X) > 0$, then, by Proposition 3.4, $V^1(\mathcal{O}_X)$ ⁽³⁾ contains at least d^{2v_1} d -torsion points for infinitely many $d \geq 1$. Hence $q(X_d) \geq q(X) + d^{2v_1} - 1$ and the claim follows.

Let now $\text{Pic}^\tau(X)$ be the variety that parameterizes isomorphism classes of holomorphic line bundles on X with torsion first Chern class. By the work of [14, Theorem 0.1] and [3, Corollaire 2.3], the irreducible components of $V^{n-1}(\omega_X)$ are related to fibrations over smooth projective curves. More precisely, any positive-dimensional irreducible component $S \subset V^{n-1}(\omega_X)$ is a component of the group

$$\text{Pic}^\tau(X, p) \stackrel{\text{def}}{=} \ker(i^*: \text{Pic}^\tau(X) \rightarrow \text{Pic}^\tau(F))$$

for some fibration $p: X \rightarrow C$ over a smooth projective curve of genus $g \geq 1$ with general fiber $i: F \hookrightarrow X$. It follows that $\dim S = g$. Moreover, if $g = 1$, then by [3, Corollaire 2.3] we have $S \neq p^* \text{Pic}^0(C)$. Therefore, by [3, Remarque 2.4], p must possess at least two multiple fibers whose multiplicities are not coprime (cf. also [38, Exercise 10.3]). This proves one of the implications.

Let now $p: X \rightarrow C$ be a fibration onto a smooth projective curve of genus $g \geq 1$. For the other direction, we note that if $g \geq 2$, then, by pulling-back line bundles from $\text{Pic}^0(C) = V^0(\omega_C)$, the fibration p gives rise to an irreducible component of $V^{n-1}(\omega_X)$ (cf. [28, Lemma 6.3]). We now prove that we reach the same conclusion even if $g = 1$, $H^2(X, \mathbb{Z})_{\text{tor}} = 0$, and the condition on the multiple fibers is verified. In fact, the condition on the multiple fibers implies that the group $\Gamma^\tau(p) \simeq \text{Pic}^\tau(X, p)/p^* \text{Pic}^0(C)$ of the connected components of $\text{Pic}^\tau(X, p)$ is non-trivial (cf. [3, Proposition 1.5,

⁽³⁾By [38, Corollary 19.2], the locus $V^1(\mathcal{O}_X)$ is a finite union of torsion translates of abelian subvarieties of $\text{Pic}^0(X)$.

Remarque 2.4] and [38, Exercise 10.3]). By [3, Section 1.6], as $H^2(X, \mathbb{Z})_{\text{tor}} = 0$, the group $\Gamma^\tau(p)$ is identified with the group $\Gamma^0(p)$ of the connected components of the following group

$$\text{Pic}^0(X, p) \stackrel{\text{def}}{=} \text{Pic}^\tau(X, p) \cap \text{Pic}^0(X).$$

Therefore $\text{Pic}^0(X, p)$ contains a connected component different from the neutral component $p^* \text{Pic}^0(C)$, which, again by [3, Corollaire 2.3], it is contained in $V^{n-1}(\omega_X)$. \square

Remark A.2. — Let S be the Cartwright–Steger surface. It follows from [8] that $H_1(S, \mathbb{Z})$ is torsion free. By the universal coefficient theorem, we know that $H^2(S, \mathbb{Z})_{\text{tor}} = 0$. Moreover, the Albanese map has no multiple fibers (cf. [7, Main Theorem]). Then Theorem A.1 implies that the unramified abelian covers of S have bounded irregularities (however much more is true for the surface S , cf. again [41] and [40] for optimal statements).

The argument of Proposition A.1 extends, in a weaker form, to all Hodge numbers of type $h^{n,i}(X)$ with $i > 0$.

PROPOSITION A.3. — *If $\limsup_{d \rightarrow \infty} h^{n,i}(X_d) = \infty$, then there exists a fibration of X onto a normal projective variety Y of dimension $0 < \dim Y \leq n - i$ such that any smooth model of Y is of maximal Albanese dimension.*

Proof. — Thanks to a calculation similar to (A.1), we can construct an irreducible component $S \subset V^i(\omega_X)$ of positive dimension. By [14, Theorem 0.1], this component induces a fibration of X onto a variety with the desired properties. \square

BIBLIOGRAPHY

- [1] M. ABERT, N. BERGERON, I. BIRINGER & T. GELANDER, “Convergence of normalized Betti numbers in nonpositive curvature”, <https://arxiv.org/abs/1811.02520>, 2018.
- [2] M. ABERT, N. BERGERON, I. BIRINGER, T. GELANDER, N. NIKOLOV, J. RAIMBAULT & I. SAMET, “On the growth of L^2 -invariants for sequences of lattices in Lie groups”, *Ann. Math.* **185** (2017), no. 3, p. 711-790.
- [3] A. BEAUVILLE, “Annulation du H^1 pour les fibrés en droites plats”, in *Complex algebraic varieties (Bayreuth, 1990)*, Lecture Notes in Mathematics, vol. 1507, Springer, 1992, p. 1-15.
- [4] ———, *Complex algebraic surfaces*, 2nd ed., London Mathematical Society Student Texts, vol. 34, Cambridge University Press, 1996, translated from the 1978 French original by R. Barlow, with assistance from N. I. Shepherd-Barron and M. Reid, x+132 pages.
- [5] A. BONDAL & D. ORLOV, “Semiorthogonal decomposition for algebraic varieties”, <https://arxiv.org/abs/alg-geom/9506012>, 1995.

- [6] N. BUDUR, “Unitary local systems, multiplier ideals, and polynomial periodicity of Hodge numbers”, *Adv. Math.* **221** (2009), no. 1, p. 217-250.
- [7] D. I. CARTWRIGHT, V. KOZIARZ & S.-K. YEUNG, “On the Cartwright-Steger surface”, *J. Algebr. Geom.* **26** (2017), no. 4, p. 655-689.
- [8] D. I. CARTWRIGHT & T. STEGER, “Enumeration of the 50 fake projective planes”, *C. R. Math. Acad. Sci. Paris* **348** (2010), no. 1-2, p. 11-13.
- [9] D. L. DEGEORGE & N. R. WALLACH, “Limit formulas for multiplicities in $L^2(\Gamma \backslash G)$ ”, *Ann. Math.* **107** (1978), no. 1, p. 133-150.
- [10] ———, “Limit formulas for multiplicities in $L^2(\Gamma \backslash G)$. II. The tempered spectrum”, *Ann. Math.* **109** (1979), no. 3, p. 477-495.
- [11] L. F. DI CERBO & L. LOMBARDI, “Moving Seshadri constants, and coverings of varieties of maximal Albanese dimension”, *Asian J. Math.* **25** (2021), no. 2, p. 305-320.
- [12] L. F. DI CERBO & M. STERN, “Price inequalities and Betti number growth on manifolds without conjugate points”, to appear in *Commun. Anal. Geom.*, <https://arxiv.org/abs/1704.06354>, 2017.
- [13] M. GREEN & R. LAZARSFELD, “Deformation theory, generic vanishing theorems, and some conjectures of Enriques, Catanese and Beauville”, *Invent. Math.* **90** (1987), no. 2, p. 389-407.
- [14] ———, “Higher obstructions to deforming cohomology groups of line bundles”, *J. Am. Math. Soc.* **4** (1991), no. 1, p. 87-103.
- [15] N. GRIEVE, “Index conditions and cup-product maps on Abelian varieties”, *Int. J. Math.* **25** (2014), no. 4, article no. 1450036 (31 pages).
- [16] M. GROMOV, “Kähler hyperbolicity and L_2 -Hodge theory”, *J. Differ. Geom.* **33** (1991), no. 1, p. 263-292.
- [17] C. HACON & R. PARDINI, “On the birational geometry of varieties of maximal Albanese dimension”, *J. Reine Angew. Math.* **546** (2002), p. 177-199.
- [18] C. HACON, M. POPA & C. SCHNELL, “Algebraic fiber spaces over abelian varieties: around a recent theorem by Cao and Păun”, in *Local and global methods in algebraic geometry*, Contemporary Mathematics, vol. 712, American Mathematical Society, 2018, p. 143-195.
- [19] R. HARTSHORNE, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 52, Springer, 1977, xvi+496 pages.
- [20] J. JOST & K. ZUO, “Vanishing theorems for L^2 -cohomology on infinite coverings of compact Kähler manifolds and applications in algebraic geometry”, *Commun. Anal. Geom.* **8** (2000), no. 1, p. 1-30.
- [21] D. A. KAJDAN, “On arithmetic varieties”, in *Lie groups and their representations (Proc. Summer School, Bolyai János Math. Soc., Budapest, 1971)*, 1975, p. 151-217.
- [22] J. KOLLÁR, “Shafarevich maps and plurigenera of algebraic varieties”, *Invent. Math.* **113** (1993), no. 1, p. 177-215.
- [23] R. LAZARSFELD, *Positivity in algebraic geometry I. Classical setting: line bundles and linear series*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*, vol. 48, Springer, 2004, xviii+387 pages.
- [24] ———, *Positivity in algebraic geometry II. Positivity for vector bundles, and multiplier ideals*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*, vol. 49, Springer, 2004, xviii+385 pages.
- [25] Y. LIU, L. MAXIM & B. WANG, “Generic vanishing for semi-abelian varieties and integral Alexander modules”, *Math. Z.* **293** (2019), no. 1-2, p. 629-645.
- [26] ———, “Topology of subvarieties of complex semi-abelian varieties”, *Int. Math. Res. Not.* (2021), no. 14, p. 11169-11208.

- [27] L. LOMBARDI, “Inequalities for the Hodge numbers of irregular compact Kähler manifolds”, *Int. Math. Res. Not.* (2013), no. 1, p. 63-83.
- [28] ———, “Derived invariants of irregular varieties and Hochschild homology”, *Algebra Number Theory* **8** (2014), no. 3, p. 513-542.
- [29] L. LOMBARDI, M. POPA & C. SCHNELL, “Pushforwards of pluricanonical bundles under morphisms to abelian varieties”, *J. Eur. Math. Soc.* **22** (2020), no. 8, p. 2511-2536.
- [30] W. LÜCK, “Approximating L^2 -invariants by their finite-dimensional analogues”, *Geom. Funct. Anal.* **4** (1994), no. 4, p. 455-481.
- [31] ———, *L^2 -invariants: theory and applications to geometry and K -theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, vol. 44, Springer, 2002, xvi+595 pages.
- [32] D. MUMFORD, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics, vol. 5, Tata Institute of Fundamental Research; Hindustan Book Agency, 2008, with appendices by C. P. Ramanujam and Yuri Manin, Corrected reprint of the second (1974) edition, xii+263 pages.
- [33] G. PARESCHI & M. POPA, “GV-sheaves, Fourier-Mukai transform, and generic vanishing”, *Am. J. Math.* **133** (2011), no. 1, p. 235-271.
- [34] M. POPA & C. SCHNELL, “Generic vanishing theory via mixed Hodge modules”, *Forum Math. Sigma* **1** (2013), article no. e1 (60 pages).
- [35] ———, “Kodaira dimension and zeros of holomorphic one-forms”, *Ann. Math.* **179** (2014), no. 3, p. 1109-1120.
- [36] ———, “On direct images of pluricanonical bundles”, *Algebra Number Theory* **8** (2014), no. 9, p. 2273-2295.
- [37] M. RAYNAUD, “Sous-variétés d’une variété abélienne et points de torsion”, in *Arithmetic and geometry, Vol. I*, Progress in Mathematics, vol. 35, Birkhäuser, 1983, p. 327-352.
- [38] C. SCHNELL, “Notes on generic vanishing”, available on the author’s webpage, 2013.
- [39] C. SIMPSON, “Subspaces of moduli spaces of rank one local systems”, *Ann. Sci. Éc. Norm. Supér.* **26** (1993), no. 3, p. 361-401.
- [40] M. STOVER, “On general type surfaces with $q = 1$ and $c_2 = 3p_g$ ”, *Manuscr. Math.* **159** (2019), no. 1-2, p. 171-182.
- [41] S. VIDUSSI, “The slope of surfaces with Albanese dimension one”, *Math. Proc. Camb. Philos. Soc.* **167** (2019), no. 2, p. 355-360.
- [42] S.-K. YEUNG, “Betti numbers on a tower of coverings”, *Duke Math. J.* **73** (1994), no. 1, p. 201-226.
- [43] T. ZHANG, “Severi inequality for varieties of maximal Albanese dimension”, *Math. Ann.* **359** (2014), no. 3-4, p. 1097-1114.

Manuscrit reçu le 26 juin 2020,
révisé le 9 décembre 2021,
accepté le 17 février 2022.

Luca F. DI CERBO
University of Florida (USA)
ldicerbo@ufl.edu
Luigi LOMBARDI
University of Milan (Italy)
luigi.lombardi@unimi.it